

and Noyes<sup>3</sup> have demonstrated a similar reaction. Extensions of these investigations are being continued.

(3) G. H. Coleman and W. A. Noyes, *THIS JOURNAL*, **43**, 2211 (1921).

CHEMICAL RESEARCH DEPT. M. E. BROOKS  
DAVISON CHEMICAL COMPANY B. RUDNER  
BALTIMORE 3, MARYLAND

RECEIVED APRIL 12, 1956

### CRYSTAL STRUCTURE AND MAGNETIC SUSCEPTIBILITY OF AMERICIUM METAL<sup>1</sup>

Sir:

We have obtained interpretable X-ray diffraction patterns of several small polycrystalline samples of americium metal of >99% purity, using Cu K $\alpha$  radiation and a 4.5-cm. radius camera. The metal was prepared on a micro scale by reduction of the trifluoride with barium vapor, in a tantalum crucible system and subsequently was annealed by slowly reducing the temperature from 800 to ca. 25° over a period of ten hours.

The powder patterns have been indexed as double hexagonal close packed,  $a = 3.642 \pm 0.005$  Å.,  $c = 11.76 \pm 0.01$  Å.

The space group is D<sub>6h</sub><sup>4</sup> and the atomic positions are: two Am in (0, 0, 0), (0, 0, 1/2); two Am in (1/3, 2/3, 3/4), (2/3, 1/3, 1/4). The Am radius is 1.82 Å. and the calculated density  $11.87 \pm 0.05$  g. cm.<sup>-3</sup>.

Relative line intensities calculated for the proposed structure agreed with visual estimates of the intensities seen in the diffraction patterns, as shown in the accompanying table.

The density calculated for the metal agrees with that observed experimentally<sup>2</sup> ( $11.7 \pm 0.3$ ) within the error of the measurements. The metallic radius is 0.02 Å. smaller than that predicted by Zachariasen<sup>3</sup> for americium metal with three valence electrons per atom. This discrepancy may indicate a small error in the predicted value, or may be due to a slight admixture of americium (IV) in the metallic state. Measurements of the magnetic susceptibility of our samples gave  $\chi_M = 1000 \pm 250 \times 10^{-6}$  cgs. units at 300° K., similar to the value of  $\sim 1000 \times 10^{-6}$  c.g.s. units for AmF<sub>3</sub>. The number of bonding electrons per atom appears to be quite close to three.

The decrease in the number of metallic bonds in going from uranium to americium affords a reasonable explanation of the corresponding decrease of some 50 kcal.<sup>4,5</sup> in the heat of vaporization.

It is interesting to note that americium is the first transactinium element which is rare earth-like in the metallic state.

Possible allotropy of the metal is now under investigation, and these studies, as well as a detailed description of the work outlined above, will be reported in a future publication.

(1) This work was performed under the auspices of the AEC.

(2) E. F. Westrum, Jr., and L. Eyring, *THIS JOURNAL*, **73**, 3396 (1951).

(3) W. H. Zachariasen, *Acta Cryst.*, **5**, 660 (1952).

(4) E. G. Rauh and R. J. Thorn, *J. Chem. Phys.*, **22**, 1414 (1954).

(5) S. C. Carniglia and B. B. Cunningham, *THIS JOURNAL*, **77**, 1502 (1955).

TABLE I

DIFFRACTION DATA FOR AMERICIUM METAL<sup>a</sup>

$hkl$	$\sin^2\theta$ calcd.	$\sin^2\theta$ obs.	$I$ calcd.	$I$ obs. <sup>b</sup>
100	0.0597	0.0592	3	vw
101	.0640	.0640	18	m
004	.0687	.0692	14	ms
102	.0769	.0769	42	s
103	.0984	.0985	10	w
104	.1284	.1283	2	t
105	.1670	.1675	5	vw
110	.1792	.1792	11	m
106	.2142	.2144	9	m
200	.2389	....	0.4	..
201	.2432	....	3	..
114	.2479	.2471	13	ms
202	.2561	.2567	7	vw
107	.2700	.2698	2	t
008	.2747	.2755	2	vww
203	.2776			
204	.3076	....	0.6	..
108	.3344	....	0.5	..
205	.3462	.3462	1	t
206	.3934	.3930	4	vww
109	.4074	.4070	1	t
210	.4181	....	0.4	..
211	.4224	.4226	2	t
212	.4353	.4353	0.6	t
207	.4492	....	1	..
118	.4539	.4531	5	m
213	.4567			
214	.4867	.4878	0.6	vw
1,0,10	.4889			

<sup>a</sup> This list includes all planes up to  $\sin^2\theta = 0.5$  for which the intensity was not calculated to be zero, by the symmetry of the special positions. <sup>b</sup> t, trace; vww, very, very weak; vw, very weak; w, weak; m, moderate; ms, moderately strong; s, strong.

**Acknowledgments.**—Peter Graf wishes to express his appreciation to Professor G. T. Seaborg for making possible for him a year's stay at the Radiation Laboratory, and to the Foundation of Fellowships in the Field of Chemistry of Switzerland for the grant of a fellowship.

PETER GRAF  
DEPARTMENT OF CHEMISTRY AND RADIATION LABORATORY  
UNIVERSITY OF CALIFORNIA  
BERKELEY, CALIFORNIA  
B. B. CUNNINGHAM  
CAROL H. DAUBEN  
J. C. WALLMANN  
D. H. TEMPLETON  
HELENA RUBEN

RECEIVED FEBRUARY 27, 1956

### SYNTHESIS OF FUSED-RING COMPOUNDS VIA ACYLATION OF KETONES

Sir:

A new method for synthesis of fused-ring compounds (II) has been found, which offers certain advantages over the well-known Pschorr,<sup>1</sup> Haworth,<sup>2</sup> Bardhan-SenGupta<sup>3</sup> and Bogert<sup>4</sup> procedures. The method consists of the reaction of cyclic ketones with phenylacetic anhydrides in the presence of boron trifluoride.<sup>5</sup> Acylation and cy-

(1) R. Pschorr, *Ber.*, **29**, 496 (1896).

(2) R. D. Haworth, *J. Chem. Soc.*, 1125 (1932).

(3) J. C. Bardhan and S. C. SenGupta, *ibid.*, 2520 (1932).

(4) M. T. Bogert, *Science*, **77**, 289 (1933).

(5) C. R. Hanser, F. W. Swamer and J. T. Adams, in R. Adams, "Organic Reactions," Vol. 8, Chapter 3, pp. 59-196, J. Wiley and Sons, Inc., New York, N. Y., 1954.