

no evidence of *dl*-epiyohimbane; the remaining chromatographic fractions amounted to 67 mg. of non-crystalline, tarry residues.

Anal. Calcd. for $C_{19}H_{24}N_2$: C, 81.38; H, 8.63. Found: C, 81.71; H, 8.35.

dl-Yohimbane Hydrochloride.—A 0.07-g. portion of the dichlorophosphate salt of Δ^3 -dehydroyohimbane was dissolved in 30 ml. of ethanol and hydrogenated at room temperature and pressure, using 0.010 g. of platinum oxide. There was a rapid uptake of hydrogen which slowed down considerably after the first 0.5 hr.; the hydrogen absorbed corresponded to 1 mole. The catalyst was filtered off, and the ethanol was evaporated to give a residue of colorless crystals which was recrystallized from ethanol. The final product melted 276–277°.

Anal. Calcd. for $C_{19}H_{23}ClN_2$: C, 72.01; H, 7.95. Found: C, 71.52; H, 8.09.

In order to obtain the free base, a solution of 0.050 g. of *dl*-yohimbane hydrochloride in 2 cc. of ethanol was added to 0.015 g. of potassium hydroxide in ethanol. The whole was thoroughly shaken and allowed to stand for 15 minutes. The ethanol was then evaporated to dryness over a steam-bath and under a current of nitrogen. To the white residue was added 4 ml. of water, and again the mixture was thoroughly shaken for 15 minutes. The colorless, crystalline precipitate of yohimbane was then filtered off and thoroughly washed with excess water and then recrystallized from 2 cc. of ethanol.

The *dl*-N-(β -3'-Indolyethyl)-*trans*-decahydroisoquinoline (IIa). A. From *dl*-Decahydroisoquinoline and Indolyethyl Bromide.—A mixture of 8 g. of decahydroisoquinoline^{3d} and 13 g. of β -3-indolyethyl bromide in 50 ml. of

ethanol was refluxed for 1 hr. The mixture was then cooled in an ice-bath; the colorless precipitate of the salt which was then collected on the filter was first washed with ether and then recrystallized from ethanol and a little acetone. In this way there was obtained 14.0 g. (67%) of colorless hydrobromide crystals, m.p. 240–242°.

Anal. Calcd. for $C_{19}H_{27}BrN_2$: C, 62.80; H, 7.49. Found: C, 63.04; H, 7.57.

The free base was prepared by treating some of the above salt with aqueous sodium hydroxide. The colorless crystals of the free base melted at 150–151°, after recrystallization from ethanol.

Anal. Calcd. for $C_{19}H_{25}N_2$: C, 80.80; H, 9.28. Found: C, 81.34; H, 9.66.

B. By Reduction of the Lactam XIII.—To a solution of 0.2 g. of lactam XIII in 50 ml. of anhydrous tetrahydrofuran was added a suspension of 0.2 g. of lithium aluminum hydride in 50 ml. of tetrahydrofuran over a period of 3 hr. The mixture was then refluxed on a steam-bath for 15 minutes. After evaporation of most of the solvent under reduced pressure, the precipitate left was taken up in methanol and filtered. Evaporation of the filtrate over a steam-bath gave yellowish crystals, m.p. 277–279°. This material was taken up in aqueous sodium hydroxide and the resulting mixture thoroughly extracted with chloroform. Evaporation of the chloroform on a steam-bath and recrystallization of the residue from methanol gave 0.04 g. (21%) of colorless crystals, m.p. 151°, which was shown, by mixed melting point determination, to be identical with the base described in A above.

MADISON, WISCONSIN

[CONTRIBUTION FROM THE DEPARTMENT OF CHEMISTRY AND RADIATION LABORATORY, UNIVERSITY OF CALIFORNIA]

Sites of Azaserine Inhibition During Photosynthesis by *Scenedesmus*¹

By S. A. BARKER,² J. A. BASSHAM, M. CALVIN AND U. C. QUARCK

RECEIVED MAY 10, 1956

L-Azaserine has been found to have a profound effect on the reservoir sizes of many of the metabolic intermediates produced during photosynthesis by *Scenedesmus*. Marked increases in levels of glutamine and the acids of the Krebs cycle were accompanied by a corresponding depletion of the amino acid reservoirs indicating that one of the major sites of azaserine action is in reactions involving transamination. In contrast, the photosynthetic carbon cycle is virtually unaffected and the rate of formation of sucrose is increased.

The success attending the use of azaserine^{3,4} as a specific inhibitor of one stage in the metabolic pathway leading to the synthesis of inosinic acid in pigeon liver prompted us to use this antibiotic in a similar attack on purine synthesis in *Scenedesmus*. However, investigation of the products resulting during photosynthesis by suspensions of these algae in the presence of azaserine showed that a more widespread interference with metabolism had occurred. The purpose of this communication is to describe the nature of these effects and to attempt to assess their importance in a general picture of the metabolic effects of azaserine.

Experimental

Experimental Procedure.—Two suspensions, each containing washed *Scenedesmus* cells (packed volume, 0.2 cc.) and KH_2PO_4 solution (0.4 cc., $3.2 \times 10^{-6} M$) in 21 cc. and one with added L-azaserine (4 mg.), were left for one hour in thin glass containers illuminated on each side by a 150 w. light (reflector flood) to achieve steady states with 4% CO_2

in air. Each suspension was then allowed to photosynthesize for five minutes with $NaH^{14}O_3$ solution (0.9 cc., 360 $\mu c.$) and then flushed with air for 1 minute. The cells were then killed by pouring into boiling ethanol (88 cc.) and the resulting 80% ethanol extract was separated from insoluble material which was then reextracted with 20% ethanol (100 cc.). The total fixation of radioactivity was determined in each case by uniformly distributing and drying 50- $\mu l.$ aliquots of the 80% ethanol suspensions plus 50 $\mu l.$ of 6 N acetic acid on aluminum discs, and counting the radioactivity with a Scott large-window Geiger-Mueller tube. The radioactivity extracted from each suspension with 80% ethanol and with 20% ethanol was determined in a similar manner. After the combined extracts of each suspension were concentrated to 3–4 cc., aliquots calculated to contain 1×10^8 counts/min. each were applied to several washed Whatman No. 4 papers and separated first in phenol-water and then in butanol-propionic acid in the manner described by Wilson and Calvin.⁵ After radioautographs of the chromatograms had been made, the various components detected on the papers were counted (Table I). Since the correction for self-absorption of radiation would be the same for each compound, no correction was applied. The results of a duplicate experiment, in which 1 mg. of azaserine was used, are also presented (Table I).

Other experiments carried out in an identical manner to those described above were: (1) a repetition of the 1-mg. azaserine experiment in which an intense photospot light was substituted for one of the reflector floods for 50 minutes

(1) The work described in this paper was sponsored by the U. S. Atomic Energy Commission.

(2) Rockefeller Research Fellow, 1955–1956.

(3) S. C. Hartman, B. Levenberg and J. M. Buchanan, *THIS JOURNAL*, **77**, 501 (1955).

(4) B. Levenberg and J. M. Buchanan, *ibid.*, **78**, 504 (1956).

(5) A. T. Wilson and M. Calvin, *ibid.*, **77**, 5948 (1955).

TABLE I
THE EFFECT OF AZASERINE ON *Scenedesmus*

	Control I	Azaserine I (4 mg.)	Control II	Azaserine II (1 mg.)
Determination on aluminum disc, counts/min.				
Total fixation	22×10^6	27×10^6	29.2×10^6	33×10^6
80% ethanol extract	45.5%	47%	47.3%	47.3%
20% ethanol extract	5.9%	7.5%	7.5%	8.5%
Determination on paper, counts/min.: equal total radioactivity placed on each paper				
Glutamine	332	1192	390	1312
Glutamic acid	2066	1227	2632	1139
Aspartic acid	4436	989	7020	762
Serine	1839	849	2411	990
Alanine	1166	272	2000	314
Glycine } area	400	574	1045	908
Threonine				
Tyrosine	261	171	444	295
Valine	1051	787	1468	1053
Sucrose	923×9.5^a	2380×9.5	937	2072
Malic acid	3922×9.5	5814×9.5	2691×9.5	4263×9.5
Citric acid	466×9.5	1118×9.5	145×9.5	400×9.5
α -Ketoglutaric acid	40	200	270	1054
Fumaric acid	269	108	381	344
Succinic acid	1117	1689	822	922
Glyceric acid	576	966	1349	845
Glycolic acid	200	3182	240	880
Lipids	1650×9.5	1259×9.5	2540×9.5	1923×9.5
Phospholipids	1229×9.5	1071×9.5	1350×9.5	853×9.5
Area X	1001×9.5	751×9.5	621×9.5	538×9.5
Sugar phosphates, PGA and origin	$14,040 \times 9.5$	$16,379 \times 9.5$	$15,071 \times 9.5$	$14,424 \times 9.5$

^a The factor 9.5 was used whenever the radioactivity was counted through aluminum foil.

of the one hour steady state period (Table II) and (2) an experiment in which one suspension contained 4 mg. of azaserine and the other both azaserine (4 mg.) and glutamine (5 mg.) (Table III).

TABLE II
THE EFFECT OF AZASERINE AND STRONG LIGHT ON
Scenedesmus

	Control III	Azaserine III (1 mg.)
Determination on aluminum disc, counts/min.		
Total fixation	20.9×10^6	14×10^6
80% ethanol extract	47.8%	57.8%
20% ethanol extract	2.5%	4.4%
Determination on paper, counts/min.		
Glutamine	529	4964
Glutamic acid	4951	2945
Aspartic acid	9089	2262
Serine	2834	1591
Alanine	2345	951
Glycine } area	949	929
Threonine		
Sucrose	1357×9.5	1907×9.5
Malic acid	3743×9.5	6479×9.5
Citric acid	254×9.5	1919×9.5
α -Ketoglutaric acid	381	1851
Fumaric acid	344	540
Succinic acid	1621	3180
Glyceric acid	1553	1312
Glycolic acid	285	506
Lipids	3582×9.5	2491×9.5
Phospholipids	2259×9.5	1665×9.5
Area X	725×9.5	506×9.5
Sugar phosphates, PGA and origin	$11,800 \times 9.5$	9312×9.5

TABLE III
ATTEMPTED REVERSAL OF AZASERINE INHIBITION WITH
GLUTAMINE

	Azaserine (4 mg.) alone	Azaserine (4 mg.) + glutamine (5 mg.)
Determination on aluminum disc, counts/min.		
Total fixation	33.6×10^6	35.8×10^6
80% ethanol extract	41.2%	52.9%
20% ethanol extract	1.15%	2.22%
Determination on paper, counts/min.		
Glutamine	2182	6572
Glutamic acid	1353	2704
Aspartic acid	666	939
Serine	735	690
Alanine	393	473
Glycine } area	503	527
Threonine		
Sucrose	1496×9.5	1111×9.5
Malic acid	3697×9.5	4631×9.5
Citric acid	586×9.5	446×9.5
α -Ketoglutaric acid	696	1052
Fumaric acid	242	249
Succinic acid	870	1308
Glycolic acid	3424	3138
Lipids	1639×9.5	2192×9.5
Phospholipids	541×9.5	1310×9.5
Area X	797×9.5	1370×9.5
Sugar phosphates, PGA and origin	$11,970 \times 9.5$	$12,383 \times 9.5$

Chromatograms from each of the above experiments were sprayed with ninhydrin to obtain a qualitative estimate of the concentrations of amino acids present. Glutamine obtained from azaserine chromatograms, as well as the glutamic acid produced from it by hydrolysis with 1 N HCl

TABLE IV
 THE EFFECT OF AZASERINE ON *Chlorella*

	Control IV	Azaserine IV (4 mg.) - air flushed 30 sec.	Control V	Azaserine V (4 mg.) - air flushed 60 sec.
Determination on aluminum disc, counts/min.				
Total fixation	25.4×10^6	30.8×10^6	23.4×10^6	34.8×10^6
80% ethanol extract	60.7%	66.4%	62%	62.8%
20% ethanol extract	8.3%	8.25%	10.6%	8.1%
Determination on paper, counts/min.				
Glutamine	250	300	623	530
Glutamic acid	2203	2290	2370	2006
Aspartic acid	2541	2700	1900	2180
Serine	2866	3361	3890	4770
Alanine	9635	8232	10,300	9670
Glycine } area	800	960	780	830
Threonine }				
Sucrose	4458×9.5	3933×9.5	4350×9.5	4330×9.5
Malic acid	2243	2409	1980	1970
Citric acid	245	250	200	220
Fumaric acid	571	428	380	300
Lipids } area	2101×9.5	1895×9.5	2800×9.5	2750×9.5
Phospholipids }				
Area X	880×9.5	858×9.5	630×9.5	640×9.5
Sugar phosphates, PGA and origin	$11,583 \times 9.5$	$12,085 \times 9.5$	$18,690 \times 9.5$	$19,140 \times 9.5$

at 100° for 2 hours, were characterized by cochromatography on two-way paper chromatograms and by ionophoresis on paper for 3 hours at 600 v. in 0.1 M propionate, pH 5.6. Sucrose, amino acids and carboxylic acids mentioned in Table I were likewise characterized by cochromatography on two-way paper chromatograms and most of them also by ionophoretic separations. The lipids, phospholipids and pigments were all moved at a similar distance in phenol-water but separated in butanol-propionic acid into two mobile areas designated lipid (containing some pigment), phospholipid (containing no visible pigment) and a stationary area designated X (containing some pigment). No radioactivity could be extracted from these areas by water elution but most of the activity was extracted with a mixture of petroleum ether and ethanol. The compounds in area X ran faster than diphosphopyridine nucleotide in phenol-water but showed the same lack of mobility in butanol-propionic acid.

The triose phosphate, pentose monophosphate, glucose cyclic 1,2-phosphate, hexose monophosphate, phosphoglyceric acid and phosphoenolpyruvate were eluted from control and azaserine chromatograms, treated with purified "Polidase" phosphatase, and the products analyzed by paper chromatography and radioautography.⁵

The effect of azaserine (4 mg.) on *Chlorella* was also determined in two experiments (Table IV) identical to that described for the *Scenedesmus* model experiment except that in one case the time of flushing with air was 30 sec. and in the other, 60 sec.

Discussion

It is evident from Table I that with *Scenedesmus* the azaserine causes a build-up of the acids (e.g., citric, malic, α -ketoglutaric, succinic) of the Krebs tricarboxylic acid cycle and a decrease in radioactivity and amount (revealed by ninhydrin) of the amino acids (glutamic acid, aspartic acid) derived therefrom by amination or transamination. Alanine, serine, valine and tyrosine, which are also known to be formed by transamination of their corresponding keto-acids, were likewise depleted. Because of the low activity of glycine and threonine and the presence of other compounds in this area the effect of azaserine on these amino acids could not be assessed. Glutamine,⁶ which is a donor of amino-groups during transamination, increases markedly in activity and amount (approximately

1 mg./g. of cells as estimated by ninhydrin) in the presence of azaserine. It is therefore suggested that azaserine interferes with transamination reactions either directly by inhibiting the actual process of transamination or indirectly by interfering with the synthesis of pyridoxal phosphate or pyridoxamine phosphate which are necessary as coenzymes of such reactions. The second possibility seems less likely because of the short time available for the inhibition to appear. Figure 1 depicts some of the suggested sites of azaserine inhibition. Kaplan and Stock⁷ have also concluded that azaserine is an inhibitor of amino acid synthesis on the basis of their observation that inhibition of growth in *E. coli* by azaserine can be overcome by addition of certain amino acids. Two azaserine-inhibited reactions, which have been recently reported, would also fit into this scheme. It has been shown that azaserine inhibits the conversion of α -N-formyl glycylamide ribotide to α -N-formylglycylamidine ribotide,⁴ and the amination of xanthosine-5'-phosphate to guanosine-5'-phosphate.⁸ L-Glutamine is the donor of amino groups in both these reactions.

An interesting feature is the build-up of glutamine despite the fact that the formation of glutamic acid (its precursor) is partially inhibited by azaserine. It is to be noted, however, that the decrease in glutamic acid is by no means as marked as the decreased formation of aspartic acid in the presence of azaserine. One explanation of this may be that while it is known⁹ that glutamic acid can be synthesized from NH_4^+ and α -ketoglutaric acid, the synthesis of aspartic acid by a similar reaction of NH_4^+ and oxalacetic acid has not been proved. This alternative pathway of glutamic acid synthesis, not involving transamination, probably operates in *Scenedesmus*.

(7) L. Kaplan and C. C. Stock, *Federation Proc.*, **13**, 239 (1954).

(8) M. Bentley and R. Adams, *ibid.*, **15**, 218 (1955).

(9) H. A. Krebs and P. D. Cohen, *Biochem. J.*, **33**, 1895 (1939).

(6) A. Meister and S. V. Tice, *J. Biol. Chem.*, **187**, 173 (1950).




Fig. 1.—Possible sites of azaserine inhibition.

Since the formation of serine is blocked partially by azaserine, smaller amounts of ethanolamine and choline (which are derived from it) will be available for incorporation into phospholipids and this may account for the decreased radioactivity found in this region of the chromatograms in the presence of azaserine. The lipid and area X also show decreased activity. This might be accounted for, at least in part, by blockage of a transamination reaction leading to glycine which is necessary for the synthesis of the heterocyclic rings in the porphyrin pigments.¹⁰

Glycolic acid, which generally accumulates⁵ at low CO_2 pressures, was considerably increased at higher levels of azaserine. This could not be caused by a lower rate of CO_2 fixation since this is increased by azaserine during photosynthesis. The increased levels of acids in the Krebs cycle may cause an increased formation of glyoxylic acid,¹¹ which with an inhibited glycine formation would be available for glycolic acid production. The increased rate of CO_2 fixation in the presence of azaserine is probably responsible for the increased level of sucrose drained from the photosynthetic cycle.

Examination of the components of the photosynthetic cycle, by analysis of the materials produced by phosphatase action, revealed the presence of glyceric acid and all the usual sugars. Several additional unidentified components were detected in the phosphatased triose phosphate and glucose cyclic 1,2-phosphate areas eluted from azaserine chromatograms.

(10) R. J. Della Rosa, K. I. Altman and K. Salomon, *J. Biol. Chem.*, **202**, 771 (1953).

(11) R. A. Smith, J. R. Stamer and I. C. Gunsalus, *Biochem. Biophys. Acta*, **19**, 567 (1956).

The total fixations in the experiments, in which an intense photospot light was substituted for one of the reflector floods for 50 minutes of the one hour steady state period (Table II), indicate that a larger proportion of the algae were killed by the bright light in the presence of azaserine than in its absence. However, control and azaserine chromatograms, each carrying a total activity of 1×10^6 counts/min., showed spots (Table II) with the same order of relative activities as those given in Table I.

An attempt to reverse the effects of azaserine inhibition with glutamine was only partially successful (Table III). Some increase in radioactivity in the lipid, phospholipid, area X and glutamic acid was observed. The apparent continued build-up of glutamine can be explained by suggesting that while the rate of conversion of C^{14} -labeled precursors to glutamine remains almost the same, the introduction of inactive glutamine causes a dilution of radiocarbon in the glutamine pool with a consequent lowering in the specific activity of the glutamine being converted to further products and hence an increase in the residual C^{14} -glutamine observed. It is interesting to note that the combined effect of azaserine and added inactive glutamine has caused almost a twenty-fold increase of C^{14} -glutamine.

Although azaserine caused a marked increase in CO_2 fixation by *Chlorella* (Table IV), analysis of aliquots containing equal amounts of radioactivity showed that the radioactivity of most of the metabolites, relative to each other, remained the same.