



TABLE I

Compound	B.p., °C.	M.p., °C.	$d^{25}$	$n_D^{25}$	Nitrogen, %		Calcd.	Found
					Calcd.	Found		
CF <sub>3</sub> C(:NH)NHCH <sub>3</sub>	47-48 (21 mm.)		1.2909	1.3919	22.20	21.85		
CF <sub>3</sub> C(:NH)N(CH <sub>3</sub> ) <sub>2</sub>	92		1.1795	1.3812	20.01	19.72		
C <sub>2</sub> F <sub>5</sub> C(:NH)NHCH <sub>3</sub>	126		1.4258	1.3664	15.91	15.63		
C <sub>2</sub> F <sub>5</sub> C(:NH)N(CH <sub>3</sub> ) <sub>2</sub>	110		1.2969	1.3678	14.75	14.43		
C <sub>3</sub> F <sub>7</sub> C(:NH)NHCH <sub>3</sub>	137		1.5140	1.3561	12.37	12.02		
C <sub>3</sub> F <sub>7</sub> C(:NH)N(CH <sub>3</sub> ) <sub>2</sub>	120		1.4242	1.3548	11.66	11.41		
CF <sub>3</sub> C(:NH)NHAg		200 dec.			12.71	12.28	48.78 <sup>a</sup>	48.40
C <sub>2</sub> F <sub>5</sub> C(:NH)NHAg		240 dec.			10.36	10.27	40.02 <sup>a</sup>	40.38
C <sub>3</sub> F <sub>7</sub> C(:NH)NHAg		277 dec.			8.74	8.50	33.60 <sup>a</sup>	32.86
(C <sub>2</sub> F <sub>5</sub> C(:NH)NH) <sub>2</sub> Hg		159 dec.			10.65	10.22	38.38 <sup>a</sup>	38.20
(C <sub>3</sub> F <sub>7</sub> C(:NH)NH) <sub>2</sub> Hg		178 dec.			9.00	8.42	32.31 <sup>a</sup>	32.60
CF <sub>3</sub> C(:NH)NH <sub>2</sub> ·CH <sub>3</sub> COOH		Subl.			16.29	15.94	172 <sup>b</sup>	174
CF <sub>3</sub> C(:NH)NH <sub>2</sub> ·C <sub>3</sub> F <sub>7</sub> COOH		Subl.			8.58	8.80	326 <sup>b</sup>	323
C <sub>3</sub> F <sub>7</sub> C(:NH)NH <sub>2</sub> ·CH <sub>3</sub> COOH		Subl.			10.20	10.26	272 <sup>b</sup>	277
C <sub>3</sub> F <sub>7</sub> C(:NH)NH <sub>2</sub> ·C <sub>3</sub> F <sub>7</sub> COOH		Subl.			6.58	6.43	426 <sup>b</sup>	420
CF <sub>3</sub> C(:S)NH <sub>2</sub>	40 (2 mm.)				10.82	10.85	24.78 <sup>c</sup>	25.13
C <sub>2</sub> F <sub>5</sub> C(:S)NH <sub>2</sub>	60 (23 mm.)	39-40			7.82	7.53	17.88 <sup>c</sup>	18.18
C <sub>3</sub> F <sub>7</sub> C(:S)NH <sub>2</sub>	69 (21 mm.)	47-48			62.1	6.01	13.97 <sup>c</sup>	13.90

<sup>a</sup> Silver (mercury), %. <sup>b</sup> Neut. equiv. <sup>c</sup> Sulfur, %.

pounds are pale yellow solids or liquids and may be stored in sealed vials without decomposition.

### Experimental

**Trifluoroacetonitrile, Pentafluoropropionitrile and Heptafluorobutyronitrile.**—These compounds were prepared by dehydration of the perfluoroamides as described by Swarts<sup>4</sup> and by Gilman and Jones.<sup>5</sup>

**N-Methylperfluoroacetamide.**—Into an 80-ml. cylindrical glass flask immersed in a Dry Ice-acetone bath and fitted with a delivery tube and Dry Ice cooled reflux condenser was condensed 11.2 g. (0.1 mole) of trifluoroacetonitrile; 6.2 g. (0.2 mole) of methylamine was added and the mixture allowed to reflux under a Dry Ice condenser for one hour. Excess amine was distilled from the mixture and the remaining high boiling liquid fractionated under reduced pressure to yield 14 g. of N-methylperfluoroacetamide, b.p. 35-36° at 11 mm.

The other N-alkyl perfluoroacetamides shown in Table I were prepared in the same manner and in comparable yields. All of this series could be fractionated at atmospheric pressure with the exception of the N-methyltrifluoroacetamide as described above.

**Trifluoroacetamide.**<sup>3</sup>—Trifluoroacetonitrile, 17.0 g. (0.18 mole), was condensed in an apparatus similar to that described above for the preparation of N-methylperfluoroacetamide. A large excess of anhydrous ammonia was condensed in the reaction vessel, the Dry Ice-acetone bath removed, and the mixture of ammonia and nitrile allowed to reflux for one hour. Excess ammonia was then removed by distillation and the product fractionated under reduced pressure in a small column packed with glass helices to yield 17.5 g. of trifluoroacetamide, b.p. 35-36° (11 mm.),  $n_D^{25}$  1.3801,  $d^{25}$  1.4940.

Pentafluoropropionamide and heptafluorobutyramidine were produced by this same procedure in 95 and 75% yields, respectively. These compounds are low melting solids and were recrystallized from an ethyl ether-petroleum ether mixture. They were found not to be hygroscopic if completely free of ammonia.

**Silver Salts of Trifluoroacetamide.**—Silver oxide, 4.60 g. (0.020 mole), was added slowly to 5 g. (0.044 mole) of trifluoroacetamide dissolved in 15 ml. of anhydrous ethyl ether. An immediate evolution of heat was noted with the formation of an insoluble white solid. The mixture was stirred vigorously to break up any lumps of silver oxide. When the black silver oxide was consumed, the white solid was filtered, washed with ethyl ether and dried to give slightly less than a quantitative yield of the silver salt of trifluoroacetamide, CF<sub>3</sub>C(NH)NHAg. This salt decomposed on heating at 200° (Table I).

(4) F. Swarts, *Bull. class. sci. acad. roy. Belg.*, [5] **12**, 692 (1926).

(5) H. Gilman and R. C. Jones, *THIS JOURNAL*, **65**, 1458 (1943).

Silver salts of pentafluoropropionamide and of heptafluorobutyramidine were prepared in the same manner.

**Mercury Salt of Heptafluorobutyramidine.**—In a test-tube immersed in an oil-bath was placed an intimate mixture of 10 g. (0.046 mole) of heptafluorobutyramidine, C<sub>3</sub>F<sub>7</sub>C(NH)NH<sub>2</sub> and 5 g. (0.023 mole) of mercuric oxide. The mixture was heated to 110° to melt the amide, then stirred vigorously. The formation of the white mercury salt could be noted and was essentially complete in 2 hours. The melt was cooled and unreacted amide removed by continuous extraction with ethyl ether. Unreacted mercuric oxide was removed by centrifugation of a suspension of the powdered reaction product in ethyl ether. The heavier mercuric oxide settled first leaving the upper layer as essentially the mercury salt of the amide. A number of centrifugations produced a low yield of the pure mercury salt of heptafluorobutyramidine, (C<sub>3</sub>F<sub>7</sub>C(NH)NH)<sub>2</sub>Hg, a white solid that decomposed on heating at 178° (Table I).

The mercury salt of pentafluoropropionamide was prepared by this same procedure. Trifluoroacetamide could not be used in this reaction since it was not sufficiently stable at elevated temperatures.

**The Trifluoroacetic Acid Derivative of Heptafluorobutyramidine.**—To 15 ml. of trifluoroacetic acid was added 5.3 g. (0.025 mole) of heptafluorobutyramidine, C<sub>3</sub>F<sub>7</sub>C(NH)NH<sub>2</sub>. The clear solution was allowed to stand one hour; the excess trifluoroacetic acid was then removed under reduced pressure. The remaining white solid was purified by sublimation to give 6.8 g. of the acid derivative of heptafluorobutyramidine, C<sub>3</sub>F<sub>7</sub>C(NH)NH<sub>2</sub>·CF<sub>3</sub>COOH (Table I.).

The other organic and perfluoro acid derivatives were prepared by a similar procedure in 75-85% yields.

**N-Benzoylheptafluorobutyramidine.**—A solution of 24.6 g. (0.174 mole) of benzoyl chloride in ether was placed in a flask protected from atmospheric moisture by drying tubes. The contents of the flask were cooled to 0° and a solution of 37 g. (0.174 mole) of heptafluorobutyramidine in ether was slowly added with stirring. Evolution of hydrogen chloride was noted and this produced a white precipitate of heptafluorobutyramidine hydrochloride. The flask was allowed to warm to room temperature and 23 g. of the hydrochloride removed by filtration. Dilution of the ethereal solution with petroleum ether and cooling precipitated 20 g. of N-benzoylheptafluorobutyramidine. After recrystallization from ethyl ether-petroleum ether, this product melted at 63°.

*Anal.* Calcd. for C<sub>11</sub>F<sub>7</sub>H<sub>7</sub>N<sub>2</sub>O: N, 8.86. Found: N, 8.58.

**Heptafluorothiobutyramide.**—A. **From Heptafluorobutyronitrile.**—Into an evacuated heavy walled Pyrex glass tube of approximately 60-ml. capacity, previously constricted for sealing, was condensed 6.8 g. (0.035 mole) of perfluorobutyronitrile, C<sub>3</sub>F<sub>7</sub>CN, and 3.4 g. (0.10 mole) of hydrogen sulfide. The tube was sealed, allowed to

remain at room temperature overnight, then opened and the volatile material removed. The remaining product was a pale yellow solid that was recrystallized from ether-petroleum ether to yield 6.3 g. of heptafluorothiobutyramide,  $C_3F_7C(S)NH_2$ , m.p.  $49^\circ$  (Table I).

Pentafluorothiopropionamide was prepared in the same manner from pentafluoropropionitrile and hydrogen sulfide at room temperature. For preparation of trifluorothioacetamide the reaction was moderated by cooling to  $0^\circ$  to prevent charring. This reaction could also be moderated by the use of ethyl ether as a solvent but reaction time was increased to 125 hours.

**Heptafluorothiobutyramide.—B.** From **Heptafluorobutyramidine.**—In a flask fitted with a delivery tube and protected by drying tubes was placed 15 g. (0.071 mole) of heptafluorobutyramidine dissolved in 25 ml. of ethyl ether. The solution was saturated with hydrogen sulfide and allowed to stand one hour. The ether was removed under reduced pressure and the remaining high boiling liquid fractionated under reduced pressure to yield 12 g. of a distillate that crystallized on cooling. Recrystallization from ether-petroleum ether gave heptafluorothiobutyramide,  $C_3F_7C(S)NH_2$ , yellow needles, m.p.  $47-48^\circ$ .

GAINESVILLE, FLORIDA

[CONTRIBUTION FROM THE ILLINOIS STATE GEOLOGICAL SURVEY]

## Aromatic Fluorine Compounds. VII. Replacement of Aromatic $-Cl$ and $-NO_2$ Groups by $-F$ <sup>1,2</sup>

By G. C. FINGER AND C. W. KRUSE

RECEIVED AUGUST 9, 1956

Replacement of  $-Cl$  by  $-F$  in aryl chlorides with potassium fluoride has been extended from 2,4-dinitrochlorobenzene to less activated halides by the use of non-aqueous solvents, especially dimethylformamide (DMF) and dimethyl sulfoxide (DMSO). Also replacement of  $-NO_2$  by  $-F$  in substituted nitrobenzenes was studied in DMF. As a direct result of this study, many aromatic fluorine compounds can now be obtained by a relatively simple synthetic route.

Unactivated aryl halides do not undergo Swarts-type organic halide-inorganic fluoride exchange reactions which are common with aliphatic and acyl halides.<sup>3,4</sup> Activation by at least two nitro groups has generally been considered necessary for exchange between aryl halides and fluoride ion.<sup>5a</sup> An outstanding example of such an exchange is the replacement of chlorine by fluorine in 2,4-dinitrochlorobenzene with anhydrous potassium fluoride at  $190-205^\circ$  in nitrobenzene solvent.<sup>6,7</sup>



A preliminary experiment has shown that even without a solvent the above reaction takes place at  $200-230^\circ$ . However, solvent studies with other non-aqueous solvents show a significant increase in reaction rate. The preceding reaction proceeds quite smoothly at steam-bath temperature in dimethylformamide (DMF) or dimethyl sulfoxide (DMSO). The accelerating effect of various solvents upon aryl halide-fluoride ion exchanges is emphasized by Table I. The decreasing reaction rates, as reflected by the longer time and higher temperatures, parallel the decreased activation from 2,4-dinitrochlorobenzene to 4-nitrochlorobenzene. Thus, aromatic halogen exchange reac-

tions have been extended from the highly activated dinitrohalobenzenes to the moderately activated mononitrohalobenzenes by the solvent studies.

TABLE I  
REPLACEMENT OF AROMATIC  $-Cl$  BY  $-F$

Reactant <sup>a</sup> (subst. $C_6H_4NO_2$ )	Solvent	Time, hr.	Temp., $^\circ C.$	Yield, % of fluoro analog	Expt. ref.
3-NO <sub>2</sub> -4-Cl-	$C_6H_5NO_2$	4	195-210	76	I-1
	No solvent	0.5	200-230	62 <sup>b</sup>	I-2
	$C_6H_5CN$	2	150-170	57 <sup>b</sup>	I-3
	$(CH_3CN)_2$	0.5	125-140	81 <sup>b</sup>	I-4
	DMF	0.5	140-150	77	I-5
	DMF	13	95-100	77	I-6
3-CF <sub>3</sub> -4-Cl-	DMSO	2.5	95-100	78	I-7
	DMF	4	160	81	II
2-Cl-5-CF <sub>3</sub> -	DMF	3.5	160	76	III
2-Cl-5-CH <sub>3</sub> OCO-	DMF	4	155	67	IV
2,4,6-Cl <sub>3</sub> -	DMSO	2	175-180	10	V
2,5-F <sub>2</sub> -4-Cl-	DMSO	2	170-175	45	VI
2,3,4-Cl <sub>3</sub> -	DMSO	4	180-190	23	VII
2,4-Cl <sub>2</sub> -	DMSO	6	180	47	VIII
2-Cl-	DMF	163	170	40 <sup>c</sup>	IX-1
	DMSO	4.5	185	38	IX-2
4-Cl-	DMSO	14	190	72	X

<sup>a</sup> Group replaced is set in bold type. <sup>b</sup> Contaminated with chloro isomer. Vacuum fractionation was ineffective for the separation of these isomers, therefore yields were estimated by index of refraction which was found to have a near linear relationship for mixtures of 2,4-dinitrochloro- and 2,4-dinitrofluorobenzenes. Values determined from known mixtures for weight-per cent. of chloro isomer *vs.*  $n_D^{20}$  were 80.4%-1.5952, 54.9%-1.5866 and 23.12%-1.5764. <sup>c</sup> Estimated.

Brown fumes and traces of lower boiling products observed in the 2,4-dinitrohalobenzene studies suggested the possibility of some  $-NO_2$  replacement. Earlier investigators<sup>6,7</sup> make no mention of such observations. Such a reaction appeared reasonable as properly activated nitro groups can be replaced by other nucleophilic reagents, as, for example, the replacement of  $-NO_2$  by  $-OCH_3$  in 2,3,5,6-tetrachloronitrobenzene.<sup>8</sup> Conclusive evi-

(8) V. S. F. Berckmans and A. F. Holleman, *Rec. Trav. chim.*, **44**, 851 (1925).

(1) Presented in part before the Organic Division, 129th Meeting of the American Chemical Society, April, 1956, Dallas, Texas. This research was supported in part by the United States Air Force under Contract No. AF 18(600)-985, monitored by the AF Office of Scientific Research of the Air Research and Development Command.

(2) Published by permission of the Chief of the Illinois State Geological Survey.

(3) "Organic Reactions," Vol. II, John Wiley and Sons, Inc., New York, N. Y., 1944, Ch. 2, p. 49.

(4) F. L. M. Pattison, *Nature*, **174**, 740 (1954).

(5) (a) J. F. Bunnett and R. E. Zahler, *Chem. Revs.*, **49**, 277 (1951); (b) p. 315.

(6) H. B. Gottlieb, *This Journal*, **58**, 532 (1936).

(7) H. G. Cook and B. C. Saunders, *Biochem. J.*, **41**, 558 (1947).