

cinic acid complexes, do not exhibit enhanced reaction rates. An "adsorption" mechanism is ruled out, for the rate of reduction of aquopentamminocobaltic perchlorate is not affected by the mere presence of cobalt-free polymer. We lean to the hypothesis that the rapid reaction corresponds to those cobalt-bearing sidechains of which the carboxyl groups overlap an extended system of hydrogen-bonded amide groups; and that electron transfer from chromium occurs *through* this system.

The products of the reduction of the polymeric complexes have been isolated by dialysis and lyophilization of the reaction mixture. They contain chromium equivalent to the initial cobalt, but the sites of this bound chromium are not known. These products are not entirely soluble in water and dissolve sluggishly in formic acid. A film of the polymer has $\nu_{C=O}$ 1650 cm^{-1} , but treatment with water at 100° does not produce any β -form (1625 cm^{-1}). It is conceivable that the rapid reduction process produces a change in the conformation of the polymer which constrains the remainder of reaction to occur by a process involving direct chromium attack at the carboxyl bearing the oxidant.

We are grateful to Professor Henry Taube for his interest in this work.

DEPARTMENT OF CHEMISTRY
UNIVERSITY OF CHICAGO
CHICAGO 37, ILLINOIS

KENNETH D. KOPPLE
GEORGE F. SVATOS

RECEIVED APRIL 25, 1960

THE ISOMERIZATION OF TRIALKYLACETIC ACIDS IN SULFURIC ACID¹

Sir:

Trialkylacetic acids have been found to isomerize in concentrated sulfuric acid under conditions which do not result in significant decarbonylation. The same mixture consisting chiefly of 2,2-dimethylvaleric acid (I)² and 2-methyl-2-ethylbutyric acid (II)² resulted from the isomerization of either I or II in concentrated sulfuric acid. Solutions of either I or II (0.27 molar) in 94.9% sulfuric acid approached the equilibrium ratio (I/II) of 2.84 with first order kinetics and a half life of 51 minutes at 26.3°. The appearance of a third component 2,2,3-trimethylbutyric acid (III),³ proceeded much slower. This acid accounted for *ca.* 3% of the mixture after three hours and became about equal to II after 48 hours. The isomerization of III under these conditions slowly yielded I and II in the ratio of 2.84.

Acid II was isomerized in sulfuric acid, the mixture of carboxylic acids separated and converted to the amides. From this mixture the pure amide of I² (identified by infrared, m.p. and mixed m.p. 94–95°) was isolated by fractional crystallization. The acid mixtures were analyzed by gas chromatography of the corresponding alcohols (obtained

(1) Presented at the 6th Tetrasectional Meeting of the Oklahoma Sections of the American Chemical Society, March 26, 1960.

(2) Compound I was prepared by oxidation of 2,2-dimethyl-1-pentanol obtained from the Carbide Chemical Corporation, and II was prepared from the Grignard. The melting points of the amides (94–95 and 74–75°) and the physical properties of the acids agree with values previously reported; D. V. N. Hardy, *J. Chem. Soc.*, 464 (1938).

(3) F. C. Whitmore and K. C. Laughlin, *THIS JOURNAL*, **55**, 3735 (1933).

by LiAlH_4 reduction) using a 12' column of Resoflex⁴ R-726 on fire brick. Infrared spectrograms of isomerized mixtures were identical with appropriate mixtures of I, II, and III.

The four nine carbon tri-*n*-alkylacetic acids were interconverted rapidly under the same conditions to yield 2,2-dimethylheptanoic acid as the major component. This emphasizes the general nature of this isomerization. The disappearance of 2,2-diethylvaleric acid, which represented *ca.* 1% of the isomerized mixture, proceeded by initially first order kinetics with a half life of less than five minutes. Slower changes occurred to yield products which are presumably analogous to III. The thermodynamically most stable species appear to predominate in isomerized mixtures of both the seven and nine carbon trialkylacetic acids.⁵

That this isomerization occurred at least in part by a decarbonylation-carbonylation process was demonstrated by the exchange of radioactive II (labeled with C^{14} in the carboxyl carbon) in concentrated sulfuric acid with unlabeled carbon monoxide.⁶ Also, slow introduction of 2-methyl-2-butene into a sulfuric acid solution of II resulted in the formation of 2,2-dimethylbutyric acid.

The author thanks Messrs. P. L. Gant and R. D. Gallagher for assistance in the tracer experiment and Dr. M. G. Ettlinger for a helpful discussion.

(4) Cambridge Industries Co., Inc., Cambridge, Mass.

(5) Similar reactions in which the more stable isomer predominated have been reported by R. E. Pincock, E. Grigat and P. D. Bartlett, *THIS JOURNAL*, **81**, 6332 (1959); G. Stork and M. Bersohn, *ibid.*, **82**, 1261 (1960).

(6) A carbonylation-decarbonylation mechanism has been demonstrated to occur in the isomerization of cinenic acid to geronic acid by J. Meinwald, H. C. Hwang, D. Christman and A. P. Wolf, *ibid.*, **82**, 483 (1960).

RESEARCH AND DEVELOPMENT DEPARTMENT
CONTINENTAL OIL COMPANY
PONCA CITY, OKLAHOMA

ALLAN LUNDEEN

RECEIVED APRIL 16, 1960

THE ISOLATION OF THE ICOSAHEDRAL $\text{B}_{12}\text{H}_{12}^{-2}$ ION

Sir:

In 1955 Longuet-Higgins and Roberts¹ predicted that the $\text{B}_{12}\text{H}_{12}$ icosahedron would be stable only as the $\text{B}_{12}\text{H}_{12}^{-2}$ ion. We now wish to report the isolation of such an ion, in very low yield, from the reaction of 2-iododecaborane with triethylamine.

Sixteen grams of pure 2-iododecaborane and 35 g. of triethylamine were dissolved in 500 ml. of dry benzene and maintained at the reflux temperature for 5 hours. The yellow semi-solid which separated was treated with 150 ml. of acetone, the resulting solid separated by filtration and washed with water to remove the triethylammonium salt² of $\text{B}_{10}\text{H}_{16}^{-2}$ formed as a by-product. The residual solid was recrystallized twice from acetonitrile and ethanol to yield 1.4 g. (3.8%) of the triethylammonium salt of $\text{B}_{12}\text{H}_{12}^{-2}$ (I). Calcd.: B, 37.54; C, 41.63; H, 12.72; N, 8.09. Found: B, 37.48; C, 40.86; H, 12.97; N, 8.08.

Treatment of I with two equivalents of potassium hydroxide in boiling water displaced triethylamine

(1) H. C. Longuet-Higgins and M. Roberts, *Proc. Royal Soc. (London)*, **A230**, 110 (1955).

(2) M. F. Hawthorne and A. R. Pitochelli, *THIS JOURNAL*, **81**, 5519 (1959).