


we have isolated 8-hydroxyphenillic acid from such fermentations. When phenylacetic acid is omitted, relatively large quantities of both 6-aminopenicillanic acid and 8-hydroxyphenillic can be isolated. When excess phenylacetic acid is present, both of these substances are produced but in much smaller concentrations.

The published literature reveals that 8-hydroxyphenillic acid has been encountered but unrecognized by other investigators. In 1959, Tardrew and Johnson<sup>14</sup> reported the isolation of "Compound VI" from precursor-free *P. chrysogenum* fermentation broths. While they failed to identify the substance, they concluded that it was a stabilization product of a biosynthetic precursor of the penicillins. In the light of our findings, a review of their results leaves little doubt that "Compound VI" was 8-hydroxyphenillic acid.

The fate of sulfur in *P. chrysogenum* fermentations can, therefore, be further defined



The anomalous results reported recently by Steinman<sup>15</sup> using the manometric assay for studying the rate of 6-APA hydrolysis by penicillinase can be explained by reaction of some of the carbon dioxide with 6-APA to form 8-HPA.

It seems likely that nearly everyone working with 6-aminopenicillanic acid will, at some time, by chance or by design prepare 8-hydroxyphenillic acid.

(14) P. L. Tardrew and M. J. Johnson, *J. Biol. Chem.*, **234**, 1850 (1959).

(15) H. G. Steinman, *Proc. Soc. Exptl. Biol. and Med.*, **106**, 227 (1961).



CHEMICAL DEVELOPMENT DIVISION  
BRISTOL LABORATORIES  
DIVISION OF BRISTOL-MYERS COMPANY  
SYRACUSE 1, NEW YORK

DAVID A. JOHNSON  
GLENN A. HARDCASTLE, JR.

RECEIVED JULY 1, 1961

#### A NEW GENERAL REACTION FOR PREPARING *gem* DINITRO COMPOUNDS: OXIDATIVE NITRATION Sir:


The methods available for preparing *gem* dinitro compounds are of limited applicability or are very inefficient.<sup>1</sup> We wish to report a new general reaction, oxidative nitration, in which salts of primary and secondary nitro compounds are converted into the corresponding *gem* dinitro derivatives by reaction with silver nitrate and inorganic nitrites in alkaline or neutral aqueous media (Equations 1 and 2). Oxidation-reduction proceeds rapidly



(1) (a) G. Born, *Ber.*, **29**, 90 (1896); (b) E. ter Meer, *Ann.*, **181**, 1 (1876); J. S. Belew and L. G. Hepler, *J. Am. Chem. Soc.*, **78**, 4005 (1956); (c) W. I. Denton, R. B. Bishop, E. M. Nygaard and T. T. Noland, *Ind. Eng. Chem.*, **40**, 381 (1948); (d) C. T. Bahner, *ibid.*, **44**, 317 (1952); (e) M. G. Chancel, *Bull. soc. chim.*, **31**, 504 (1879); M. Fileti and G. Ponzio, *J. prakt. Chem.*, **55**, 195 (1897).

from homogeneous solution at 0–30° to yield an addition-complex which decomposes into *gem* dinitro compound and silver. Primary, secondary, and functionally-substituted dinitro compounds such as 1,1-dinitroethane, 1,1-dinitropropane, 2,2-dinitropropane, 2,2-dinitrobutane,<sup>2a</sup> 1,1-dinitrocyclohexane, 2,2,4,4-tetranitropentane,<sup>2b</sup> 2,3-dimethyl-2,4,4-trinitropentane,<sup>2c</sup> 2,2-dinitro-1-propanol, 1,1-dinitro-2-propanol, 2,2-dinitro-1,3-propanediol, 4,4-dinitropentanal and methyl 3,3-dinitropropionate may be prepared efficiently (60–95%) from their corresponding nitro derivatives. Sensitive or hindered compounds such as 3,3-dinitro-2-butanol,<sup>2d</sup> 2,2-dimethyl-1,1,3-trinitropropane<sup>2e</sup> and 1-cyclopropyl-1,1-dinitroethane,<sup>2f</sup> substances which cannot be prepared satisfactorily by other known methods, have been synthesized.

Dinitromethane has been obtained as its potassium salt from nitromethane<sup>3</sup> or much better from 1-nitro-2-propanol via base-catalyzed decomposition of 1,1-dinitro-2-propanol. The most practical method for preparing potassium dinitromethane (>53%) or potassium 2,2-dinitroethanol (99%) is by controlled alkaline demethylation (Equations 3 and 4) of 2,2-dinitro-1,3-propanediol ob-



tained by oxidative nitration (70–80%) of 2-nitro-1,3-propanediol. Under different conditions 2,2-dinitro-1,3-propanediol is converted by potassium hydroxide to dipotassium 1,1,3,3-tetranitropropane<sup>2g,4</sup>; this salt apparently is formed by reaction of potassium dinitromethane and potassium hydroxide with 1,1-dinitroethylene<sup>5</sup> generated by decomposition of potassium 2,2-dinitroethanol.

Oxidative nitration of salts of 1,1-dinitro compounds does not give 1,1,1-trinitromethyl derivatives.  $\alpha$ -Arylalkanenitronates yield *vicinal* dinitro compounds,  $\text{R}_2\text{C(NO}_2)\text{C(NO}_2)\text{R}_2$ , by oxidative dimerization along with carbonyl derivatives and *gem* dinitro compounds. Thus phenylnitromethane gives phenyldinitromethane (19%), benzaldehyde (36%) and *meso* and *d,l*-1,2-dinitro-1,2-diphenylethanes (12 and 25%, respectively); 9-nitrofluorene yields 9,9-dinitrofluorene (8%), fluorenone (8%), and 9,9'-dinitrodifluorenyl (76%). The effects of other functional groups on the oxidative nitration reaction are being studied.

The silver obtained may be separated easily and recovered essentially quantitatively as silver nitrate. Mercuric nitrate has been successfully sub-

(2) New compounds: (a) b.p. 78° (10 mm.): C, 32.18; H, 5.37; (b) M.p. 87.5°: C, 23.95; H, 3.18; N, 22.39. (c) M.p. 83°: C, 36.15; H, 5.86; N, 17.71. (d) B.p. 73–75° (2 mm.): C, 29.10; H, 4.75; N, 17.02. (e) M.p. 122°: C, 29.67; H, 4.15; neut. equiv., 207. (f) B.p. 99° (10 mm.): C, 38.08; H, 4.91; N, 17.45. (g) C, 11.77; H, 0.67; N, 18.11; K, 25.50.

(3) The yield is poor because alkaline solutions of nitromethane are rapidly converted to salts of methazonic acid and because the acid, dinitromethane, is unstable.

(4) This also has been observed independently by K. Klager, Aerojet-General Corporation, Azusa, California.

(5) See L. Zeldin and H. Shechter, *J. Am. Chem. Soc.*, **79**, 4708 (1957), and M. B. Frankel, *J. Org. Chem.*, **23**, 813 (1958).

