


Although no mechanistic studies have been undertaken, it is reasonable to postulate a sequence¹⁰ involving (a) Lewis acid-Lewis base interaction of the diazo ketone with the trialkylborane; (b) rapid 1,2-alkyl shift from boron to carbon with simultaneous expulsion of nitrogen; (c) boron-carbon bond fission of the functionalized derivative.



The method is illustrated for the preparation of 2-nonanone. 1-Hexene (60 mmoles) was converted⁷ to trihexylborane by treatment with BH_3 (20 mmoles) in tetrahydrofuran. An azotometer was connected to the reaction vessel. A solution of diazoacetone (20 mmoles) in 15 ml of tetrahydrofuran was added over a period of *ca.* 20 min while the magnetically stirred reaction mixture was kept at 20° by ice cooling. Over 90% of the theoretical amount of nitrogen was evolved at this stage. The mixture was stirred for 30 min at room temperature, then refluxed for 30 min to liberate the remaining nitrogen. The solution was cooled in an ice bath and treated with 20 ml of 3 *N* potassium hydroxide solution. After being stirred at room temperature for 2 hr, the mixture was poured into water and extracted with pentane (three 60-ml portions). Glpc analysis at this stage indicated an 89% yield of 2-nonanone and a trace amount of 1-hexanol. The solution was dried (Drierite), and the residue remaining after removal of solvent was distilled to afford 1.86 g (65%) of 2-nonanone, identical in all respects with an authentic sample.

Currently we are exploring the scope of the reactions of organoboranes with a wide variety of mono- and difunctionally substituted diazoalkanes, N_2CHX and N_2CXY , respectively. Preliminary studies indicate that ethyl diazoacetate reacts with trialkylboranes at a faster rate than does diazoacetone and provides the two-carbon homologated ester. Detailed results of this and related studies will be dealt with in future publications.

Acknowledgment. We wish to thank the National Research Council of Canada for financial support of this work.

(10) Analogous proposals have appeared to account for the reactions of trialkylboranes with ylides¹ and carbanions.² See also ref 3.

(11) Postdoctoral Research Fellow, 1967-1968.

John Hooz, Siegfried Linke¹¹

Department of Chemistry, University of Alberta
Edmonton, Alberta, Canada

Received July 15, 1968

Organic Synthesis by Electrolysis. II. Anodic Methoxylation of Isocyanide


Sir:

Anodic methoxylation of some aromatic compounds or aliphatic unsaturated compounds has been studied by several workers¹ and most of the mechanisms of

(1) S. D. Ross, M. Finkelstein, and R. C. Petersen, *J. Am. Chem. Soc.*, **88**, 4657 (1966), and references cited therein.


these reactions have been elucidated by the formation of a cationic intermediate. On the other hand, the nucleophilic character of isocyanide has been well established² and, furthermore, substantial reactivity of isocyanide toward free radicals has been observed in our previous study.³ Thus, it seemed interesting to study the anodic formation of a cationic species from an isocyanide or the attack of an anodically generated methoxy radical on an isocyanide. In this communication, we wish to report the anodic methoxylation of cyclohexyl isocyanide (I) to yield some unusual products of intriguing synthetic potentialities. In addition, a novel one-step methoxylation reaction of aliphatic ethers is suggested by the present study.

The electrolysis (1 A, 20 V, 20 hr) was carried out with carbon electrodes at the boiling temperature of the solvent. The concentration of I was 1 mole/l. in methanol containing sodium methoxide (0.5 mole/l.). The electrolysis gave seven products (II-VIII) and each product was isolated by fractional distillation or preparative gas chromatography.



Compound II had bp 115° (25 mm); nmr spectrum, τ 8.47 (singlet, 10 H), 6.23 (singlet, 3 H), 5.98 (singlet, 2 H); mol wt (mass spectrum), 169; ir spectrum, 1675 cm^{-1} (C=N). *Anal.* Calcd for $C_9H_{15}NO_2$: C, 63.88; H, 8.94; N, 8.28. Found: C, 63.72; H, 9.13; N, 8.06. The 3-oxazoline (II) was synthesized independently by the route⁴ shown in Scheme I and the structure of II was thereby established.

Scheme I



Compound III had bp 120° (25 mm); nmr spectrum, τ 8.43 (singlet, 10 H), 6.60 (singlet, 3 H), 6.20 (singlet, 3 H), 4.88 (singlet, 1 H); ir spectrum, 1695 cm^{-1} (C=N). *Anal.* Calcd for $C_{10}H_{17}NO_3$: C, 60.28; H, 8.60; N, 7.03. Found: C, 60.13; H, 8.62; N, 7.25. The fact that the electrolysis of II under the same reaction condition as that for I gave III strongly supports the structure proposed for III.

Compound IV had bp 61° (15 mm); nmr spectrum, τ 8.1-8.9 (multiplet, 10 H), 7.0 (broad singlet, 1 H), 6.40 (singlet, 3 H), 2.49 (singlet, 1 H); ir spectrum, 1660

(2) I. Ugi, *Angew. Chem.*, **74**, 9 (1962).

(3) T. Shono, M. Kimura, Y. Ito, K. Nishida, and R. Oda, *Bull. Chem. Soc. Japan*, **37**, 635 (1964).

(4) H. O. L. Fischer, D. Dangschat, and H. Stettiner, *Chem. Ber.*, **65**, 1032 (1932).

