

basic alumina for 16 hr led in high yield to **2** (mp 158–161°), identical with authentic **2** (mp 157–160°). Finally, reduction of **2** with an excess of lithium aluminum hydride in ether at –65° for 90 min²² afforded 50% of **1** (mp 238–241°), identical with natural bufalin (mp 239–242°).

Acknowledgments. We are indebted to the Koninklijke Nederlandsche Gist-en Spritusfabriek N.V. (Delft) for a generous supply of **4**, and to Professors K. Meyer (Basel), G. R. Pettit (Tempe), and T. Reichstein (Basel) for kindly providing samples of natural bufadienolides. We also thank Dr. J. Mathieu (Romainville) for valuable correspondence.

(22) See H. Kondo and S. Ohno, U. S. Patent 3,134,772 (*Chem. Abstr.*, **61**, 5736f (1964)).

(23) Author to whom inquiries may be addressed at the Chemistry Department, University College, London, W.C.1, England.

(24) Honorary NATO Fellow, 1964–1965; Imperial Chemical Industries Fellow, 1964–1966.

(25) Science Research Council Postdoctoral Fellow, 1966–1968.

F. Sondheimer,²³ W. McCrae,²⁴ W. G. Salmond²⁵

University Chemical Laboratory
Cambridge, England

Received December 23, 1968

The Preparation of 2-Methoxypentaborane (9), a Novel Example of an Alkoxy Polyborane

Sir:

We have recently found that 1-iodopentaborane(9), 1-IB₅H₈, reacts with dimethyl ether to produce 2-methoxypentaborane(9), 2-(CH₃O)B₅H₈, in moderate yield. The only other examples of neutral alkoxy polyboranes have the general formula ROB₁₀H₁₃ and are prepared by a complex reaction between NaB₁₀H₁₃ and a solution of I₂ in various ethers.¹ The position of attachment of the alkoxy group is not known, but has been discussed.² A previous study of the reaction of 1-BrB₅H₈ with dimethyl ether resulted in the preparation of 2-BrB₅H₈ and 1-CH₃B₅H₈, but no tractable alkoxy derivatives of B₅H₉ were observed.³

In a typical preparation of 2-(CH₃O)B₅H₈, 1.317 g (6.97 mmoles) of 1-IB₅H₈ was allowed to react with 20.7 mmoles of liquid (CH₃)₂O for 15 hr at –12°. High-vacuum fractional distillation of the pale yellow reaction mixture yielded 0.161 g (1.73 mmoles, 25%) of 2-(CH₃O)B₅H₈. There was 4.9 mmoles of (CH₃)₂O consumed. Substantial quantities of B₅H₉, B(OCH₃)₃, and CH₃I, a relatively small amount of HB(OCH₃)₂, and traces of H₂ were also produced in the reaction. A side product of low volatility has prevented quantitative estimation of the 1-IB₅H₈ recovered from the reaction.

This new B₅H₉ derivative is a colorless liquid which freezes as a glass at –196°. Upon warming the glass begins to flow and then crystallizes. The melting point is in the vicinity of –100°. Representative vapor pressures are 4.4 ± 0.2 mm at 0° and 15.5 ± 0.2 mm at 20°. Moderately rapid decomposition occurs in the liquid phase such that the vapor pressures are of qualitative

value only. In the gas phase at low pressure (~7 mm) no decomposition can be observed after 12 hr at ambient temperature (infrared).

The mass spectrum of 2-MeOB₅H₈, obtained using AEI MS-9 and CEC 21-103 spectrometers, is consistent with that expected for an alkoxy pentaborane(9). The cutoff at *m/e* 94 corresponds to the parent ion ¹²CH₃¹⁶O¹¹B₅H₈⁺: calcd *m/e* 94.12754; found 94.12751 ± 0.00010 (estimated error range). The most intense peak in the spectrum at *m/e* 43 corresponds to ¹²CH₃¹⁶O¹¹BH⁺: calcd *m/e* 43.03552; found 43.03549 ± 0.00010.

The ¹¹B nmr spectrum of 2-(CH₃O)B₅H₈ (at 32.1 MHz) is very similar in appearance to that of 2-FB₅H₈.⁴ The chemical shifts (δ in parts per million from BF₃O-(C₂H₅)₂ ± 0.2), coupling constants (*J* in Hz ± 5), and relative areas are given in Table I. The extreme separation between the B(2) and B(4) resonances suggests that 2-(CH₃O)B₅H₈ is more closely related, electronically, to 2-FB₅H₈ than to any other known B(2)-substituted B₅H₉ derivative.

Table I

	δ	<i>J</i>	Area (rel)
B(2)-OCH ₃	-14.1		1.00
B(3, 5)-H	+16.8	158	1.99
B(4)-H	+31.5	160	0.99
B(1)-H	+55.0	170	1.04

The ¹H nmr spectra of 2-(CH₃O)B₅H₈ (at 60 and 100 MHz) show the presence of two bridge hydrogen regions, as does the spectrum of 2-FB₅H₈.⁴ Overlap of other areas of the spectra makes assignments for H-B_{3, 5}, H-B₄, and H-B₁ ambiguous, but the general appearance is similar to that of 2-FB₅H₈. The methoxy resonance of 2-(CH₃O)B₅H₈ at –3.56 ppm is substantially shifted from that of (MeO)₃B at –3.09 ppm.

The gas-phase infrared spectrum of 2-(CH₃O)B₅H₈ contains major bands (cm⁻¹ ± 10) at 3005 (w), 2960 (w), 2870 (w), 2600 (s), 1985 (w, br), 1850 (w, br), 1475 (m), 1315 (s, br), 1005 (m), 950 (w), 875 (m), 825 (w). The two broad bands at 1985 and 1850 cm⁻¹ are tentatively attributed to the two types of bridge hydrogens indicated in the ¹H nmr spectra. The band at 1475 cm⁻¹ is attributed to the methoxy methyl deformation, and the band at 1005 cm⁻¹ is probably due to a C–O stretch.

Boron trichloride reacts with 2-(CH₃O)B₅H₈ to form what appears to be a 1:1 complex. Solutions of the complex in CS₂ and BCl₃ exhibit ¹¹B nmr spectra similar to those expected for a mixture of 2-ClB₅H₈ and ROBCl₂ (which appears to undergo rapid exchange with excess BCl₃). Subsequent isolation of 2-ClB₅H₈, however, is not possible unless the mixture is heated to ~50°. Further studies of this behavior are in progress.

Acknowledgments. Support of this research by the National Science Foundation is gratefully acknowledged. We wish to thank Bill Stebbings for the exact *m/e* mass spectral determinations.

(4) A. B. Burg, *ibid.*, **90**, 1407 (1968).

Donald F. Gaines

Department of Chemistry, University of Wisconsin
Madison, Wisconsin 53706

Received November 30, 1968

(1) M. F. Hawthorne and J. J. Miller, *J. Am. Chem. Soc.*, **82**, 500 (1960).

(2) R. E. Williams, *Inorg. Chem.*, **4**, 1504 (1965).

(3) A. B. Burg and J. S. Sandhu, *J. Am. Chem. Soc.*, **87**, 3787 (1965).