


an active site residue. The exact nature of the linkage is now under investigation.

Studies were conducted with propargylglycine in intact mice. Groups of three mice were injected intraperitoneally with 2 and 5 μmol of D,L-propargylglycine, respectively, while three control mice received the same volume of saline solution. After 24 hr, the mice were sacrificed, and the livers were removed and assayed for γ -cystathionase. The livers from the treated mice had 10–20 and 0%, respectively, of the enzyme levels from control livers. Propargylglycine inactivation of the enzyme *in vivo* indicates induction of a condition similar to that found in the genetic defect cystathionuria¹⁰ in which liver γ -cystathionase is absent or defective.

An essential feature of the mechanism of action of γ -cystathionase is the abstraction of a proton from the β position.⁶ We, therefore, tentatively propose the mechanism in Scheme I for the inactivation by pro-

Scheme I



pargylglycine. The allene formed after proton abstraction is in conjugation with the ketimine and should be capable of ready Michael addition by an enzyme active site nucleophile to produce covalent labeling of the active site (III). (B_1 and B_2 are active site basic groups and PyCHO is pyridoxal-P.)

In the proposed mechanism, the acetylenic linkage is essential. This is consistent with finding that allylglycine (Sigma Chemical Co.) (2-amino-4-pentenoic acid) does not inactivate γ -cystathionase. Also, only those pyridoxal-dependent enzymes which abstract substrate β -hydrogens should catalyze their own destruction by propargylglycine. Preliminary experiments¹¹ with threonine deaminase and transaminase show no inactivation by propargylglycine.

In summary, then, we have shown for the first time that an acetylenic amino acid will irreversibly inactivate a pyridoxal-P dependent enzyme, rat liver γ -cystathionase, *in vitro* and *in vivo*, presumably *via* a reactive allene intermediate. This extends our recent observations that several flavine coenzyme-dependent enzymes^{3,12} are irreversibly inactivated by acetylenic substrates. It confirms the expectation that, in general, acetylenic substrate analogs should be potent and

(10) G. W. Frimpter, *Science*, **149**, 1095 (1965); F. C. Brown and P. H. Gordon, *Biochim. Biophys. Acta*, **230**, 434 (1971).

(11) W. Washtien and R. H. Abeles, unpublished experiments.

(12) C. T. Walsh, R. H. Abeles, and H. R. Kaback, *J. Biol. Chem.*, **247**, 7858 (1972).

specific irreversible inactivators in enzymatic catalyses where the carbon-bound hydrogen adjacent to the acetylenic moiety is abstracted as a proton.

Acknowledgments. Acknowledgment is made to the donors of the Petroleum Research Fund (C. T. W.), administered by the American Chemical Society, for partial support of this work, and to the National Institutes of Health (Grant No. GM 12633-10) and Massachusetts Institute of Technology Chemistry Department Research Funds (C. T. W.). We also wish to acknowledge the competent technical assistance of Miss Donna Ozog and Mrs. Carole Foxman.

Robert H. Abeles*

Graduate Department of Biochemistry, Brandeis University
Waltham, Massachusetts 02154

Christopher T. Walsh*

Departments of Chemistry and Biology
Massachusetts Institute of Technology
Cambridge, Massachusetts 02139

Received May 30, 1973

Highly Specific Enzyme Inhibitors. Inhibition of Plasma Amine Oxidase


Sir:

We have recently suggested¹ that substrate analogs containing an acetylenic group adjacent to a carbon-bound hydrogen should be generally useful as enzyme inactivators when this hydrogen is abstracted as a proton during catalysis.

Based upon earlier work² from the laboratory of Bloch, it was suggested that this inactivation proceeds through rearrangement of the acetylene to the allene which then reacts with a nucleophile at the active site. Results obtained in our studies with beef plasma amine oxidase³ suggest that an early step in the oxidation involves proton abstraction⁴ from the carbon atom which is oxidized. Therefore, this enzyme might be susceptible to inactivation by appropriate acetylenic substrates.

As reported below, we have found that 1-amino-2-alkynes are capable of inactivating plasma amine oxidase. Furthermore, if the proposed mechanism of inactivation is correct, other substrates which can form allenes during the catalytic process should also inactivate the enzyme.

2-Chloroallylamine (I), which could undergo allene



formation by expulsion of chloride after the C-1 hydrogen is abstracted as a proton, is found to inactivate the enzyme irreversibly.

(1) R. H. Abeles and C. T. Walsh, *J. Amer. Chem. Soc.*, **95**, 6124 (1973).

(2) M. Morisaki and K. Bloch, *Biochemistry*, **11**, 309 (1972).

(3) Plasma amine oxidase is a nonflavoprotein oxidase, which catalyzes the oxidation of certain primary amines according to the equation $\text{RCH}_2\text{NH}_2 + \text{O}_2 + \text{H}_2\text{O} \rightarrow \text{RCHO} + \text{H}_2\text{O}_2 + \text{NH}_3$. The enzyme is believed to contain Cu^{II} and pyridoxal phosphate: H. Yamada and K. T. Yasunobu, *J. Biol. Chem.*, **237**, 3077 (1962); **238**, 2669 (1963). We feel, however, that the presence of pyridoxal phosphate is not firmly established. The enzyme used in these studies was purified and assayed according to the procedure of H. Yamada and K. T. Yasunobu, *J. Biol. Chem.*, **237**, 1511 (1962).

(4) R. Hevey and R. H. Abeles, unpublished results.