

measurements of  $1/\tau_2$  were carried out on a Durrum stopped-flow spectrophotometer. Determinations of  $1/\tau_1$  and initial rate measurements were carried out on a Gilford Model 2000 spectrophotometer. All relaxation times reported in Tables I, II, and IV represent averages of three (in a few cases two) determinations.

**Acknowledgment.** This work was supported by a grant from the Petroleum Research Fund administered by the American Chemical Society. Grateful acknowledgment is made to the donors of this fund.

## The Kinetics of Carbonium Ion Formation from Anhydroretinol

Vernon C. Bulgrin\* and George L. Lookhart

*Contribution from the Department of Chemistry, University of Wyoming, Laramie, Wyoming 82071. Received February 22, 1974*

**Abstract:** The initial stages of the reaction between anhydroretinol and anhydrous hydrogen chloride in 1,2-dichloroethane were studied using stopped-flow spectrophotometry. The results are explained in terms of a five-step mechanism, involving two carbonium ions, with the anhydroretinylic cation forming much more rapidly than the more stable retinylic cation. Evidence is presented indicating that the former cation may be a precursor of the latter.

Highly conjugated polyenes when treated with acid generally give rise to absorption maxima in the visible region. These absorptions have been attributed to charge transfer or to carbonium ion formation. Wasserman<sup>1</sup> showed by conductance measurements that, in the case of protonic acids, carbonium ions are present, and presumably are responsible for the color observed.

The carbonium ions from *all-trans*-retinol and related compounds have been of particular interest because their spectral properties are similar to those of visual pigments, and they may be meaningful as models of visual pigments. Blatz and Pippert<sup>2</sup> showed the retinylic cation (Figure 1,II) to be the only cation observed on protonating *all-trans*-retinyl acetate or retinol; their results also demonstrated that  $\lambda_{\text{max}}$  of the cation is only a weak function of solvent and independent of the nature and concentration of the protonic acid. Protonation of anhydroretinol (I) gave rise to a longer wavelength absorption, which was assigned to the anhydroretinylic cation (III) by Blatz on the basis of protonation at the C<sub>15</sub> position.<sup>3</sup> (Protonation at the other most likely point of attack, C<sub>4</sub>, would give rise to the retinylic cation.) Symmetry considerations also favored the assignment of the longer wavelength absorption to the anhydroretinylic cation. Baumgartner<sup>4</sup> observed both cations to form when anhydroretinol is protonated by HCl in 1,2-dichloroethane. It could not be determined which of the cations formed first.

A kinetic study of the rate of formation of carbonium ions from anhydroretinol was undertaken, in order to establish the mechanism of the reaction, to ascertain the relative rates of formation of the cations, to indicate the relative ease of protonation at the C<sub>4</sub> and C<sub>15</sub> positions of anhydroretinol, and to give some indication as to the relative stabilities of the two cations.

The reaction was studied at 10.0, 21.2, and 30.0°, using HCl as protonating agent in 1,2-dichloroethane solvent. Only the initial (up to several minutes) portion of the reaction was studied in detail, since it has been shown by Sorenson,<sup>5</sup> Deno,<sup>6</sup> and Blatz<sup>7</sup> that cyclization and polymerization of such carbonium ions take place at longer times. In all of the studies, a very large excess of HCl was used in order to achieve first-order kinetics. The concentration of HCl was varied to obtain the HCl dependency.

### Results and Discussion

Typical action spectra taken on the Cary 14R at -30° are shown in Figure 2. During the time period of these observations (0.5-4 min), it appears that the first-formed anhydroretinylic cation (~620 nm) is decreasing slightly with time, while the retinylic cation (~580 nm) is increasing.

Typical reaction profiles from the stopped-flow data are shown in Figure 3, which also illustrates the effect of temperature. The data are plotted as absorbance vs. log time for convenience. Allowing for some overlap in the spectra of the two cations, there appear to be four regions in time corresponding to four rate processes. These are labeled  $\phi_1$ ,  $\phi_2$ ,  $\phi_d$ , and  $\phi_3$ , with overlap in rates occurring between  $\phi_2$  and  $\phi_d$ .  $\phi_2$  and  $\phi_3$  were assigned to the formation of anhydroretinylic and retinylic cations, respectively,  $\phi_1$  to the formation of a precursor which absorbs rather broadly over the 580-620-nm region, and  $\phi_d$  to a process involving the disappearance of the anhydroretinylic cation. The  $\phi_2$ ,  $\phi_d$ , and  $\phi_3$  assignments were made after comparing the stopped-flow data with the Cary 14 action spectra. The time scale of the action spectra suggests a correspondence with processes occurring in the  $\phi_d$ - $\phi_3$  region of the stopped-flow data. The  $\phi_1$  assignment, that of a reaction leading to a precursor to the carbonium ions,

(1) A. Wasserman, *J. Chem. Soc.*, 245 (1954).

(2) P. E. Blatz and D. L. Pippert, *J. Amer. Chem. Soc.*, **90**, 1296 (1968).

(3) The numbering system is that used in the carotenoid literature.

(4) N. R. Baumgartner, Ph.D. Dissertation, University of Wyoming, 1969.

(5) T. S. Sorenson, *Can. J. Chem.*, **43**, 2744 (1965).

(6) N. C. Deno, C. U. Pittman, and J. O. Turner, *J. Amer. Chem. Soc.*, **87**, 2153 (1965).

(7) P. E. Blatz and D. L. Pippert, *Tetrahedron Lett.*, 1117 (1966).




Figure 1. Carbonium ions from anhydroretinol.





Figure 2. Action spectra, Cary 14R,  $-30^\circ$ . Anhydroretinol concentration,  $8 \times 10^{-4} M$ ; HCl concentration, about 0.01 M. Curves A, B, C, and D taken at 0.5, 1, 2, and 4 min, respectively.

was more difficult, but was made on the basis that the process was observable at both 580 and 620 nm with identical rate constants within the experimental error of about  $\pm 10\%$ . A detailed analysis of each region was then made.

**$\phi_1$  Region.** The time scale of the  $\phi_1$  reaction was such as to push the stopped-flow instrument to the limit of its capabilities. Nevertheless, good first-order plots were obtained from both the 580- and 620-nm data, with the first-order rate constants at the two wavelengths agreeing at each temperature and HCl concentration to within 10% or less. Plots of reciprocal rate constants  $\phi_1$  vs. reciprocal HCl concentration are linear (Figure 4), consistent with a two-step process



where A is anhydroretinol, C an intermediate of unknown structure, and AHCl an anhydroretinol-HCl species absorbing rather weakly and broadly over the 580-620-nm region. It is assumed that in the AHCl species, taken to be the immediate precursor to the anhydroretinylic cation, the HCl molecule is attached at the  $C_{15}$  end of the anhydroretinol molecule, perhaps as a  $\pi$  complex as discussed later. In the mechanism above,  $k_1$  is the reciprocal of the intercepts in Figure 4; the ratio of the intercept to the slope gives  $K$ . Values of  $k_1$  and  $K$  are given later.

**$\phi_2$ - $\phi_d$  Region.** First-order rate constants were de-




Figure 3. Reaction profile, stopped-flow data. Anhydroretinol concentration,  $7.45 \times 10^{-4} M$ ; HCl concentration, 0.0144 M.




Figure 4. Reciprocal first-order rate constant,  $\phi_1$  vs. reciprocal HCl. Intercepts are  $k_1^{-1}$ ; slopes/intercepts are  $K$ .

termined from the 620-nm data in this region, where overlap of two rate processes occurs. The method for separating the two rate constants  $\phi_2$  and  $\phi_d$  is well known. It involves obtaining  $\phi_d$  (the smaller rate constant) from the slope of a plot of  $\ln(A - A_\infty)$  vs. time at longer times as shown in Figure 5a. Here,  $A$  is the observed absorbance at time  $t$  and  $A_\infty$  the absorbance (very nearly constant) at the completion of the  $\phi_d$  process. The straight line is extrapolated back to zero time, and values of  $A_{\text{calcd}}$  were obtained for the early ( $\phi_2$ ) part of the region. Plots of  $\ln(A - A_{\text{calcd}})$  vs. time are linear, as shown in Figure 5b, with slopes equal to negative  $\phi_2$ .

Figure 6 shows plots of  $\phi_2$  vs. the concentration of HCl at the three temperatures studied, demonstrating a first-order HCl dependency. The slope of each line is  $k_2$ , the rate constant for the reaction of the AHCl complex with HCl to give the anhydroretinylic cation. In the solvent used the anhydroretinylic cation probably exists as an ion pair with chloride, *i.e.*,  $\text{AH}^+\text{Cl}^-$ . Thus, the formation of the anhydroretinol cation from anhydroretinol requires two HCl molecules. This is in




Figure 5. First-order plots in  $\phi_2$ - $\phi_d$  region. Figure 5b corresponds to first 5 min of 5a.

agreement with the observation that in hydrohalogenation of olefins,<sup>8</sup> involving formation of carbonium ions, two molecules of HCl are required to form the carbonium ion intermediate, the first leading to the formation of an HCl  $\pi$  complex to the olefin (as in the  $\phi_1$  step of this reaction), the second removing the  $\text{Cl}^-$  ion as (presumably)  $\text{HCl}_2^-$  (as in the  $\phi_2$  step of this reaction). In the anhydroretinol reaction, however, it is possible to observe, spectroscopically, the formation of the carbonium ion, because of the conjugated polyene system leading to an absorption in the visible region.

The  $\phi_d$  rate constants obtained were found to be nearly independent of HCl concentration. For example, although there appeared to be a slight increase in  $\phi_d$  with HCl, at 30°, for example, the average deviation of  $\phi_d$  for all five HCl concentrations over a three-fold change was only  $\pm 5\%$ . Therefore,  $\phi_d$  was taken to be  $k_d$ , the HCl-independent rate constant for the disappearance of the anhydroretinylic cation, possibly *via* cyclization or polymerization as mentioned earlier.

It appears that a suitable mechanism describing the  $\phi_1$ ,  $\phi_2$ , and  $\phi_d$  regions is



where P is a cyclization or polymerization product. Evidence for the reversibility of the last step is presented later.

**$\phi_3$  Region.** A relatively slow zero-order process was found from the 580-nm data in this region, plots of absorbance *vs.* time giving very good straight lines. Figure 7 shows plots of  $\phi_3$  *vs.* the hydrogen chloride concentration, the resulting straight lines indicating a first-order dependency. The slope of each line is  $k_3$ , the rate constant for the formation of the retinylic cation (III). Zero-order kinetics under the conditions of the experiment (large excess of HCl) would be expected if the precursor of the retinylic cation were in very nearly a steady-state concentration, or if the process were photochemical. The latter was ruled out by the observation that the rate of growth of the 580-nm peak was the same whether irradiation in the stopped-flow

(8) F. R. Mayo and C. Walling, *Chem. Rev.*, **27**, 351 (1940).




Figure 6. Plot showing first-order HCl dependency for  $\phi_2$ .





Figure 7. Plot showing first-order HCl dependency for  $\phi_3$ .

spectrophotometer was continuous or intermittent. An approximate steady state in the anhydroretinylic cation in this region is indicated by the Cary action spectra (Figure 2). Therefore, one conclusion is that the retinylic cation is formed by attack of HCl at the  $\text{C}_4$  position of the anhydroretinylic cation, with subsequent or simultaneous elimination of a proton from  $\text{C}_{15}$ . Alternatively, the precursor to the retinylic cation could be an earlier intermediate such as C or AHCl. No way has been found to distinguish between these alternatives, although the latter would seem to require a higher degree of reversibility in the mechanism than is observed. A third possibility is that the precursor is the unknown reaction product P of the anhydroretinylic cation.

A mechanism which is consistent with the experimental observations is



The constants for this mechanism, as well as the thermodynamic parameters, are given in Table I.

Why the retinylic cation appears to be formed primarily by HCl attack on the anhydroretinylic cation rather than on anhydroretinol is not understood at this time. The latter case would be expected to give first-order kinetics rather than the zero order observed.

The necessity for reversibility of the  $k_d$  step is indicated from the observed near steady state in  $\text{AH}^+\text{Cl}^-$  and the accompanying zero-order formation of  $\text{RH}^+\text{Cl}^-$ .

**Table I.** Constants<sup>a</sup> and Thermodynamic Parameters<sup>b</sup> for the Mechanism in Eq 1

Con- stant <sup>a</sup>	Temp, °C			$\Delta H^\circ$ , kcal	$\Delta S^\circ$ , cal mol <sup>-1</sup> deg <sup>-1</sup>
	10.0	21.2	30.0		
<i>K</i>	250	90	39	-19.3	-57
<i>k</i> <sub>1</sub>	260	640	1500	15.7	85
<i>k</i> <sub>2</sub>	15.7	18.4	21.1	4.9	43
<i>k</i> <sub>d</sub> + <i>k</i> <sub>-d</sub>	0.026	0.045	0.054	6.3	35
<i>k</i> <sub>3</sub> <sup>c</sup>	0.0074	0.0154	0.0210	9.8	44

<sup>a</sup> *K* in l. mol<sup>-1</sup>; *k*<sub>1</sub> and *k*<sub>d</sub> in sec<sup>-1</sup>; *k*<sub>2</sub> and *k*<sub>3</sub> in M<sup>-1</sup> sec<sup>-1</sup>.

<sup>b</sup> All ln (*K* or *k*) vs. *T*<sup>-1</sup> plots gave good straight lines except for the  $\phi_d (= k_d + k_{-d})$  plot which showed curvature, as might be expected for the sum of two rate constants. <sup>c</sup> Thermodynamic parameters for *k*<sub>3</sub>, although listed here, have no absolute significance in that the *k*<sub>3</sub> determined is a product of a second-order rate constant times an extinction coefficient times a cell path length times the concentration of a steady-state precursor.

It is also likely that the *k*<sub>1</sub>, *k*<sub>2</sub>, and *k*<sub>3</sub> steps are to some degree reversible. The major evidence in support of this is the fact that anhydroretinol is prepared (albeit in very small yields) by action of HCl on retinol or retinyl acetate. Preliminary stopped-flow experiments following the protonation of retinol indicate that only one carbonium ion, the retinyl cation, is present in observable quantity early in the reaction; Cary data, however, indicate a small amount of the anhydroretinyl cation is present in the later stages. Thus, it appears that the retinyl cation is capable of being converted to anhydroretinol, probably *via* the anhydroretinyl cation. The stopped-flow data on retinol cannot directly confirm this, however, because of the difficulty of observing a small amount of anhydroretinyl cation (620 nm) in the presence of a very large amount of retinyl cation (590 nm), due to the (small) amount of overlap in the peaks.

The zero-order formation of the retinyl cation, coupled with the near steady-state in anhydroretinyl cation in this ( $\phi_3$ ) time region, seems to be rather compelling evidence that the anhydroretinyl cation is a necessary intermediate in the formation of retinyl cation from anhydroretinol. That two molecules of HCl are required to form the anhydroretinyl cation is consistent with the generally accepted mechanism of hydrohalogenation of olefins. Other aspects of the reaction are more difficult to explain, for example, the nature of the intermediate C, and the apparent lack of direct protonation of anhydroretinol to give retinyl cation.

It is concluded, however, that direct protonation of anhydroretinol at the C<sub>15</sub> position is kinetically favored over protonation at C<sub>4</sub>. Conversely, the retinyl cation is ultimately formed in much larger quantity than the anhydroretinyl, and therefore appears to be more stable than the latter. The mechanism proposed also explains why the yields of anhydroretinol obtained on protonating retinol or retinyl acetate are always very low.

## Experimental Section

**Polyenes.** *all-trans*-Retinyl acetate was obtained in crystalline

form from Sigma Chemical Co. in vacuum-sealed 100-g ampoules. After opening the ampoules, the crystals were dried under vacuum for 3 hr and stored under dry nitrogen at -30° in total darkness. *all-trans*-Anhydroretinol was prepared from retinyl acetate by a modification of the procedure of Shantz;<sup>9</sup> 10 g of retinyl acetate was dissolved in 500 ml of anhydrous ethanol, and 250 ml of an approximately 3% solution of HCl in anhydrous ethanol was added. The reaction mixture was neutralized after 5 min with a saturated aqueous solution of sodium bicarbonate. This time of reaction was selected, by monitoring the uv spectrum at 1-min intervals, as that giving the maximum concentration of the desired material. The mixture was extracted with low-boiling petroleum ether; the extracts were combined, washed with water, dried over anhydrous MgSO<sub>4</sub>, and reduced in volume. This was followed by twice chromatographing under N<sub>2</sub> in dim red light on a 4 × 30 cm column packed with 6% water-deactivated Woelm neutral alumina, using petroleum ether as eluent. The second band through the column was the *all-trans*-anhydroretinol. The first and third (minor) bands have been ascribed to cis isomers.<sup>10</sup> The fractions with the proper relative peak heights<sup>11</sup> were combined, reduced in volume (under dry nitrogen) to near oil, and stored in total darkness at -30°. After 3 days, crystallization was complete; the yellow-orange crystals were dissolved in a small amount of petroleum ether; after reduction in volume, crystallization was complete in 24 hr at -30°. The crystals were rinsed with a small amount of very cold petroleum ether, dried under vacuum for 8 hr, and stored under dry nitrogen in the dark at -30°. The yield was 500 mg of yellow-orange crystals melting at 75.5-76.5°. Solutions of the crystals in 1,2-dichloroethane showed the characteristic triple peaks at 356, 375, and 398 nm, with relative peak heights of 0.68, 1.00, and 0.90, respectively. The molar absorptivity for the 375-nm peak was found to be 7.83 × 10<sup>4</sup> (±0.53, average deviation for 16 preparations). Orosnik<sup>12</sup> reported a value of 7.8 × 10<sup>4</sup> in ethanol. Solutions of anhydroretinol in 1,2-dichloroethane were found to be stable for up to a week if stored in the dark under dry nitrogen at -30°.

**Solvents.** 1,2-Dichloroethane was washed several times with concentrated sulfuric acid, then with water, a saturated solution of sodium bicarbonate, and water again. After successively drying with anhydrous MgSO<sub>4</sub> and P<sub>2</sub>O<sub>5</sub>, it was distilled under dry nitrogen from P<sub>2</sub>O<sub>5</sub>. Petroleum ether was purified in a similar manner. The solvents, after purification, were stored in glass under dry nitrogen.

**Hydrogen Chloride Solutions.** Solutions of hydrogen chloride in 1,2-dichloroethane were prepared by passing anhydrous HCl from Matheson Gas Products through Drierite and then through the solvent. The solution was analyzed titrimetrically and used as a stock solution for preparing the standard HCl solutions used in the rate determinations.

**Action Spectra.** The action spectra shown in Figure 3 were obtained on the Cary 14R in the following way. Two milliliters of an 8.5 × 10<sup>-4</sup> M solution of anhydroretinol in 1,2-dichloroethane was placed in one of two matched 10-mm Suprasil quartz cells, pure solvent in the other. The cells were cooled to -30° and a small amount of HCl solution (constituting a large excess of HCl) was added to the first. The spectrum was rapidly scanned repetitively until the peaks began to disappear.

**Rate Determinations.** The rate of the reaction was studied using a Durrum-Gibson D-110 stopped-flow spectrophotometer, equipped with a 20-mm path length Kel-F cell and Kel-F flow components. Temperature was maintained at 10.0, 21.2, and 30.0° to within ±0.1° by means of a thermostated bath, using anhydrous ethanol as circulating liquid. The flow system, including all syringes, was thoroughly rinsed with dry acetone, anhydrous ethanol, and purified anhydrous 1,2-dichloroethane. The concentration of anhydroretinol used was 7.45 × 10<sup>-4</sup> M (after mixing in the stopped-flow apparatus); concentrations of HCl used (after mixing) were 0.0048, 0.0072, 0.0096, 0.0120, and 0.0144 M. For each concentration at each temperature, the reaction was followed at 580 nm and repeated at 620 nm, corresponding to the absorption peaks of the retinyl and anhydroretinyl cations, respectively. All runs were made in dim red light.

(9) E. M. Shantz, J. D. Cawley, and N. D. Embree, *J. Amer. Chem. Soc.*, **65**, 901 (1943).

(10) R. C. Christensen and B. E. Kohler, *Photochem. Photobiol.*, **18**, 293 (1973).

(11) E. M. Shantz, *J. Biol. Chem.*, **182**, 515 (1950).

(12) W. Orosnik, G. Karmes, and A. D. Mebane, *J. Amer. Chem. Soc.*, **74**, 295 (1952).