

Protonation Rate Constants and Optical Absorption Spectra of Benzyl Carbanion Pairs in Tetrahydrofuran Solution

Bradley Bockrath and Leon M. Dorfman*

Contribution from the Department of Chemistry, The Ohio State University, Columbus, Ohio 43210. Received September 26, 1974

Abstract: The benzyl carbanion paired with lithium cation and with tetrabutylammonium ion has been formed in tetrahydrofuran solution and investigated by the pulse radiolysis method. Absolute rate constants for the protonation of both of these species by methanol and by *tert*-butyl alcohol have been determined. These data, and previous data for benzyl sodium, show that the ion paired species is more reactive, in these protonations, than is the free carbanion. The optical absorption spectra of both $\text{PhCH}_2^-\text{Li}^+$ and $\text{PhCH}_2^-\text{NBu}_4^+$ were determined. The latter exhibits a band maximum at 342 nm, the former, in agreement with data from equilibrium studies, at 330 nm. The effect of ion pairing upon the absorption spectrum of PhCH_2^- by Li^+ , Na^+ , and NBu_4^+ is shown.

Effects of cation pairing upon the reactivity of aromatic radical anions²⁻⁶ and carbanions⁷⁻⁹ in solution have been demonstrated. In such studies both enhancement of rates and reduction of rates as a result of cation pairing have been found. In our recent report⁷ of the formation and observation of benzyl carbanion on a submicrosecond time scale by the pulse radiolysis method,^{10,11} it has been shown how such studies may be extended to highly reactive carbanions. With this method the carbanions may be generated in either the unpaired state or in a cation paired state. In this report we extend these fast reaction studies of the benzyl carbanion, a relatively simple carbanion of particular interest in organic chemistry, to cover the effects of ion pairing with lithium cation and with tetrabutylammonium cation. These effects of cation pairing are concerned with changes in the optical absorption spectrum as well as with the reactivity of the species in elementary reactions. The particular reactions under investigation are protonation reactions with aliphatic alcohols.


Experimental Section

The source of the electron pulse, as in our earlier studies,¹² was a Varian type V-7715A electron linear accelerator, delivering 3-4 MeV electrons at a pulse current of about 300 mA for pulse duration of 100-1500 nsec and about 600 mA for pulse duration less than 80 nsec. The transient optical absorptions were observed using an H.T.V. 196 detector with an S-1 response or an R.C.A. 1P28 with an S-5 response. The 10-90% rise time of these electronic detection systems was less than 10 nsec. A Bausch and Lomb grating monochromator, type 33-86-26, $f/3.5$ was used. Corning filters were selected to eliminate second-order components from the analyzing light beam. Our standard reaction cells,¹² with high-purity silica windows and a cell length of 20.0 mm, were used with a double pass of the analyzing light beam.

Tetrahydrofuran, dibenzylmercury, methanol, and *tert*-butyl alcohol all were purified as previously described.⁷ Lithium perchlorate (Alfa Products, 99.5%) and tetrabutylammonium perchlorate (Eastman) were used as supplied after thorough drying under vacuum.

Results and Discussion

The formation of benzyl carbanion by electron irradiation of appropriate solutions occurs, as we have shown,⁷ when attachment of the solvated electron produces a dissociative ionization, as with dibenzylmercury in tetrahydrofuran.





The rate constant⁷ for this reaction is $2.7 \times 10^{10} \text{ M}^{-1} \text{ sec}^{-1}$. Since this dissociation is extremely rapid, the use of an electron pulse with duration of 100 nsec or less thus pro-

vides a "pulse" of benzyl carbanion. The carbanion may thus be formed in the absence of any pairing counterion or in the presence of selected cations by introducing, into the solution, dissociative salts. In the presence of a dissociative salt, M^+X^- , the corresponding ion pair of the carbanion may be formed in either of two ways:



or



The particular path will depend upon the ratio $[\text{M}^+]/[(\text{PhCH}_2)_2\text{Hg}]$.

In the present study, the ion pairs of benzyl carbanion with lithium ion and with tetrabutylammonium ion have been formed by using LiClO_4 and NBu_4ClO_4 as the dissociative salt. In previous work¹³⁻¹⁶ we have reported the absorption spectra of the primary reducing species in its various forms in THF, namely e_s^- , (Na^+, e_s^-) , and (Li^+, e_s^-) , as well as the values of the absolute rate constant, k_5 , for both Na^+ ($7.9 \times 10^9 \text{ M}^{-1} \text{ sec}^{-1}$) and Li^+ ($1.8 \times 10^{10} \text{ M}^{-1} \text{ sec}^{-1}$). There has been no evidence presented as yet that a process analogous to reaction 4 proceeds with NBu_4^+ in THF. In our system, the formation of $\text{PhCH}_2^-\text{NBu}_4^+$, for which we present evidence, occurs exclusively by the sequence of reactions 1 and 3.

The transient optical absorption spectra observed at the end of a pulse in solutions of dibenzylmercury and either LiClO_4 or NBu_4ClO_4 are shown in Figures 1 and 2, respectively. The following evidence supports the identification of these absorption bands with the respective ion pairs, $\text{PhCH}_2^-\text{Li}^+$ (the spectrum of which is known in stable solutions) and $\text{PhCH}_2^-\text{NBu}_4^+$. The perchlorate salt concentration in these experiments was made great enough so that, whichever mechanism applies, the formation of the ion pair was complete by the end of the pulse. In the case of NBu_4^+ , the ion pairing occurred by reaction 3. In the case of Li^+ , the ion pair formation occurred by way of the concurrent reactions 3 and 5. No continued formation of the ion pair was observed after the electron pulse.

The spectrum of $\text{PhCH}_2^-\text{Li}^+$, shown in Figure 1, was derived from rate curves observed at various wavelengths following the irradiation of dibenzylmercury solutions with LiClO_4 (0.03 *F*) present. In these solutions, which also contained either methanol or *tert*-butyl alcohol as carbanion scavengers, the absorption decayed in accord with a first-




Figure 1. Normalized absorption spectrum of benzyllithium in THF obtained from decay curves following irradiation of dibenzylmercury solutions with LiClO_4 (0.03 F) present. The difference was taken between the absorption at $t = 0$ after the pulse and at $t = 1 \mu\text{sec}$, when the protonation of the carbanion by the scavenger alcohol was complete: (O) methanol ($7.5 \times 10^{-3} M$); (●) *tert*-butyl alcohol ($2.0 \times 10^{-3} M$). From the combined data, the wavelength of maximum absorption is 330 nm, in agreement with the value¹⁷ obtained in stable solutions.




Figure 2. Normalized absorption spectrum of $\text{PhCH}_2\text{-NBu}_4^+$ in THF obtained from rate curves following irradiation of dibenzylmercury solutions with both NBu_4ClO_4 (0.01 to 0.02 F) and a scavenger alcohol present: (O) methanol ($2.4 \times 10^{-3} M$); (●) *tert*-butyl alcohol ($5.2 \times 10^{-3} M$). From these data, the wavelength of maximum absorption is 342 nm.

order rate law, in less than $1 \mu\text{sec}$ to form a plateau. The decay is due to the protonation of the carbanion by the alcohol. The absorption plateau is undoubtedly due to radiolysis products other than the carbanion which decay with a much longer time constant than that observed for the foregoing protonation reaction. The absorption due only to the carbanion was determined by subtracting the absorbance values after the plateau is established from those at $t = 0$ after the pulse. The combined data from two separate experiments, using either alcohol, define an absorption band with a maximum at 330 nm which is undoubtedly due to $\text{PhCH}_2\text{-Li}^+$ since the absorption maximum coincides with that reported for benzyllithium from studies¹⁷ in equilibrium systems. The potential in our method of formation lies in the ability to produce general ion pairs, R^-Li^+ , in the *monomeric* form for the study not only of the spectra but of the absolute reactivity.

The transient absorption spectrum shown in Figure 2 was obtained from dibenzylmercury solutions with NBu_4ClO_4

Table I. Optical Absorption Band Maxima for the Free Carbanion and for Ion Pairs, with Various Cations, of Benzyl and Fluorenyl Ion in THF

Benzyl carbanion		Fluorenyl ion ^b	
Ion or ion pair	λ_{max} , nm	Ion or ion pair	λ_{max} , nm
PhCH_2^-	362 ^a	Fl^-	374
$\text{PhCH}_2^-\text{Na}^+$	355 ^c	$(\text{Fl}^-, \text{M}^+)_s$	373 ^d
$\text{PhCH}_2^-\text{NBu}_4^+$	342	$\text{Fl}^-\text{NBu}_4^+$	368
$\text{PhCH}_2^-\text{Li}^+$	330	Fl^-Na^+	356
		Fl^-Li^+	349

^a Reference 7. ^b Reference 18. ^c Reference 19. ^d $(\text{Fl}^-, \text{M}^+)_s$ indicates a solvent separated ion pair for $\text{M}^+ = \text{Li}^+$ or Na^+ , produced at slightly reduced temperatures.

Table II. Rate Constants ($M^{-1} \text{sec}^{-1}$) for the Protonation of the Benzyl Carbanion in Its Various Forms by Aliphatic Alcohols in THF at 24°

Benzyl carbanion	CH_3OH	$t\text{-C}_4\text{H}_9\text{OH}$
$\text{PhCH}_2^-\text{Na}^+$ ^a	5.8×10^9	1.3×10^9
$\text{PhCH}_2^-\text{NBu}_4^+$	6.0×10^8	4.6×10^8
$\text{PhCH}_2^-\text{Li}^+$	3.4×10^8	9.7×10^7
PhCH_2^- ^a	2.3×10^8	1.6×10^7

^a Taken from ref 7.

(0.01 to 0.02 F) present, in a manner similar to that described above using alcohol scavengers. The absorption maximum of the band is at 342 nm. On the basis of the manner of formation of this transient absorption and its reactivity toward the proton donors, we suggest that the species is $\text{PhCH}_2\text{-NBu}_4^+$. The absorption is considerably removed from that for PhCH_2^- (362 nm) and is different from the absorption of either $\text{PhCH}_2\text{-Li}^+$ or $\text{PhCH}_2\text{-Na}^+$. Different absorption spectra, depending upon the particular cation involved in ion pairing, have been found in earlier work with other carbanions.¹⁸ The shift upon formation of the lithium ion pair is the largest observed among the three coupling cations, Na^+ , Li^+ , and NBu_4^+ . Our spectra, although complete only on the low-energy side because of cut-off of the transmission by dibenzylmercury, show that pairing with Li^+ narrows the band considerably. The half-width (on the low-energy side) at half-height for $\text{PhCH}_2\text{-Li}^+$ is about 2000 cm^{-1} , while for PhCH_2^- it is about 4000 cm^{-1} . Ion pairing with Na^+ also narrows the absorption band, but not to as great an extent as with Li^+ . Pairing with NBu_4^+ on the other hand, produces no discernible narrowing of the band.

Table I contains the absorption maxima of the three benzyl ion pairs and the free benzyl carbanion, along with similar data which have been reported for fluorenyl¹⁸ ion pairs. It is noteworthy that, with the addition of our new data for $\text{PhCH}_2\text{-NBu}_4^+$ to this series, a somewhat different order is found for the benzyl ion pairs than for the fluorenyl ion pairs. Sodium ion pairing apparently has a greater relative effect upon λ_{max} for the Fl^-M^+ pairs than for the $\text{PhCH}_2^-\text{M}^+$ pairs.


The correlation between the nature of the cation and λ_{max} of the bands for Fl^-M^+ , which have been identified as contact ion pairs, has been discussed¹⁸ in terms of the radius of the counterion. It has been pointed out that greater shifts from λ_{max} for the free ion are induced by cations of smaller size. However, in the case of solvent separated ion pairs formed with alkali metal cations, the pairing induces somewhat less of a shift than has been found with NBu_4^+ , a bulky ion with radius 4.5 Å, comparable to the dimension of the solvated cation in a solvent separated pair. If such a correlation, which involves relatively small shifts, may be ap-

plied to the ion pairs of benzyl carbanion, it reveals that Na^+ occupies the same relative position in the series as would a solvent separated ion pair, from which we infer that $\text{PhCH}_2^-\text{Na}^+$ in THF is a solvent separated pair. It is noteworthy that previous research²⁰ on polystyryl sodium, a somewhat similar carbanion pair, has led to the conclusion that PS^-Na^+ in THF is a contact ion pair. Apparently factors such as the degree of alkyl substitution at the benzylic position may determine the nature of the ion pair formed.

Absolute rate constants were determined for the proton transfer reaction of methanol and *tert*-butyl alcohol to $\text{PhCH}_2^-\text{Li}^+$



and to $\text{PhCH}_2^-\text{NBu}_4^+$



The decay of the absorption of the carbanion pair was monitored, after an electron pulse, at the maximum of either carbanion species in both the absence and the presence of the proton donor. The solutions of dibenzylmercury irradiated had a sufficiently high salt concentration, $[\text{LiClO}_4] = 0.04 F$ or $[\text{NBu}_4\text{ClO}_4] = 0.02 F$, so that formation of the ion pair was complete before the proton transfer reaction was observed. The alcohol concentrations were sufficiently high to make the decay of the carbanion cleanly first order. The pseudo-first-order rate constants obtained are linear in the alcohol concentration, as may be seen in Figures 3 and 4. The absolute rate constants, k_6 and k_7 (with an uncertainty of less than $\pm 20\%$) taken from the slopes of these plots, are shown in Table II along with previously determined values for PhCH_2^- and $\text{PhCH}_2^-\text{Na}^+$ for comparison.

It is evident from these data that factors which determine the rate constant for each alcohol persist for all four forms of the benzyl carbanion which we have investigated. Thus, $k_{\text{CH}_3\text{OH}} > k_{t\text{-C}_4\text{H}_9\text{OH}}$ is the same order as the acidity of these alcohols as measured in DMSO,²¹ indicating that the acid-base nature of the reaction is rate determining. Further, the ion paired species, in all three cases, is more reactive than the free carbanion. The range of the kinetic effect produced by the ion pairing is considerably greater than that resulting from the dependence upon the proton lability for these two alcohols. It is interesting that the rate constants for protonation of a carbanion derived from even so weakly acidic a hydrocarbon as toluene are lower than the diffusion controlled limit.

It is interesting that sodium, while it has the largest rate enhancing effect of the three counterions, has the smallest effect upon the absorption spectrum. This slight effect upon λ_{max} was attributed to the formation of a loose, rather than a contact, ion pair. We advance the speculation that the large rate enhancement may reflect the possibility that alcohol molecules may enter the primary solvation shell of the cation in a selective solvation, and are thus available for reaction; this close approach to the center of positive charge does not exist in the case of NBu_4^+ . Such specific solvation might also be expected for Li^+ which, to the contrary, shows a relatively small rate enhancement. A different effect, the degree of polarity of $\text{PhCH}_2^-\text{M}^+$, may be the determining factor for $\text{PhCH}_2^-\text{Li}^+$. NMR studies²² show that some degree of covalency exists for benzyl lithium, while we have reasoned that $\text{PhCH}_2^-\text{Na}^+$ is solvent separated, and therefore completely ionic in character. It seems reasonable to suggest that the ability of Li^+ to catalyze the proton transfer is reduced by the partial covalent nature of




Figure 3. Plot of pseudo-first-order rate constants for decay of benzyl carbanion pairs vs. *tert*-butyl alcohol concentration: (O) $\text{PhCH}_2^-\text{Li}^+$; (●) $\text{PhCH}_2^-\text{NBu}_4^+$. Second-order rate constants taken from the slopes are $k_6 = 9.7 \times 10^7$, and $k_7 = 4.6 \times 10^8 M^{-1} \text{sec}^{-1}$, for *tert*-butyl alcohol at 24° in THF.




Figure 4. Plot of pseudo-first-order rate constants for decay of benzyl carbanion pairs vs. methanol concentration: (O) $\text{PhCH}_2^-\text{Li}^+$; (●) $\text{PhCH}_2^-\text{NBu}_4^+$. Second-order rate constants taken from the slopes are $k_6 = 3.4 \times 10^8$ and $k_7 = 6.0 \times 10^8 M^{-1} \text{sec}^{-1}$, for methanol at 24° in THF.

$\text{PhCH}_2^-\text{Li}^+$ since this reduces the effective positive charge upon Li^+ .

Acknowledgment. We are indebted to Ed Ray for maintaining the linac and the electronic detection equipment.

References and Notes

- (1) This work was supported by the U.S. Atomic Energy Commission under Contract No. AT(11-1)-1763.
- (2) M. Szwarc, "Carbanions, Living Polymers, and Electron Transfer Processes", Interscience, New York, N.Y., 1968.
- (3) S. Bank and B. Bockrath, *J. Am. Chem. Soc.*, **93**, 430 (1971); **94**, 6076 (1972).
- (4) B. Bockrath and L. M. Dorfman, *J. Phys. Chem.*, **77**, 2618 (1973).
- (5) A. Rainis, R. Tung, and M. Szwarc, *J. Am. Chem. Soc.*, **95**, 659 (1973).
- (6) E. R. Minnich, L. D. Long, J. M. Cerasa, and J. L. Dye, *J. Am. Chem. Soc.*, **95**, 1061 (1973).
- (7) B. Bockrath and L. M. Dorfman, *J. Am. Chem. Soc.*, **96**, 5708 (1974).
- (8) L. L. Chan and J. Smid, *J. Phys. Chem.*, **76**, 695 (1972).
- (9) T. E. Hogen-Esch and J. Smid, *J. Am. Chem. Soc.*, **89**, 2764 (1967).
- (10) M. S. Matheson and L. M. Dorfman, *J. Chem. Phys.*, **32**, 1870 (1960).

- (11) L. M. Dorfman, "Techniques of Chemistry", Vol. VI, Part II, G. G. Hammes, Ed., Wiley, New York, N.Y., 1974, pp 463-519.
- (12) W. D. Felix, B. L. Gall, and L. M. Dorfman, *J. Phys. Chem.*, **71**, 384 (1967).
- (13) L. M. Dorfman, F. Y. Jou, and R. Wageman, *Ber. Bunsenges. Phys. Chem.*, **75**, 681 (1971).
- (14) F. Y. Jou and L. M. Dorfman, *J. Chem. Phys.*, **58**, 4715 (1973).
- (15) B. Bockrath and L. M. Dorfman, *J. Phys. Chem.*, **77**, 1002 (1973).
- (16) B. Bockrath and L. M. Dorfman, *J. Phys. Chem.*, in press.
- (17) R. Waack and M. A. Doran, *J. Am. Chem. Soc.*, **85**, 1651 (1963).
- (18) J. Smid, "Ions and Ion Pairs in Organic Chemistry", Vol. 1, M. Szwarc, Ed., Wiley-Interscience, New York, N.Y., 1972, pp 85-151.
- (19) R. Asami, M. Levy, and M. Szwarc, *J. Chem. Soc.*, 361 (1962).
- (20) Reference 2, pp 424-434.
- (21) C. D. Ritchie, "Solute-Solvent Interactions", J. F. Coetzee and C. D. Ritchie, Ed., Marcel-Dekker, New York, N.Y., 1969, p 230.
- (22) R. Waack, M. A. Doran, E. B. Baker, and G. A. Olah, *J. Am. Chem. Soc.*, **88**, 1272 (1966).


Influence of Solvent on the Ultrafast Kinetics of Electron Photoejection

W. S. Struve,^{1a} T. L. Netzel,^{1a} P. M. Rentzepis,^{1a} G. Levin,^{1b} and M. Szwarc^{*1b}


Contribution from Bell Laboratories, Murray Hill, New Jersey 07974, and the Department of Chemistry, SUNY College of Environmental Science and Forestry, Syracuse, New York 13210. Received June 3, 1974

Abstract: Picosecond spectroscopy revealed that sodium salt of tetraphenylethylene dianion ($T^{2-}, 2Na^+$) is bleached by a 530-nm pulse whether dissolved in dioxane or tetrahydrofuran (THF). It was established that in either solvent $T^{2-}, 2Na^+$, and not $T^{\cdot-}, Na^+$, is the photolyzed species. The original absorbance returns within about 10 psec when dioxane solution is photolyzed, but the bleached state persists for at least a nanosecond when the reaction occurs in THF. The different structure of the aggregate, tighter in dioxane and looser in THF, is reflected in the different behavior of the excited state. The absorbance of $T^{\cdot-}$ radical anion ($\lambda_{max} = 660$ nm) is not observed in either system within the time scale of our experiments. Although electron ejection follows the photolysis of $T^{2-}, 2Na^+$ in THF, either only a small fraction of excited aggregates undergoes autoionization (making the detection of $T^{\cdot-}$ hardly possible) or the resulting $T^{\cdot-}$ retains the geometry of T^{2-} for at least 60 psec (the long time of our observations) and the distorted radical anion apparently does not absorb in the expected region. To differentiate between these alternatives, sodium salt of tetracene dianion was flash photolyzed in dioxane and in THF. Bleaching and the appearance of tetracene radical anion was observed in both solvents. It is probable, therefore, that the substantially different shapes of T^{2-} and of $T^{\cdot-}$ prevent the observation of $T^{\cdot-}$ formation in dioxane.


Flash photolysis of sodium salt of tetraphenylethylene dianions, $T^{2-}, 2Na^+$, in tetrahydrofuran (THF) leads to electron photoejection²



The reaction is manifested by bleaching of the 480-nm absorption band of the dianion and by the simultaneous appearance of a 660-nm transient arising from the formation of the tetraphenylethylene radical anion, $T^{\cdot-}, Na^+$. The latter is produced by two processes: (1) by the primary photolytic act, and (2) through capture of the ejected electron by tetraphenylethylene, T



The latter reaction was rapid since the concentration of tetraphenylethylene in the photolyzed system was relatively large. In the following dark period, the disproportionation regenerated the dianions




and the systems returned every time precisely to its initial state.

The above experiments were performed with visible light ($\lambda > 420$ nm), the radiation of shorter wavelength being quantitatively absorbed by suitable filters. In a typical flash, about 20% of the 480-nm absorption had been bleached and the degree of bleaching had not been affected by the addition of even a large excess of sodium tetraphenylboride, the presence of which greatly reduced the concentration of T^{2-}, Na^+ . One concludes therefore that the photolysis involves the $T^{2-}, 2Na^+$ aggregate and *not* the half-


dissociated T^{2-}, Na^+ .

Repetition of these experiments with dioxane solution of $T^{2-}, 2Na^+$ led to surprising results.² Although the experimental conditions were basically the same as those maintained in the previous experiments, no bleaching whatsoever of the 480-nm band had been observed, even when substantially more powerful flashes were employed.

It is known that the ionic aggregates are tighter in dioxane than in tetrahydrofuran and therefore the different behavior of this salt in THF and in dioxane has to be attributed to the difference of their structures. Apparently the absorbed light excites the $T^{2-}, 2Na^+$ aggregate to its autoionizing state, and the excited species formed from the looser aggregate present in THF has a finite, although probably small, probability to undergo autoionization leading to electron ejection into the bulk of the liquid. Such an electron may escape the attractive field of the sodium ions and has then a chance to be trapped by T. On the other hand, the excitation of the tight aggregate present in dioxane leads not to its autoionization but probably to intramolecular electron transfer to a sodium cation



The reverse reaction, a kind of cage process, occurs rapidly, e.g.



and therefore the relatively slow flash-photolysis technique is incapable of revealing the occurrence of this process. To test these ideas, the photolysis of $T^{2-}, 2Na^+$ in THF and in dioxane was reinvestigated by the method of picosecond absorption spectroscopy.³