

- (28) Leffler, J. E.; Grunwald, E. "Rates and Equilibria of Organic Reactions"; Wiley: New York, 1963; p 192. See also ref 27a for a related discussion.
- (29) *f* would have to assume the unrealistic value of 0.8 to predict a relative rate of 1.45 on the basis of a one-terminus mechanism.
- (30) The ρ value for [2 + 2] cycloaddition is comparable with that for reaction by the ene³¹ and by the [2 + 4]³² modes.
- (31) Foote, C. S.; Denny, R. W. *J. Am. Chem. Soc.* **1971**, *93*, 5162.
- (32) Young, R. H.; Martin, R. L.; Chinh, N.; Mallon, C.; Kayser, R. H. *Can. J. Chem.* **1972**, *50*, 932.

K. A. Zaklika, Bashir Kaskar, A. Paul Schaap*

Department of Chemistry, Wayne State University
Detroit, Michigan 48202

Received June 20, 1979




Figure 1. ESR spectrum of DCA^{•-} obtained by photolysis of a solution of DCA (5×10^{-4} M) and 1,1-diphenylethylene (10^{-2} M) in dry acetonitrile under Ar.⁶

Mechanisms of Photooxygenation. 2. Formation of 1,2-Dioxetanes via 9,10-Dicyanoanthracene-Sensitized Electron-Transfer Processes





Sir:

Electron-transfer quenching of singlet excited sensitizers by olefins has recently received considerable attention.¹ Radical-ion intermediates have been characterized by transient absorption spectroscopy² and CIDNP methods.³ 9,10-Dicyanoanthracene (DCA) is a typical sensitizer for photogeneration of radical ions. Electron-transfer quenching of the fluorescence of DCA by a variety of substrates has been postulated from correlations of quenching rate constants and free energies of electron transfer,⁴ from solvent dependent exciplex emissions,^{4a} and on the basis of distinctive photochemistry.⁵

We now provide direct evidence for the photochemical formation of the DCA radical anion (DCA^{•-}) with the observation of the ESR spectrum of this species upon irradiation of solutions of DCA and various quenchers in deoxygenated acetonitrile.⁶ Figure 1 shows an ESR spectrum obtained with 1,1-diphenylethylene (**1**). The spectrum is identical with that reported by Brunner and Dörr for DCA^{•-} prepared by metal/ammonia reduction.⁷ Under conditions of high modulation, the spectrum becomes similar to that obtained by Happ and Janzen following addition of cyanide ion to an air-saturated solution of 9-cyanoanthracene.⁸ The ESR spectrum of DCA^{•-} was also obtained upon electrochemical reduction of DCA^{9,10} or on irradiation of DCA in the presence of quenchers **2-6a-d** (Chart I), all of which can reduce singlet excited DCA exothermically.¹¹ Additionally, DCA^{•-} could be photogenerated in dichloromethane and acetone in the presence of **6a**.







The radical cation of the quencher should also be a primary product of electron transfer to singlet excited DCA (Scheme I, eq 3). We have characterized by ESR spectroscopy a variety

Chart I



- a. R₁=R₂=*p*-MeOC₆H₄
b. R₁=R₂=2-naphthyl
c. R₁=H; R₂=C₆H₅
d. R₁=C₆H₅; R₂=2-(6-dimethylaminonaphthyl)
e. R₁=R₂=*m*-MeOC₆H₄
f. R₁=R₂=*p*-MeC₆H₄

Scheme I



of radical cations of the general structure **6** produced by electrochemical oxidation¹² or by chemical oxidation with tris(*p*-bromophenyl)ammoniumyl tetrafluoroborate (**7**).¹³ Although ESR signals due to **6**^{•+} are readily detected, we have so far been unable to observe them upon irradiation of mixtures of DCA and **6**.¹⁴ Flash experiments with 1,1-diphenylethylene (**1**) and the related sensitizer methyl *p*-cyanobenzoate² are, however, reported to result in a transient absorption ascribable to **1**^{•+}.

While the photochemical generation of DCA^{•-} is of general significance to electron-transfer quenching, it is particularly relevant to the mechanism of DCA-sensitized photooxygenation recently proposed by Foote.^{5a} A related reaction had been described earlier by Farid.^{5b,15} Foote observed carbonyl products suggestive of 1,2-dioxetane intermediates when olefins were photooxygenated under these conditions. We find that oxygenation of olefins of structure **6** using DCA or 9,10-dicyano-2-methylantracene (MeDCA) as sensitizers leads to substantial yields of 1,2-dioxetanes.¹⁶ Figure 2 shows the time evolution of the photooxygenation of **6a** using MeDCA in acetone. It is evident that, unless the MeDCA is efficiently quenched by the olefin, it sensitizes the cleavage of the 1,2-dioxetane¹⁷ which can therefore escape observation. In another experiment **6a** yielded 45% dioxetane (characterized by ¹H NMR spectroscopy and chemiluminescence¹⁸), whereas the yields were 53% with **6e** and 61% with **6f**. Pho-




Figure 2. Time evolution of the MeDCA-sensitized photooxygenation of 2,3-di(*p*-methoxyphenyl)-1,4-dioxene (**6a**) at 0 °C in acetone-*d*₆. ¹H NMR spectroscopy was used to determine the concentration of **6a** (●), the corresponding 1,2-dioxetane (▲), and the resultant cleavage product (■).

photooxygenation of adamantylideneadamantane (**5**) in dichloromethane led to isolation of the stable dioxetane¹⁹ in 88% yield. Similar treatment of olefins **1** and **2** gave the corresponding benzophenones in 30²⁰ and 37% isolated yield.

Competition kinetics between olefins of structure **6** suggest singlet oxygen (¹O₂), derived from triplet DCA, is not involved.²¹ Relative rates are distinct from singlet oxygen rates and are, moreover, a function of photooxygenation time. For instance, the relative rate for **6a/6f** is 1.64 for reaction with ¹O₂ but ranges from 6.7 (16% **6a** reacted) to 2.4 (94% **6a** reacted) in the DCA system.²⁴ Reaction of **6** with ¹O₂ produced by eq 1 and 2 (intersystem crossing)²⁵ or by eq 3, 4, and 2 (annihilation)²⁶ appears therefore to be excluded. Since DCA^{-•} is efficiently quenched by oxygen under our conditions (eq 5), as evidenced by ESR spectroscopy and cyclic voltammetry, we focused attention on eq 6, the final step of the Foote mechanism (eq 3, 5, and 6).

Treatment of **6a** with 1 equiv of tris(*p*-bromophenyl)-ammonium tetrafluoroborate (**7**) in dichloromethane at -78 °C under argon generated a deeply violet solution of **6a**⁺. Following addition at -78 °C of 1 equiv of potassium superoxide dissolved in dichloromethane using 18-crown-6, the color was discharged within 5 min upon warming to -20 °C.²⁷ Chemiluminescence resulted when the reaction mixture was heated after dilution with a solution of 9,10-dibromoanthracene in *o*-xylene. The luminescence decayed at a rate identical with that from the 1,2-dioxetane formed from **6a** and singlet oxygen,¹⁸ at several temperatures spanning a 20 °C range. Addition of authentic dioxetane to the spent chemiluminescence solutions at known concentration and integration of the emitted light showed that this dioxetane was formed in 0.4% yield from **6a**⁺ and O₂^{-•}. Analysis of the reaction mixture revealed tris(*p*-bromophenyl)amine, recovered **6a** (42%), the diester cleavage product of the dioxetane, ethylene glycol di(*p*-methoxybenzoate) (**8**, 5%), and three other products derived from **6a** which are as yet not fully characterized. Recovery of substantial amounts of **6a** suggests that electron transfer from O₂^{-•} to **6a**⁺ is a competitive reaction. In a control experiment, where oxygen was bubbled into the solution at -20 °C in place of addition of superoxide, 60 min were required for complete decolorization of the solution.²⁸ The yield of dioxetane was diminished 4.5-fold; 35% **6a** was recovered while 2% **8** was found along with the other products. In a

similar experiment with **6a** and **7** at -20 °C with acetonitrile as solvent, the solution decolorized as soon as superoxide was added. Dioxetane was again found in 0.4% yield, together with **6a** (14%), **8** (19%), and the other products. Admission of oxygen without addition of superoxide²⁹ using either 0.5 or 1 equiv of **7** reduced the yield of dioxetane 18-fold. Dioxetane yields have not been optimized.

The longer reaction times and reduced yields of dioxetane in the absence of superoxide make it unlikely that the bulk of the dioxetane can arise by a chain mechanism (eq 7 and 8) similar to that proposed³⁰ for an oxygenation first described by Barton et al.^{13c,31} In the case of **6a**, formation of ¹O₂ and olefin by electron transfer within the solvent cage is a possibility³² (eq 9). Cycloaddition would then be competitive with diffusion into the bulk solution where quenching by superoxide would predominate.³³ Based on the known rate constants and on production of ¹O₂ with unit efficiency, the maximum yield of dioxetane is predicted to be 2.1%.³⁵ However, ¹O₂ is formed with an efficiency much less than 1 in analogous reactions³² and **4** will undergo DCA-sensitized photooxygenation, even though it is unreactive toward ¹O₂.^{5a} Equation 9 is, therefore, a poor contender for the major pathway of dioxetane formation in this instance. However, some contribution from ¹O₂ cannot be excluded at present.

Another mechanism, additional to that of Foote, which could account for formation of dioxetane from chemically or photochemically generated **6a**⁺ is the sequence described by eq 10, 11, and 8. Disproportionation of ions **6**⁺ appears to be thermodynamically, but not necessarily kinetically, favored.³⁶ This possibility is under investigation. Work is in progress to further define the scope and mechanistic basis of the experiments reported here. Nevertheless, the results presently available show clearly that oxygenation mediated by electron transfer can be a viable mechanism for a variety of olefinic substrates.

Acknowledgments. Support of this research by the U.S. Army Research Office and the National Science Foundation is gratefully acknowledged. We also thank Professor John F. Endicott for helpful discussions.

References and Notes

- (a) Smothers, W. K.; Schanze, K. S.; Saltiel, J. *J. Am. Chem. Soc.* **1979**, *101*, 1895. (b) O'Connor, D. V.; Ware, W. R. *Ibid.* **1979**, *101*, 121. (c) Maroulis, A. J.; Shigemitsu, Y.; Arnold, D. R. *Ibid.* **1978**, *100*, 535. (d) Mariano, P. S.; Stavinoha, J. L.; Pépe, G.; Meyer, E. F. Jr., *Ibid.* **1978**, *100*, 7114. (e) Miyamoto, T.; Tsujimoto, Y.; Tsuchinaga, T.; Nishimura, Y.; Odaira, Y. *Tetrahedron Lett.* **1978**, 2155. (f) McCullough, J. J.; Miller, R. C.; Fung, D.; Wu, W.-S. *J. Am. Chem. Soc.* **1975**, *97*, 5942. (g) Arnold, D. R.; Humphreys, R. W. R. *Ibid.* **1979**, *101*, 2743.
- Neunteufel, R. A.; Arnold, D. R. *J. Am. Chem. Soc.* **1973**, *95*, 4080.
- (a) Arnold, D. R.; Wong, P. C. *J. Am. Chem. Soc.* **1979**, *101*, 1894. (b) Roth, H. D.; Schilling, M. L. M. *Ibid.* **1979**, *101*, 1898, and references therein.
- (a) Ware, R. W.; Holmes, J. D.; Arnold, D. R. *J. Am. Chem. Soc.* **1974**, *96*, 7861. (b) Eriksen, J.; Foote, C. S. *J. Phys. Chem.* **1978**, *82*, 2659.
- (a) Eriksen, J.; Foote, C. S.; Parker, T. L. *J. Am. Chem. Soc.* **1977**, *99*, 6455. (b) Brown-Wensley, K. A.; Mattes, S. L.; Farid, S. *Ibid.* **1978**, *100*, 4162. (c) Ando, W.; Nagashima, T.; Saito, K.; Kohmoto, S. *J. Chem. Soc., Chem. Commun.* **1979**, 154.
- ESR spectra were acquired on a Varian E109E spectrometer equipped with a Nicolet 535 signal averager. Photochemical radical formation was studied by irradiating at ambient temperature, under argon, solutions of the quencher (10⁻²-10⁻³ M) and DCA (10⁻³-10⁻⁴ M) in MeCN which had been distilled from CaH₂ and P₂O₅. The illumination of the solution was carried out in the ESR cavity with a Varian EIMAC 150-W illuminator through a filter solution.¹⁶
- Brunner, E.; Dörr, F. *Ber. Bunsenges. Phys. Chem.* **1964**, *68*, 468.
- Happ, J. W.; Janzen, E. G. *J. Org. Chem.* **1970**, *35*, 96.
- Controlled potential electrolysis of DCA (10⁻³ M) in MeCN containing 0.1 M tetraethylammonium perchlorate was carried out at -1.2 V vs. SCE under Ar. A flow system was used with external generation of DCA^{-•} at a Pt electrode.
- The ESR spectrum of electrochemically generated DCA^{-•} has recently been described: Yildiz, A.; Baumgärtel, H. *Ber. Bunsenges. Phys. Chem.* **1977**, *81*, 1177.
- See ref 4 and 5a,b for discussions of the energetics of electron transfer from quenchers to singlet excited DCA.
- (a) See note 9. Electrolyses of olefins **6** were carried out at potentials of (E_{1/2} + 0.2) V. (b) For the half-wave potentials of **6**, see Zaklika, K. A.:

- Kaskar, B.; Schaap, A. P. *J. Am. Chem. Soc.*, preceding paper in this issue.
- (13) (a) Nelson, R. F.; Adams, R. N. *J. Am. Chem. Soc.* **1968**, *90*, 3925. (b) Bell, F. A.; Ledwith, A.; Sherrington, D. C. *J. Chem. Soc. C* **1969**, 2719. (c) Barton, D. H. R.; Haynes, R. K.; Leclerc, G.; Magnus, P. D.; Menzies, I. D. *J. Chem. Soc., Perkin Trans. 1* **1975**, 2055.
- (14) The lack of an ESR signal for $6^{+\bullet}$ could be due to electron transfer from $6^{+\bullet}$ to singlet excited DCA to yield $DCA^{-\bullet}$ and the dication of **6**. Cyclic voltammetry studies of olefins **6** indicate that this reaction would be exothermic.
- (15) Farid, S.; Brown, K. A.; Doty, J. C.; Hartman, S. E.; Williams, J. L. R. *Abstr. Pap. Int. Conf. Photochem., 8th, 1975* **1975**, Abstract Y1.
- (16) Typically photooxygenations were performed at 0 °C with 10^{-2} M substrate and 5×10^{-4} M sensitizer under an oxygen atmosphere. Concentrations were monitored by $^1\text{H NMR}$ or GC. A 450-W medium-pressure Hg lamp was used for irradiation with a 1-cm path-length filter solution made from 27 g of $\text{CuSO}_4 \cdot 5\text{H}_2\text{O}$, 30 g of NaNO_2 , and 50 mL of concentrated NH_4OH diluted with water to 1000 mL.
- (17) Wilson, T.; Schaap, A. P. *J. Am. Chem. Soc.* **1971**, *93*, 4126.
- (18) Zaklika, K. A.; Thayer, A. L.; Schaap, A. P. *J. Am. Chem. Soc.* **1978**, *100*, 4916.
- (19) (a) Wieringa, J. H.; Strating, J.; Wynberg, H.; Adam, W. *Tetrahedron Lett.* **1972**, 169. (b) Schuster, G. B.; Turro, N. J.; Steinmetzer, H.-C.; Schaap, A. P.; Falser, G.; Adam, W.; Liu, J. C. *J. Am. Chem. Soc.* **1975**, *97*, 7110.
- (20) Isolated as the 2,4-dinitrophenylhydrazones.
- (21) Attempts were made to quench any reaction via $^1\text{O}_2$. The quenchers, β -carotene²² and bis(dithiobenzil)nickel,²³ were unstable to the photooxygenation conditions. 1,4-Diazabicyclo[2.2.2]octane (Dabco)²² did not quench within experimental error at a concentration where 75% of an authentic $^1\text{O}_2$ reaction was quenched. Dabco ($E_{1/2} = 0.64$ V vs. SCE) could, however, reduce singlet excited DCA.
- (22) For a discussion of singlet oxygen quenchers see: (a) Belluš, D. "Singlet Oxygen", Rånby, B.; Rabek, J. F., Eds.; Wiley: New York, 1978; Chapter 9. (b) Denny, R. W.; Nickon, A. *Org. React.* **1973**, *20*, 133.
- (23) Zweig, A.; Henderson, W. A. Jr., *J. Polym. Sci.* **1975**, *13*, 717.
- (24) There is some evidence from ESR experiments that this may be related to electron transfer from **6a** to $6^{+\bullet}$.
- (25) Unlikely because DCA is efficiently fluorescent: Melhuish, W. J. *J. Phys. Chem.* **1961**, *65*, 2291.
- (26) (a) Orbach, N.; Patoshnik, R.; Ottolenghi, M. *J. Phys. Chem.* **1972**, *76*, 1133. (b) Schulton, K.; Staerk, H.; Weller, A.; Werner, H.-J.; Nickel, B. *Z. Phys. Chem. (Frankfurt am Main)* **1976**, *101*, 371. (c) Mataga, N.; Migita, M.; Wishimura, T. *J. Mol. Struct.* **1978**, *47*, 199. (d) Hayashi, H.; Nagakura, S. *Chem. Phys. Lett.* **1978**, *53*, 201. (e) Tachikawa, H.; Faulkner, L. R. *J. Am. Chem. Soc.* **1976**, *98*, 5569. (f) Park, S. M.; Bard, A. J. *Ibid.* **1975**, *97*, 2978.
- (27) No apparent color change occurred over a period of 1.7 h at -78 °C.
- (28) No apparent color change occurred over a period of 1.3 h at -78 °C.
- (29) In this case the solution decolorized within 3 min.
- (30) Tang, R.; Yue, H. J.; Wolf, J. F.; Mares, F. *J. Am. Chem. Soc.* **1978**, *100*, 5248.
- (31) For related studies see also: (a) Saeva, F. D.; Olin, G. R. *J. Chem. Soc., Chem. Commun.* **1976**, 943. (b) Anderson, C. P.; Salmon, D. J.; Meyer, T. J.; Young, R. C. *J. Am. Chem. Soc.* **1977**, *99*, 1980.
- (32) Mayeda, E. A.; Bard, A. J. *J. Am. Chem. Soc.* **1973**, *95*, 6223.
- (33) $\text{O}_2^{-\bullet}$ quenches³⁴ $^1\text{O}_2$ with $k_q = 7 \pm 6 \times 10^9 \text{ M}^{-1} \text{ s}^{-1}$, whereas k_q for **6a** is $6.6 \times 10^7 \text{ M}^{-1} \text{ s}^{-1}$. The triarylamine and 18-crown-6 are expected to quench less efficiently still than **6a**.²²
- (34) Guiraud, H. J.; Foote, C. S. *J. Am. Chem. Soc.* **1976**, *98*, 984.
- (35) Calculated for **6a** using a reaction rate with $^1\text{O}_2$ of $12.6 \times 10^7 \text{ M}^{-1} \text{ s}^{-1}$ (see ref 12b) and a diffusion rate constant of $12.6 \times 10^{10} \text{ M}^{-1} \text{ s}^{-1}$ for acetonitrile at -20 °C.
- (36) Compounds **6** show a second oxidation wave with a peak potential 0.2–0.5 V more positive than the half-wave potential for one-electron oxidation (see ref 12b).

A. Paul Schaap,* K. A. Zaklika
Bashir Kaskar, L. W.-M. Fung

Department of Chemistry, Wayne State University
Detroit, Michigan 48202


June 20, 1979

Electron-Transfer Photooxidation. 3. Detection of Radical-Ion Intermediates in the Cyanaromatic-Sensitized Photooxidation of *trans*- and *cis*-Stilbene¹

Sir:

Radical ions have been proposed as intermediates in a number of photochemical systems,² including photooxidation reactions involving cyanaromatic sensitizers and electron-rich substrates.^{3,4} The proposed mechanism for these photooxidation reactions (Scheme I) involves electron transfer from

Scheme I



the substrate to the excited singlet sensitizer; the radical anion of the sensitizer then reacts with oxygen to produce superoxide ion, which reacts with the substrate radical cation to produce the oxidation products.⁵

We now report the direct observation of the radical-ion intermediates in the photooxidation of *trans*-stilbene (TS) and *cis*-stilbene (CS) using 9,10-dicyanoanthracene (DCA) and 9-cyanoanthracene (CA) as sensitizers. The absorption spectra were obtained by laser flash photolysis⁸ using a Nd-YAG laser as the excitation source at 355 nm and monitoring with a xenon flash lamp. Figure 1 shows the absorption spectrum (1 μs after laser pulse) produced from 2.0×10^{-4} M CA and 0.05 M TS in N_2 -saturated dry CH_3CN . The absorption (λ_{max} 470 nm) is very similar to that of $\text{TS}^{\bullet+}$, previously published.^{9–11} The spectrum also shows an absorption at λ_{max} 587 nm that corresponds well to the absorption spectra of $\text{CA}^{-\bullet}$.¹² The transient decay rate at 587 nm increases on going from N_2 - to air- to oxygen-saturated solution. In N_2 , the decay appears to be second order, while, in air and O_2 , there is an initial fast component and a slower decay at longer times. In contrast, the decay rate at 470 nm shows mixed first- and second-order kinetics. The decay rate under air is equal to or only slightly faster than that under N_2 and is larger under O_2 (Figure 2).




Figure 1. Absorption spectrum of 0.05 M TS + 2.0×10^{-4} M CA in CH_3CN , 1.0 μs after flash.