

- (30) (a) An equilibrium constant for the addition of hydrogen cyanide to acetophenone in 96% ethanol at 70 °C is estimated^{30b} to be $\sim 10^{-1} \text{ M}^{-1}$. The equilibrium constant for the addition of aliphatic amines to aldehydes in water at 20–25 °C is $\sim 10^{-3}$ smaller than that observed for hydrogen cyanide and is independent of amine basicity.²⁸ Since the ratio of equilibrium constants for the addition of different nucleophiles to aldehydes is independent of electron-withdrawing groups²⁸ on the aldehyde, the ratio probably applies to acetophenone as well. To the extent that solvent and temperature effects on the ratio are small, the equilibrium constant for the addition of amines to acetophenone to give the neutral carbinolamine will be near 10^{-4} M^{-1} . Whereas the change in solvent might raise the estimated equilibrium constant, the increased steric requirements^{30c} and resonance effect²⁸ would be expected to lower it. (b) Lapworth, A.; Manske, R. *J. Chem. Soc.* **1930**, 1976. Evans, D. P.; Young, J. R. *Ibid.* **1954**, 1310. (c) Prelog, V.; Kobelt, M. *Helv. Chim. Acta* **1949**, 32, 1187.
- (31) Miller, D. L.; Westheimer, F. H. *J. Am. Chem. Soc.* **1966**, 1507, 1511, 1514. Miller, D. L.; Ukena, T. *Ibid.* **1969**, 91, 3050.
- (32) DiSabato, G.; Jencks, W. P. *J. Am. Chem. Soc.* **1961**, 83, 4400. Benkovic, S. J.; Schray, K. J. *Biochemistry* **1968**, 7, 4090. Haake, P.; Allen, G. *Proc. Natl. Acad. Sci. U.S.A.* **1971**, 68, 2691; *J. Am. Chem. Soc.* **1973**, 95, 8080; *Ibid.* **1976**, 98, 4990.
- (33) Kirby, A. J.; Warren, S. G. "The Organic Chemistry of Phosphorus"; Elsevier: Amsterdam, 1967; p 281 ff. Bruce, T. C.; Benkovic, S. "Bioorganic Mechanisms"; W. A. Benjamin: New York, 1966; Vol. 2, pp 22–25, 157–159. Benkovic, S. J.; Schray, K. J., *Enzymes* **1973**, 8, 201. Jencks, W. P. "Catalysis in Chemistry and Enzymology"; McGraw-Hill: New York, 1969; pp 81–83, 112–115, 151, 160–161, 608. Butcher, W. W.; Westheimer, F. H. *J. Am. Chem. Soc.* **1955**, 77, 2420. Barnard, P. W. C.; Bunton, C. A.; Llewellyn, D. R.; Oldham, K. G.; Silver, B. L.; Vernon, C. A. *Chem. Ind. (London)* **1955**, 760. Todd, A. R. *Proc. Natl. Acad. Sci. U.S.A.* **1959**, 45, 1389. Kirby, A. J.; Varvoglis, A. G. *J. Am. Chem. Soc.* **1967**, 89, 415. Haake, P.; Ossip, P. S. *Ibid.* **1971**, 93, 6924. Gorenstein, D. G. *Ibid.* **1972**, 94, 2523. Kluger, R. *J. Org. Chem.* **1973**, 38, 2721.
- (34) Jencks, W. P.; Gilchrist, M. *J. Am. Chem. Soc.* **1964**, 86, 1410. Traylor, P. S.; Westheimer, F. H. *Ibid.* **1965**, 87, 553. Kirby, A. J.; Jencks, W. P. *Ibid.* **1965**, 87, 3209. Benkovic, S. J.; Benkovic, P. A. *Ibid.* **1967**, 89, 4714. Kirby, A. J.; Varvoglis, G. A. *J. Chem. Soc. B.* **1968**, 135. Guthrie, J. P. *J. Am. Chem. Soc.* **1977**, 99, 3991.
- (35) Mildvan, A. *Adv. Enzymol.* **1979**, 49, 103.
- (36) Rose, I. A. *J. Biol. Chem.* **1960**, 235, 1170. Robinson, J. L.; Rose, I. A. *Ibid.* **1972**, 247, 1096. Kuo, D. J.; Rose, I. A. *J. Am. Chem. Soc.* **1978**, 100, 6288. Kuo, D. J.; O'Connell, E. L.; Rose, I. A. *Ibid.* **1979**, 101, 5025.
- (37) Lowe, G.; Sproat, B. S. *J. Chem. Soc., Chem. Comm.* **1978**, 783; *J. Chem. Soc., Perkin Trans. 1* **1978**, 1622.
- (38) Rose, I. A. *Adv. Enzymol.* **1979**, 50, 361. See addendum.
- (39) Abbott, S. J.; Jones, S. R.; Weinman, S. A.; Knowles, J. R. *J. Am. Chem. Soc.* **1978**, 100, 2558. Blattler, W. A.; Knowles, J. R. *Ibid.* **1979**, 101, 510; *Biochemistry* **1979**, 18, 3927.
- (40) Warren, W. A. *Biochim. Biophys. Acta* **1968**, 156, 340. Horiguchi, M.; Rosenberg, H. *Ibid.* **1975**, 404, 333.
- (41) Westheimer, F. H. *Acc. Chem. Res.* **1968**, 1, 70.
- (42) (a) Buchanan, J. *Adv. Enzymol.* **1973**, 39, 91. Greenberg, D. M.; Wynston, L. K.; Nagabhushanam, A. *Biochemistry* **1965**, 4, 1872. (b) Lieberman, I. *J. Biol. Chem.* **1956**, 223, 327. Koshland, D. E., Jr. *Biochemistry* **1971**, 10, 3365. Markham, G. D.; Reed, G. H. *J. Biol. Chem.* **1978**, 253, 6184.
- (43) Page, M. I.; Jencks, W. P. *Proc. Natl. Acad. Sci. U.S.A.* **1971**, 68, 1678.
- (44) van de Werf, P.; Orłowski, M.; Meister, A. *Proc. Natl. Acad. Sci. U.S.A.* **1971**, 68, 2982.
- (45) Roberts, J. W.; Roberts, C. W.; Craig, N. L. *Proc. Natl. Acad. Sci. U.S.A.* **1978**, 75, 4714. Bridges, B. A. *Nature (London)* **1979**, 514.
- (46) Herskko, A.; Ciechanover, A.; Rose, I. A. *Proc. Natl. Acad. Sci. U.S.A.* **1979**, 76, 3107.
- (47) De Martino, G. N.; Goldberg, A. L. *J. Biol. Chem.* **1979**, 254, 3712.
- (48) Lipmann, F. *Adv. Enzymol.* **1941**, 1, 99.

Dioxygen Transfer from 4a-Hydroperoxyflavin Anion. 2. Oxygen Transfer to the 10 Position of 9-Hydroxyphenanthrene Anions and to 3,5-Di-*tert*-butylcatechol Anion

Shigeaki Muto and Thomas C. Bruce*

Contribution from the Department of Chemistry, University of California at Santa Barbara, Santa Barbara, California 93106. Received December 10, 1979


Abstract: The reaction of the peroxy anion of *N*⁵-ethyl-4a-hydroperoxy-3-methylflavin (4a-FIET₂⁻) with the anions of 3,5-di-*tert*-butylcatechol (VIII), 10-ethoxy-9-phenanthrol (Ia), and 10-methyl-9-phenanthrol (Ib) has been investigated. All products may be accountable through the transfer of O₂ from the 4a-FIET₂⁻ reactant to the phenolate anions with the production of reduced flavin anion (FIET⁻) and a hydroperoxycyclohexadienone. From VIII⁻(*t*-BuOH) there was obtained 3,5-di-*tert*-butyl-*o*-quinone (IX) and Ib⁻ yielded (*t*-BuOH or CH₂CN) 10-hydroxy-10-methyl-9,10-dihydro-9-phenanthrone (IIIb), while Ia⁻ provided both 9,10-phenanthrenequinone (V) and monoethyl 1,1'-diphenate (IVa) (the ratio of V:IVa being solvent dependent). The mechanisms for the decomposition of intermediate peroxide anions to products are discussed. The conversion of Ia⁻ to IVa by oxygen transfer from 4a-FIET₂⁻ amounts to a catalysis by FIET⁻ of the reaction of ³O₂ with Ia and serves as a biomimetic reaction of flavoenzyme dioxygenase. The kinetics for the reaction of VIII⁻ with 4a-FIET₂⁻ require the formation of an intermediate. Since the rate constants for the reaction of both VIII⁻ and 2,6-di-*tert*-butyl-4-methylphenolate anion with 4a-FIET₂⁻ are identical under saturating conditions by these phenolate ions, it is concluded that the intermediate is formed in a unimolecular reaction from 4a-FIET₂⁻ ($k = 0.36 \text{ s}^{-1}$) as in eq 19. Dissociation of 4a-FIET₂⁻ to FIET⁻ + O₂ and reaction of phenolate ions with O₂ may be discounted since the second-order rate constants for the reaction of phenolate ions with O₂ are less than required for the kinetic competency of this process. Dissociation of 4a-FIET₂⁻ to yield neutral flavin radical (FIET[•]) + O₂^{-•} followed by reduction of FIET[•] by phenolate ion to provide FIET⁻ and phenoxy radical with the coupling of the latter with O₂^{-•} is also improbable. Thus, though the second-order rate constants for le⁻ reduction of FIET[•] by the various phenolate species are sufficiently large to allow the kinetic competency of this step, there exists no evidence that O₂^{-•} can couple with any radical species to provide a hydroperoxide. The oxygen-donating intermediate formed from 4a-FIET₂⁻ is suggested to be the 4a,10-dioxetane (XII) or an oxygen molecule more loosely associated with FIET⁻. The equilibrium constant for the formation of such an intermediate may be as small as 10⁻⁵ if the rate of reaction of phenolate ion with this species approaches a diffusion-controlled process.

Of much present concern are the mechanisms by which the mono- and dioxygenase enzymes combine with triplet molecular oxygen to provide species capable of transferring one or both of the oxygen atoms to a substrate molecule. The only non-metal-requiring oxygenases require flavin molecules as cofactors. This laboratory has been engaged in mechanistic


studies of the reaction of oxygen and oxygen species with flavins and the problem of oxygen activation by flavins.^{1–8} The transfer of molecular oxygen in toto by metallo-dioxygenase enzymes was first recognized by Hayaishi.⁹ The mechanisms of these reactions are at least partially understood and metal-centered biomimetic systems have been explored.¹⁰

Some understanding of flavin monooxygenases exists. Thus, flavin biomimetic systems for the activation of $^3\text{O}_2$ and transfer of a single oxygen atom to a substrate have been reported.^{1,2,4,8}


The identity or similarity of the spectrum of 4a-FIEtO₂H^{2.5} to that of the initial product obtained upon the reaction of molecular oxygen with reduced flavomonooxygenases¹¹ has established the importance of 4a-hydroperoxyflavins in the mechanisms of these enzymes. It may only be presumed that 4a-hydroperoxyflavins are also involved in flavoenzyme dioxygenase reactions.¹² In a previous publication from this laboratory (part 1 of these studies)⁷ there is described the reaction between 3-methyl-5-ethylflavin-4a-peroxy anion (4a-FIEtO₂⁻) and 2,6-di-*tert*-butyl-4-methylphenolate ion (eq 1). In this reaction (*t*-BuOH solvent) 4a-FIEtO₂⁻ becomes



reduced to the 1,5-dihydroflavin and the phenolate ion is converted into the corresponding dienone hydroperoxide. The reaction is very facile (stopped flow) and the products are obtained in ~95% yield. Since reduced *N*⁵-ethylflavin reacts with molecular oxygen to yield 4a-FIEtO₂⁻ the reaction of eq 1 represents one-half of the catalytic cycle of eq 2. We



report herein extensions of the 4a-FIEtO₂⁻ dioxygenation reaction to other substrates. It is our purpose both to gain an understanding of the mechanism of this reaction and to provide examples wherein dioxygenation by 4a-FIEtO₂⁻ will lead to a dioxygenase reaction (eq 3).



Experimental Section

General. All melting points were measured on a Thomas Model 40 micro hot stage apparatus and are uncorrected. Proton magnetic resonance (¹H NMR) spectra were recorded on a Varian T60 spectrometer using tetramethylsilane (Me₄Si) as internal reference. Chemical shifts were expressed as δ values (parts per million downfield from Me₄Si). All spectrophotometric determinations were performed on a Cary 118 spectrophotometer thermostated at 30.0 ± 0.2 °C. Rapid spectral changes were followed with a Durham stopped-flow spectrophotometer under an oxygen-free N₂ atmosphere.

Materials, 9,10-Phenanthrenequinone (V), 3,5-di-*tert*-butylcatechol (VIII), and 3,5-di-*tert*-butyl-*o*-quinone (IX) were obtained from Aldrich Chemical Co. *tert*-Butyl alcohol was distilled under nitrogen. Dioxane was refluxed over benzophenone and Na for 3 h and distilled under nitrogen. Tetrahydrofuran was first distilled from Na and then from LiAlH₄ under nitrogen. Acetonitrile was refluxed over P₂O₅ for 6 h and distilled under nitrogen. All solvents were kept under a dry

nitrogen atmosphere. The preparation of all solutions, weighing of solids, etc., were carried out in a dry nitrogen atmosphere.

***N*⁵-Ethyl-4a-hydroperoxy-3-methylflavin (4a-FIEtOOH).** 4a-FIEtOOH has been synthesized in this laboratory in 85 to >99% purity,^{1,2} λ_{max} 370 nm (ε₃₇₀(*t*-BuOH) 8000 M⁻¹ cm⁻¹).

10-Ethoxy-9-phenanthral (Ia). Ia was synthesized according to a literature procedure.¹³ The product was isolated by silica gel column chromatography (chloroform-*n*-hexane (1:1)) and its purity determined by thin layer chromatography. Recrystallization from *n*-hexane under N₂ and drying over P₂O₅ at 64 °C for 6 h in vacuo provided Ia free of ethanol: mp 74–76 °C (lit.¹³ 73–76 °C); ¹H NMR (CDCl₃) δ 1.65 (t, 3 H), 4.27 (q, 2 H), 6.30 (s, 1 H), 7.50–8.87 (m, 8 H).

10-Methyl-9-phenanthrol (Ib). According to the method of Eistert and El-Chahawi,¹⁴ Ib was obtained by the deethylation of 10-methyl-9-ethoxyphenanthrene, which in turn was obtained by the reaction of 9-fluorenone with excess diazoethane: mp 122–123 °C (lit.¹⁴ 122–123 °C); NMR (CDCl₃) δ 2.68 (s, 3 H), 5.32 (br s, 1 H), 7.50–8.8 (m, 8 H).

Monoethyl 1,1'-Diphenate (IVa). Diphenic anhydride was first prepared by treatment of 1,1'-diphenic acid (0.51 g, 0.0021 mol) with excess acetic anhydride according to the method reported by Roberts et al.¹⁵ The anhydride so obtained was employed in the syntheses of IVa without further purification.

The anhydride (0.38 g, 0.0017 mol) was added to a solution which was previously prepared by dissolving 1 g of Na in 10 mL of EtOH. The solution was poured into a mixture of 100 mL of 2 N HCl and 200 mL of dichloromethane after stirring for 10 min at room temperature. The organic layer was separated, washed with water twice, and dried over Na₂SO₄ overnight. Evaporation of the solvent gave a light yellow solid which was purified by column chromatography (silica gel) using CHCl₃-ether (1:1) as an eluent. The product eluted before impurities. Recrystallization from *n*-hexane twice afforded monoethyl 1,1'-diphenate (0.2 g); yield 34% based on diphenic acid used: mp 99–100 °C; NMR (CDCl₃) δ 0.95 (t, 3 H), 4.06 (q, 2 H), 7.10–7.60 (m, 6 H), 7.95–8.20 (m, 2 H), and 10.28 (br s, 1 H); UV (*t*-BuOH) λ_{max} 283 nm (ε₂₈₃ 3090 M⁻¹ cm⁻¹).

2-(2'-Acetylphenyl)benzoic acid (IVb) was prepared after a method of Reid and Conte.¹⁶ IVb was synthesized by reaction of 95% formic acid with 7-oxo-5-[bis(ethoxycarbonyl)methylene]-5,7-dihydrodibenz[*c,e*]oxepin, which was obtained by treatment of 1,1'-diphenic anhydride with diethyl malonate in the presence of triethylamine and acetic anhydride: mp 121 °C (lit.¹⁶ 120–121 °C); UV (*t*-BuOH) λ_{max} 288 nm (ε₂₈₈ 2920 M⁻¹ cm⁻¹); NMR (CDCl₃) δ 2.22 (s, 3 H), 7.00–8.10 (m, 8 H), 10.37 (s, 1 H).

10-Hydroxy-10-methyl-9,10-dihydro-9-phenanthrone (IIIb). Rapid bubbling of O₂ through a solution of Ib⁻ (0.209 g, 0.001 mol) in 5 mL of ethanol containing KOH (0.1 g in 5 mL of H₂O) for 10 min yielded (after pouring into ice-water, collecting, and recrystallization from *n*-hexane) a 60% yield of IIIb: mp 86 °C (lit.¹⁷ 87 °C); UV (*t*-BuOH) λ_{max} 326 nm (ε 3050 M⁻¹ cm⁻¹), 278 (10 100), and 270 (9500); NMR (CDCl₃) δ 1.54 (s, 3 H), 3.97 (s, 1 H), 7.18–8.10 (m, 8 H).

1-Hydroperoxy-1,2,3,4-tetrahydronaphthalene (VI). VI was prepared by autoxidation of tetrahydronaphthalene initiated by AIBN at 60 °C for 48 h. Purity of the hydroperoxide was determined by conventional iodometric titration (97%).

2,6-Di-*tert*-butyl-4-hydroperoxy-4-methyl-2,5-cyclohexadien-1-one (VII). VII was synthesized by the method of Kharasch and Joshi.¹⁸ Iodometric titration showed that the hydroperoxide was 85% pure.

Product Analysis. In a typical experiment, solutions of Ia or Ib containing the appropriate concentrations of *t*-BuO⁻K⁺ were prepared in a glovebox under dry N₂. Reactions were initiated by the addition of an aliquot (5 mL) of the basic solution of Ia or Ib to a weighed portion of solid 4a-FIEtOOH. Typical concentrations were as follows: [Ia] = 7.8 × 10⁻³ M, [*t*-BuO⁻K⁺] = 7.7 × 10⁻³ M, and [4a-FIEtOOH] = 3.3 × 10⁻⁴ M. After the mixture was stirred for several minutes, 0.05 mL of glacial acetic acid was added. In order to determine the yield of FIEt⁻ a portion of the acidified solution was transferred to a Thunberg cuvette and mixed with a solution of the nitroxide radical (4-hydroxy-2,2,6,6-tetramethylpiperidin-1-oxyl) which is known to convert FIEtH to FIEt⁻.⁶ The concentration of FIEt⁻ (= [FIEtH]) was determined by its absorbance at 640 nm (ε₆₄₀ 5000 M⁻¹ cm⁻¹).⁷ λ_{max} and extinction coefficients of products follow: IX, 398 nm (ε 1890 M⁻¹ cm⁻¹); V, 414 nm (ε 1620 M⁻¹ cm⁻¹); IVa, 283 nm (ε 3090 M⁻¹ cm⁻¹); IVb, 288 nm (ε 2920 M⁻¹ cm⁻¹); IIIb, 278 nm (ε 10 100 M⁻¹ cm⁻¹).

The yield of products from the substrates was determined by LC.

Table I. Reaction of 4a-FIEtOOH with 10-Methyl-9-phenanthrol in the Presence of *tert*-Butoxide

| 4a-FIEtOOH, ^a ×10 ⁴ M | Ib, ×10 ³ M | <i>t</i> -BuOK, solvent reaction time, | | products, % ^b | |
|--|---------------------------|--|---------------------------------|--------------------------|------|
| | | ×10 ³ M | min | FIEtH | IIIb |
| 3.14 | 7.00 | 6.82 | <i>t</i> -BuOH | 69 | 35 |
| 3.14 | 7.00 | 6.82 | <i>t</i> -BuOH | 53 | 42 |
| 2.77 | 7.98 | 7.93 | <i>t</i> -BuOH | 57 | 59 |
| 2.72 | 8.98 | 6.91 | <i>t</i> -BuOH | 46 | 65 |
| 3.86 | 8.55 | 11.6 | <i>t</i> -BuOH | 57 | 28 |
| 1.79 | 7.85 | 7.61 | <i>t</i> -BuOH | 40 | 89 |
| 3.63 | 8.02 | 7.37 | CH ₃ CN ^c | 13 | 52 |






^a The purity of 4a-FIEtOOH used was 85–93%. ^b The yields of FIEtH and IIIb were based on 4a-FIEtOOH employed. ^c No formation of IVb was observed by LC analysis.

The LC analysis was carried out with a Du Pont Instrument reverse phase column (Lichrosorb 5RP-8, 25 cm, 4.6 mm), using acetonitrile–acetic acid–water, 25:25:50 (v/v), and acetonitrile–water, 40:60 (v/v), as solvents for Ia and Ib, respectively, at a flow rate of 1.2 mL/min. The products were monitored at 283 nm (=λ_{max} of IVa and IVb). The retention times (using CH₃CN–AcOH–H₂O) of Ia, V, and IVa, were 26.5, 6.6, and 9.5 min, respectively, and the retention times (using CH₃CN–H₂O) of Ib, IVb, IIIb, and IIIb were 35.0, 6.8, 11.3, and 12.3 min, respectively.

Results

The kinetics for the reaction of 4a-FIEtO₂[−] with the phenanthrene substrates of this study could not be followed spectrophotometrically. This is due to the fact that the absorbances of substrate anions under the condition of [substrates[−]] ≫ [4a-FIO₂[−]] exceeded greatly the absorbance of 4a-FIEtO₂[−]. For this reason, the oxygen-transfer reactions were allowed to proceed for a set time period (generally 10 min), the reaction was quenched, and products were determined. Kinetic analyses of partial reactions were feasible and have been carried out. A means was designed to follow dioxygen transfer from 4a-FIEtO₂[−] to 3,5-di-*tert*-butylcatechol. All reactions were investigated at 30 °C under strictly anaerobic conditions and unless stated to the contrary in absolute *t*-BuOH.

10-Methyl-9-phenanthrol anion (Ib[−]) reacts with molecular oxygen in strongly basic ethanol to yield 10-hydroxy-10-methyl-9,10-dihydro-9-phenanthrone (IIIb). The intermediate peroxide anion (IIIb[−]) is too unstable under basic conditions for its isolation (eq 4). The experimental results obtained for the reaction of Ib[−] with 4a-FIEtO₂[−] are tabulated in Table I.



The products of reaction were found to be FIEt[−] and IIIb. No evidence for the presence of the 10-hydroperoxy compound IIB could be obtained by LC and comparison to authentic IIB. The inability to detect IIB is likely due to its instability in the basic reaction solution and possibly also to its reduction by FIEt[−]. The fact that FIEt[−] is not consumed in the reaction (Table I) may be explained by the finding that both FIEt[−] and FIO_x⁺Et are reduced to FIEt[−] by Ib[−] which is present in excess (vide


Table II. Reaction of 4a-FlEtOOH with 10-Ethoxy-9-phenanthrol in Various Solvents

| 4a-FlEtOOH, ^a ×10 ⁴ M | Ia, ×10 ³ M | <i>t</i> -BuOK, ×10 ³ M | solvent | reaction time, min | products, % ^b | | |
|--|---------------------------|---------------------------------------|--------------------|-----------------------|--------------------------|----|--------------|
| | | | | | FlEt | V | IVa |
| a. 5.70 | 6.43 | 6.03 | CH ₃ CN | 10 | FlEt: 32 | 53 | 43 |
| b. 3.73 | 7.86 | 7.47 | CH ₃ CN | 10 | FlEt: 20 | 45 | 29 |
| c. 2.95 | 6.50 | 6.79 | CH ₃ CN | 10 | FlEt: 22 | 36 | 32 |
| d. 3.98 | 5.77 | 5.99 | <i>t</i> -BuOH | 20 | 58 | 66 | ^c |
| e. 3.25 | 7.70 | 8.04 | <i>t</i> -BuOH | 20 | 54 | 92 | ^c |
| f. 3.33 | 7.98 | 7.86 | <i>t</i> -BuOH | 30 | 50 | 86 | 0.7 |
| g. 3.33 | 7.89 | 7.86 | <i>t</i> -BuOH | 150 | 28 | 94 | 0.9 |
| h. 4.52 | 6.06 | 5.88 | THF | 10 | no data | 51 | ^c |
| i. 4.52 | 6.06 | 5.88 | THF | 120 | no data | 98 | ^c |
| j. 5.04 | 8.00 | 7.96 | dioxane | 30 | no data | 78 | 10 |

^a 4a-FlEtOOH used was 85–92% pure. ^b The yields were based on 4a-FlEtOOH employed. ^c No formation of IVa was detected by LC.

infra). The reactions of the hydroperoxides VI and DTBP-O₂H with FlEt⁻ were investigated in *t*-BuOH under anaerobic conditions with [organic peroxide] = 3.6×10^{-3} M > [FlEt⁻] = 7.8×10^{-4} M. The products in each case were determined to be FlEt· (λ_{max} 640 nm) and the corresponding alcohol (LC comparison to authentic alcohol).


In the reaction of 4a-FlEtO₂⁻ with Ib⁻ special attention was given to the presence of 2-(2'-acetylphenyl)benzoic acid (IVb) in the course of LC analysis in which instance authentic IVb was employed. No trace of this compound could be detected. This negative result finds ready interpretation in the reactions of eq 9. Thus, the decomposition of IIb⁻ → IIIb⁻ is a rapid reaction and dioxetane formation, with its accompanying 40 kcal mol⁻¹ thermodynamic barrier,¹⁹ should be slow when compared to the rate of opening of the dioxetane ring by elimination to provide IIb⁻.



10-Ethoxy-9-phenanthrol anion (Ia⁻) reacts with 4a-FlEtO₂⁻ to provide 9,10-phenanthroquinone (V) and monoethyl 1,1'-diphenate (IVa). The ratio of V:IVa is dependent upon the polarity of the solvent (Table II). In acetonitrile an almost equal yield of both components is obtained but little or no IVa is obtained in dioxane, THF, or *t*-BuOH. Formation of products from the substrate Ia⁻ is accountable through the intermediacy of a common peroxy anion. Yields of FlEt⁻ (nitroxide assay, see Experimental Section) of about 60% based on [4a-FlEtO₂H]_{initial} were obtained in *t*-BuOH and THF. Longer reaction times were advantageous to the production of V but resulted in lower yields of FlEt⁻ (~30%).

For the reaction carried out in acetonitrile solvent, the radical FlEt· appeared on acidification with acetic acid without addition of nitroxide. In run a the yield of FlEt· amounted to 35%. A like appearance of FlEt· on acidification of the spent solution resulting from the reaction of VIII⁻ with 4a-FlEtO₂⁻ was also observed. The explanation afforded (vide infra) for this observation is applicable to the present case.


3,5-Di-*tert*-butylcatechol anion (VIII⁻) at 4×10^{-3} M reacts with 4a-FlEtO₂⁻ ([4a-FlEtO₂H] = 1.2×10^{-4} M;




[*t*-BuO⁻K⁺] = 4×10^{-3} M) in *t*-BuOH under anaerobic conditions. Acidification of the reaction solution by a small amount of deoxygenated acetic acid provided FlEt· and 3,5-di-*tert*-butyl-*o*-quinone (IX) in yields of ~70 and 60%, respectively. The yields of product were not sensitive to the alteration in [*t*-BuO⁻K⁺] of fourfold employed. Product analysis (LC) indicated the absence of such known oxidation products of VIII as γ -lactone, 3,5-di-*tert*-butyl-5-(carboxymethyl)-2-furanone, and 3,5-di-*tert*-butyl-5-(carboxyhydroxymethyl)-2-furanone.²⁰

On completion of the reaction of VIII⁻ with 4a-FlEtO₂⁻, and prior to acidification, the reaction solution possessed a pronounced blue color. The visible spectrum of the reaction solution beyond 390 nm proved to be identical (λ_{max} 730 nm) with that of 3,5-di-*tert*-butyl-*o*-benzosemiquinone (which was generated independently from VIII by disproportionation with an equal concentration of the quinone IX). In addition, the blue solution exhibited no absorbance at 490 and 398 nm, indicating the absence of FlEt· and the quinone IX. On acidification of the reaction solution the absorbance due to the di-*tert*-butyl-*o*-benzosemiquinone disappeared and in its place there was found to be absorbance by FlEt·. These observations may be explained by the reactions of Scheme I. Reaction c, Scheme I, was studied in a separate experiment by reacting VIII⁻ (4×10^{-3} M), IX (4×10^{-4} M), and *t*-BuO⁻K⁺ (8×10^{-3} M), which provided the semiquinone (λ_{max} 730 nm, ε 680 cm⁻¹ M⁻¹) in 100% yield (no IX remained). By use of the extinction coefficient of the semiquinone it was then calculated that this species was found in 96% yield in the reaction between 4a-

Scheme I




FIET^- and VIII^- . Reaction e of Scheme I was also established in a separate experiment in which a solution of the semiquinone (8.22×10^{-4} M) in *t*-BuOH containing $t\text{-BuO}^- \text{K}^+$ was acidified with acetic acid to provide a 48% yield of IX. On comproportionation the semiquinone should produce a 50% yield of IX. Reaction d of Scheme I could also be established in a separate experiment. When a *t*-BuOH solution of FIET^- (1.93×10^{-4} M) containing semiquinone (4.24×10^{-4} M) and $t\text{-BuO}^- \text{K}^+$ (8.52×10^{-3} M) was acidified with acetic acid, there were immediately formed FIET^- (1.83×10^{-4} M) in 95% yield and IX (1.03×10^{-4} M) in 43% yield. Since the reactions of eq d and e would both be in operation under the concentrations used (i.e., eq 11) the yield of quinone IX is






86% of theory. From Scheme I, the theoretical yield of IX is 50% and that of FIET^- is 100%. These expectations may be compared to the determined product yields of 60% for IX and 70% for FIET^-

Summing reactions a, b, and c of Scheme I provides eq 12.



Since reactions b and c are very rapid the transfer of dioxygen from $4a\text{-FIET}^-$ to VIII^- (reaction a, Scheme I) could be followed by monitoring the appearance of semiquinone with time at 730 nm. In performing the stopped-flow kinetic experiments one drive syringe contained a *t*-BuOH solution of VIII and $t\text{-BuO}^- \text{K}^+$ at equal concentrations and the other syringe contained a *t*-BuOH solution of $4a\text{-FIET}^-$ at 1.45×10^{-4} M. At $[\text{VIII}^-] \gg [4a\text{-FIET}^-]$ the appearance of semiquinone followed the first-order (k_{obsd}) rate law. The kinetic study was repeated by mixing on the stopped-flow bench a solution of $4a\text{-FIET}^-$ (1.45×10^{-4} M) and solutions of VIII (0.324 to 6.48×10^{-3} M) plus $t\text{-BuO}^- \text{K}^+$ at a concentration of 6.64×10^{-3} M. Plots of k_{obsd} (determined by either procedure) vs. $[\text{VIII}^-]$ exhibit saturation in $[\text{VIII}^-]$ and a plot of $1/k_{\text{obsd}}$ vs. $1/[\text{VIII}^-]$ is linear (Figure 1). This observation establishes the presence of an intermediate formed from $4a\text{-FIET}^-$ plus VIII^- (see Discussion).

Reaction of phenolate ions with N^5 -ethyl-3-methylflavin radical (FIET^-). were investigated by stopped-flow spectrophotometry under an inert atmosphere using *t*-BuOH as solvent. All reactions were carried out by mixing a solution of phenolate ion (phenol plus equivalent $t\text{-BuO}^- \text{K}^+$) with one containing FIET^- prepared by mixing equimolar solutions of $\text{FI}_{\text{ox}}^+ \text{Et}$ and FIET^- at 2×10^{-4} M. Phenolate ion was employed in excess of FIET^- and under these conditions the disappearance of FIET^- (640 nm) followed good pseudo-first-order kinetics and plots of k_{obsd} vs. [phenolate anion] were found to be linear. From the slopes of the plots of Figure 2 there were calculated the second-order rate constants of eq 13-15.



Reactions of phenolate ions with O_2 were allowed to proceed in absolute *t*-BuOH containing $t\text{-BuO}^- \text{K}^+$ and continually saturated with air (constant $[\text{O}_2] \approx 3.15 \times 10^{-3}$ M) at 30 °C for a set period of time. The reactions were then quenched by the addition of acetic acid to slight acidity and product analysis was performed. Results are presented in Table III. The percentage yields reported in Table III are based upon the initial $[\text{O}_2]$ and must be considered maximal since $[\text{O}_2]$ is constant.

The kinetics for the reaction of O_2 with VIII^- were investigated by monitoring the appearance of semiquinone at 730 nm (eq 16). Reactions were initiated by rapid mixing a solution

Table III. Reaction of Phenolate Ions with Oxygen (30 °C, *t*-BuOH)^a

| | substrate, ×10 ³ M | [<i>t</i> -BuO ⁻] ×10 ³ M | reaction time, min | product % yields ^b | | | |
|---|----------------------------------|--|-----------------------|-------------------------------|---------------------------|---------------------------|------|
| | | | | IX | V | IVa | IIIb |
| 1 | Ia (8.05) | 7.99 | 10 | | <12 | 0 | |
| 2 | Ia (8.04) | 7.85 | 30 | | <33 | trace | |
| 3 | Ia (7.03) | 6.93 | 10 | | <8.9 × 10 ⁻⁴ M | <3.4 × 10 ⁻⁴ M | |
| 4 | Ib (7.98) | 7.70 | 10 | | 0 | | <9 |
| 5 | VIII (4.01) | 3.81 | 10 | <47 | | | |
| 6 | VIII (4.45) | 7.83 | 10 | <46 | | | |

^a The solvent for entry 3 is acetonitrile. ^b Percent yield based on initial [³O₂].




Figure 1. Plot of the reciprocal of the pseudo-first-order rate constants ($1/k_{\text{obs}}$) vs. the reciprocal of the concentration of 3,5-di-*tert*-butylcatechol (VIII) for the reaction of VIII⁻ with 4a-FlEtO₂⁻ (1.45×10^{-4} M). Reaction was carried out in absolute *t*-BuOH under inert atmosphere with [*t*-BuOK] = 6.64×10^{-3} M at 30 °C.






Figure 2. Plot of pseudo-first-order rate constants (k_{obs}) vs. phenolate ion concentrations for the $1e^-$ reduction of *N*⁵-ethyl-3-methylflavin radical ([FlEt⁻] = 9.80×10^{-5} M, *t*-BuOH solvent, 30 °C).



containing equimolar concentrations of VIII and *t*-BuO⁻K⁺ in absolute *t*-BuOH with an air-saturated ($[\text{O}_2] = 3.15 \times 10^{-4}$ M) solution of *t*-BuOH. With [VIII⁻] in excess over [O₂] there was obtained good pseudo-first-order kinetics. The second-order rate constant for reaction of O₂ with VIII⁻ was determined ($2.32 \text{ M}^{-1} \text{ s}^{-1}$) from a plot (Figure 3) of $k_{\text{obs}}(\text{s}^{-1})$ vs. [VIII⁻]. This value may be compared to the second-order rate constant ($0.348 \text{ M}^{-1} \text{ s}^{-1}$) for the reaction of DTBP⁻ with O₂ under the same conditions.⁷

Discussion

A result of this study which is of biochemical relevance is the finding that the 4a-hydroperoxide obtained from the reaction of an *N*⁵-alkylflavin with ³O₂ is capable of transferring both oxygen atoms to a catechol monoether anion to provide the monoester of a dicarboxylic acid (eq 17) with the regeneration of reduced flavin. We have, in effect, discovered a dihydroflavin-catalyzed dioxygenase reaction. The oxidation of Ia⁻ to provide IVa undoubtedly occurs through the intermediacy of a peroxide formed upon dioxygen transfer from 4a-FlEtO₂⁻ to Ia⁻, followed by a Hock-type rearrangement (eq 10). When the OEt substituent of IX is replaced by Me, the resultant peroxide (IIb) does not provide a ring-cleavage product (IVb) but decomposes to a tertiary alcohol, IIIb (eq 4). This may be accounted for by (1) the instability of IIb in base; (2) the greater stability of X as compared to XI; (3) the inoperability of the dioxetane path of eq 9. Rearrangements of peroxide or perester intermediates similar to that of eq 10 have been suggested for the oxidative cleavage of phenols,²¹



β -naphthoquinone,²² and *trans*-decalin peresters.²³ Hamilton²⁴ has previously provided thermodynamic arguments in favor of this type of mechanism over one involving the formation and cleavage of a dioxetane. In addition, the presently favored mechanism for intradiol metallocatechol dioxygenases






Figure 3. Plot of the pseudo-first-order rate constants (k_{obs}) vs. the concentration of 3,5-di-*tert*-butylcatecholate anion (VIII^-). Oxygen concentration is 3.15×10^{-4} M in *t*-BuOH (30 °C).

involves rearrangement through anhydride intermediate.^{25,26}

Of key interest is the mechanism whereby the peroxy anion moiety of $4a\text{-FIEtO}_2^-$ is transferred to a phenolate anion. The first example of this reaction is shown in eq 1.⁷ The reaction was found to be first order in both $[4a\text{-FIEtO}_2^-]$ and $[\text{DTBP}^-]$ at low values of the latter and to be independent of $[\text{DTBP}^-]$ at its higher molarity. In the original study⁷ a *t*-BuOH solution of $4a\text{-FIEtO}_2^-$ was stopped flow mixed with an excess of the phenol DTBP in *t*-BuOH. In a series of kinetic runs the mole fraction of DTBP present as DTBP^- was increased by including increasing concentrations of *t*-BuO⁻K⁺. To be certain that saturation in $[\text{DTBP}^-]$ was not due to a salt effect (or saturation by trace of metal) the kinetic studies of this reaction have now been repeated with a constant value of $[\text{K}^+]$. This was accomplished by stopped-flow mixing of a solution of $4a\text{-FIEtO}_2^-$ (5.85×10^{-5} M) with solutions containing *t*-BuO⁻K⁺ at a constant concentration of 6.64×10^{-3} M and varying concentrations of DTBP (0.324 to 6.48×10^{-3} M). Again, saturation in DTBP^- was obtained. Plots of $1/k_{\text{obs}}$ vs. $1/\text{DTBP}^-$ were found to be indistinguishable when employing both sets of experimental data (Figure 4). Thus, the dioxygen transfer reaction is not unduly sensitive to the change in μ and the formation of an intermediate species from $4a\text{-FIEtO}_2^-$ is verified. An intermediate might arise by either the covalent combination or complexation of $4a\text{-FIEtO}_2^-$ and DTBP^- (eq 18) or in a reaction involving only $4a\text{-FIEtO}_2^-$ (eq 19):



No reasonable structures are apparent for an intermediate compound which could be formed between $4a\text{-FIEtO}_2^-$ and DTBP^- and it was argued⁷ that complex formation between the two negatively charged species would not be favorable in *t*-BuOH. In addition, evidence of a kinetic nature could not be found for the complexation of the neutral phenol with neutral $4a\text{-FIEtO}_2\text{H}$. In this study we find that the reaction of VIII^- with $4a\text{-FIEtO}_2^-$ is also first order in $[\text{VIII}^-]$ at its lower values, becoming independent of $[\text{VIII}^-]$ at higher concentrations. Assuming the sequence of eq 19 with X_2 at steady state provides the equation

$$1/k_{\text{obs}} = k_2/k_1k_3[\text{I}^-] + 1/k_1 \quad (20)$$

Plots of $1/k_{\text{obs}}$ vs. $1/[\text{VIII}^-]$ and $1/k_{\text{obs}}$ vs. $1/[\text{DTBP}^-]$ are provided in Figures 1 and 4. The derived rate constants are given in Table IV. Within experimental error the values of k_1 determined with DTBP^- and VIII^- are identical. This finding





Figure 4. The reciprocal of the pseudo-first-order rate constants ($1/k_{\text{obs}}$) vs. the reciprocal of the concentration of the anion of 2,6-di-*tert*-butyl-4-methylphenol ($1/[\text{DTBP}^-]$) for the reaction of DTBP^- with the anion of *N*⁵-ethyl-4*a*-hydroperoxy-3-methylflavin ($4a\text{-FIEtO}_2^-$) in *t*-BuOH at 30 °C: (●) mixing of a solution of $4a\text{-FIEtO}_2\text{H}$ (5×10^{-5} M) with solutions of *t*-BuOK in the presence of DTBP (5.5×10^{-3} M) (ref 7); (Δ) mixing of a solution of $4a\text{-FIEtO}_2\text{H}$ (6×10^{-5} M) with solutions of DTBP containing *t*-BuOK at 6.6×10^{-3} M.

Table IV


| | k_1, s^{-1} | $k_2/k_1k_3, \text{s M}$ | $k_2/k_3, \text{M}$ |
|--|----------------------|--------------------------|----------------------|
| | 0.36 | 6.2×10^{-4} | 2.2×10^{-4} |
| | 0.37 | 7.7×10^{-4} | 2.8×10^{-4} |

supports the sequence of eq 19. It is of some little surprise to find that the values of the partition coefficients (k_2/k_3) determined with both phenolate ions are also very similar. Assuming that eq 19 represents the sequence of events, k_2 is common to both reactions and VIII^- undergoes reaction with X_2 about 1.3 times slower than does DTBP^- . With the liberal assumption that the finding with DTBP^- and VIII^- of a common value of k_1 will prove general, one might draw the conclusion that the sequence of eq 19 is correct and that $4a\text{-FIEtO}_2^-$ is spontaneously converted into an intermediate or intermediates which peroxidize the phenolate anions with about the same rate. The various possibilities will be considered.

Migration of the peroxy anion moiety from the 4*a* to the 10*a* position to yield X_2 as a 10*a*-peroxyflavin (XIII) would be




highly endothermic and should not be expected. This view is supported by the observation that the rate constants for reaction of the 4a-peroxy anion of 10-(2',6'-dimethylphenyl)-5-ethyl-3-methylisoalloxazine (XIV) and 4a-FIEtO₂⁻ with DTBP⁻ do not differ greatly.²⁷ The 10a position of XIII is sterically hindered²⁸ to nucleophilic attack and rearrangement to the 10a-peroxyflavin would be even more endothermic than in the rearrangement of 4a-FIEtO₂⁻. The hypothetical dioxetane (XII) may be considered as X₂. The structure of XII is sterically feasible even when a bulky substituent, such as the 2',6'-dimethyl moiety, occupies the 10 position. This is due to the fact that the >C=C< bond between C_{4a} and C₄ provides a butterfly configuration much as in dihydroflavin and steric compression between the 10a and 10 substituents is greatly reduced. Nucleophilic attack of PhO⁻ upon the dioxetane XII




to produce ⁻OOPh=O would require C—O rather than O—O bond scission. We know of no instances in which the opening of a dioxetane has been shown to involve C—O bond cleavage. However, the structure of XII is rather unique possessing, as it does, an electron sink in the >C=O function at the 4a position of the isoalloxazine ring.

In the mechanism of eq 19 the species X₂ might represent (O₂ + FIEt⁻) or (O₂⁻ + FIEt⁻). The formation of these intermediates could occur simply by the reversal of the reaction of ³O₂ with 1,5-dihydroflavin:⁵




If X₂ represented completely dissociated FIEt⁻ + O₂, the peroxidation would take place in a second-order reaction between phenolate anion and O₂. This possibility was eliminated in the previous study by an independent determination of the rate constant for the reaction of eq 23b. The second-order rate



constant so obtained was ~100-fold less than required to be compatible with the observed kinetics for peroxidation of DTBP⁻ by 4a-FIEtO₂⁻. From the second-order rate constant for reaction of O₂ with VIII⁻ and a knowledge of k₁ for X₂ formation it can be calculated that the rate of reaction of O₂ with VIII⁻ is ca. 40-fold too slow to allow the kinetic competency of X₂ = FIEt⁻ + O₂ in the peroxidation of VIII⁻. Though the second-order rate constants for the reaction of Ia⁻

and Ib⁻ with O₂ were not obtained, arguments against the reaction of these phenolate ions with O₂ being involved in their peroxidation by 4a-FIEtO₂⁻ can be made. At a constant concentration of O₂ equal to ten times that of 4a-FIEtO₂⁻ employed Ib⁻ provides IIIb in <9% of theory based upon [O₂]_{initial} (see Results for details). No other product could be detected. Given that the peroxidation reactions with 4a-FIEtO₂⁻ and O₂ were carried out for the same period of time and that the concentration of O₂ was at a constant concentration which exceeded that for the initial concentration of 4a-FIEtO₂⁻ by >tenfold, the efficiency of peroxidation by 4a-FIEtO₂⁻ exceeds that for O₂ by >100-fold. The reaction of Ia⁻ with O₂ under like conditions of concentration provided <12% yield of V in 10 min and a <33% yield of V in 30 min. No diphenic acid monoester (IVa) could be detected in the 10-min run and only a possible trace of this substance could be detected after 30 min. Since the steady-state oxygen concentration exceeded the initial concentration of 4a-FIEtO₂⁻ employed by >tenfold one may conclude that the efficiency of the reactions with oxygen is <3–4% than seen with 4a-FIEtO₂⁻. In acetonitrile Ia⁻ reacts with O₂ to yield both V and IVa. Thus, both 4a-FIEtO₂⁻ and O₂ provide with Ia⁻ only V in *t*-BuOH but both V and IVa in acetonitrile. Like experiments with VIII⁻ establish that the efficiency of the reaction of this phenolate anion with O₂ amounts to <4–5% of that seen with 4a-FIEtO₂⁻. The observation that the efficiency of O₂ as a peroxidizing agent, for the various phenolate anions, is far less than with 4a-FIEtO₂⁻ establishes that peroxidation by FIEtO₂⁻ is not the result of the dissociation of a 4a-FIEtO₂⁻ into FIEt⁻ + O₂ and subsequent bimolecular reaction of phenolate ion with the free O₂ species (eq 23).


The possibility that the reaction sequence involves eq 19 with X₂ representing solvent-separated FIEt⁻ and O₂⁻ should be considered:




To account for the saturation in the pseudo-first-order rate constant by phenolate anion it would be assumed that eq 24c would be rate limiting at low [phenolate] and eq 24a rate limiting at high [phenolate]. The reactions 24b and 24c may be studied independently.

All the phenolate ions which have been shown to be oxidized by 4a-FIEtO₂⁻ have also been shown to reduce FIEt⁻ to FIEt⁻. The second-order rate constants (*t*-BuOH) follow: DTBP⁻, minimally 10⁷ M⁻¹ s⁻¹;⁷ Ia⁻, 1.02 × 10³ M⁻¹ s⁻¹; Ib⁻, 9.97 M⁻¹ s⁻¹; VIII⁻, 1.95 × 10⁴ M⁻¹ s⁻¹. At phenolate concentrations employed in the reaction with 4a-FIEtO₂⁻ the pseudo-first-order rate constants for phenolate ion reduction of FIEt⁻ may be calculated as: Ia (at 6 × 10⁻³ M), 6.12 s⁻¹; Ib (at 7 × 10⁻³ M), 6.98 × 10⁻² s⁻¹; VIII (at 4 × 10⁻³ M), 7.8 × 10¹ s⁻¹. The rate constant for spontaneous decomposition of 4a-FIEtO₂⁻ has been determined⁷ to be 4.6 ± 0.2 × 10⁻² s⁻¹ (*t*-BuOH). These results establish that, at the concentrations of the various phenolate ions used in the reactions with 4a-FIEtO₂⁻, the rate of reduction of FIEt⁻ by Ia⁻, VIII⁻, and DTBP⁻ is more than adequate to assure that eq 24b could be a kinetically competent component of the peroxidation reaction. In the case of Ib⁻, the rate of reduction of FIEt⁻ exceeds the rate of spontaneous decay of 4a-FIEtO₂⁻ by a factor of 1.5. This is sufficient to account for the 70% yield of FIEt⁻ obtained in the reaction of 4a-FIEtO₂⁻ with Ib⁻. The coupling of O₂⁻ with phenol radical appears to be the weak link in the overall mechanism of eq 24. Thus, it has been reported²⁹ that in the presence of excess O₂⁻ · K⁺ (crown) the radical DTBP⁻ is re-


duced to DTBP⁻. There appears to be in the literature no documented example of O₂⁻· coupling with a radical species to yield a peroxide.³⁰ Formation of the phenol-derived peroxides would not, so it appears, result from the reaction of molecular oxygen derived from 4a-FlEtO₂⁻ with phenolate anion nor from the coupling of the radical, derived by 1e⁻ abstraction from phenolate anion, with superoxide. The intermediate X₂ must, therefore, represent a compound or complex of oxygen and dihydroflavin. The dioxetane mechanism of eq 21 must be considered as a possibility. Another possibility is that phenolate ion reacts with an O₂ species which has not dissociated from the flavin moiety (eq 25).



The oxygen would not, in this instance, be as closely associated with the flavin as in the dioxetane XII and might possess some singlet character. Charge-transfer complexes of oxygen have been proposed to be involved in bimolecular and intramolecular oxygen transfer reactions.³¹ Because the values of *k*₃ determined for DTBP⁻ and VIII⁻ are quite similar the capture of X₂ by phenolate ion may be diffusion controlled.



An alternative would involve the preassociation of 4a-FlEtO₂⁻K⁺ with PhO₂⁻K⁺ in a loose complex within a solvent cage. Dissociation of 4a-FlEtO₂⁻ within this solvent cage could then result in the peroxidation reaction prior to solvent rearrangement (eq 27). If this were the case, then the equilibrium



constant for complex formation would be only slightly dependent upon the nature of the phenolate anion (DTBP⁻ vs. VIII⁻) while the equilibrium constants for complex formation would be 5.85 × 10² for DTBPK⁺ and 3.63 × 10² for VIII⁻K⁺.

Acknowledgment. This work was supported by grants provided by the National Institutes of Health and the National

Science Foundation. Suggestions and comments by Professors D. Sawyer and W. H. Okamura are gratefully acknowledged.

References and Notes

- (1) Kemal, C.; Bruice, T. C. *Proc. Natl. Acad. Sci. U.S.A.* **1976**, *73*, 995.
- (2) Kemal, C.; Chan, T. W.; Bruice, T. C. *Proc. Natl. Acad. Sci. U.S.A.* **1977**, *74*, 405.
- (3) Chan, T. W.; Bruice, T. C. *J. Am. Chem. Soc.* **1977**, *99*, 2387.
- (4) Kemal, C.; Bruice, T. C. *J. Am. Chem. Soc.* **1977**, *99*, 7064.
- (5) Kemal, C.; Chan, T. W.; Bruice, T. C. *J. Am. Chem. Soc.* **1977**, *99*, 7272.
- (6) Chan, T. W.; Bruice, T. C. *J. Am. Chem. Soc.* **1977**, *99*, 7287.
- (7) Kemal, C.; Bruice, T. C. *J. Am. Chem. Soc.* **1979**, *101*, 1635.
- (8) Ball, S.; Bruice, T. C. *J. Am. Chem. Soc.* **1979**, *101*, 4017.
- (9) Hayaishi, O.; Katagiri, M.; Rothberg, S. *J. Am. Chem. Soc.* **1955**, *77*, 5450.
- (10) Nishinaga, A. *Chem. Lett.* **1975**, 273. Nishinaga, A.; Tojo, T.; Matsuura, T. *J. Chem. Soc., Chem. Commun.* **1974**, 896. Tsuji, J.; Takayanagi, H. *J. Am. Chem. Soc.* **1974**, *96*, 7449. Tsuji, J.; Takayanagi, H.; Sukai, J. *Tetrahedron Lett.* **1975**, 1245. Rogic, M. M.; and Demmin, T. R. *J. Am. Chem. Soc.* **1978**, *100*, 5472. Brown, D. G.; Beckmann, L.; Ashby, C. H.; Vogel, G. C.; Reinprecht, J. R. *Tetrahedron Lett.* **1977**, 1363. Matsuura, T.; Matsushima, H.; Kato, S.; Saipo, I. *Tetrahedron* **1972**, *28*, 5119. Moro-Oka, Y.; Foote, C. S. *J. Am. Chem. Soc.* **1976**, *98*, 1510. Lee-Ruff, E.; Lever, A. B. P.; Rigaudy, J. *Can. J. Chem.* **1976**, *54*, 1837. Ferrand, J. C.; Johnson, D. C. *J. Org. Chem.* **1971**, *36*, 3606. Pandell, B. L. *ibid.* **1976**, *41*, 3922.
- (11) Spencer, T.; Massey, V. *J. Biol. Chem.* **1972**, *247*, 5632. Entsch, B.; Ballou, D. P.; Massey, V. *ibid.* **1976**, *251*, 2550. Stickland, S.; Massey, V. *ibid.* **1973**, *248*, 2953. Hastings, J. W.; Balny, C.; Le Peuch, C.; Dousou, P. *Proc. Natl. Acad. Sci. U.S.A.* **1973**, *70*, 3468.
- (12) Sparrow, L. G.; Ho, P. P. K.; Sundaram, T. K.; Zach, D.; Nyns, F. J.; Snell, E. E. *J. Biol. Chem.* **1969**, *244*, 2590.
- (13) Müller, E.; Schurr, K.; Scheffler, J. *Justus Liebigs Ann. Chem.* **1959**, 627, 132.
- (14) Eistert, B.; El-Chahawi, M. A. *Monatsh. Chem.* **1967**, *98*, 941.
- (15) Roberts, R. C.; Johnson, T. B. *J. Am. Chem. Soc.* **1925**, *47*, 1399.
- (16) Reid, W.; Conte, R. *Chem. Ber.* **1971**, *104*, 1573.
- (17) Eistert, B.; Klein, L. *Chem. Ber.* **1968**, *101*, 900.
- (18) Kharasch, M. S.; Joshi, B. S. *J. Org. Chem.* **1957**, *22*, 1439.
- (19) O Neal, H. E.; Richardson, W. H. *J. Am. Chem. Soc.* **1970**, *92*, 6553.
- (20) Moro-oka, Y.; Foote, C. S. *J. Am. Chem. Soc.* **1976**, *98*, 1510.
- (21) Norman, R. O. C.; Smith, J. R. L. In "Oxidases and Related Redox Systems", King, T. E., Mason, H. S., Morrison, M., Eds.; Wiley: New York, **1965**; Vol. 1, p 131.
- (22) Karrer, P.; Schnelder, L. *Helv. Chim. Acta* **1947**, *30*, 859.
- (23) Crlegee, R.; Kaspar, R.; Dietrich, W. *Justus Liebigs Ann. Chem.* **1948**, 560, 127.
- (24) Hamilton, G. A. "Molecular Mechanisms of Oxygen Activation", Hayaishi, O., Ed.; Academic Press: New York, **1974**; p 405.
- (25) Que, L. Jr.; Lipscomb, J. D.; Münck, E.; Wood, J. M. *Biochim. Biophys. Acta* **1977**, *485*, 60.
- (26) Loehr, T. M.; Keyes, W. E.; Loehr, J. S. "Proceedings of the 3rd International Symposium on Oxidase and Related Oxidation Systems", King, T. E., Mason, H. S., Morrison, M., Eds.; University Park Press: Baltimore, in press.
- (27) Miller, A.; Bruice, T. C. *J. Chem. Soc., Chem. Commun.*, **1979**, 896; Bruice, T. C., Miller, A., *ibid.*, in press.
- (28) Smith, S. B.; Bruice, T. C. *J. Am. Chem. Soc.* **1975**, *97*, 2875.
- (29) Nishinaga, A.; Itahara, T.; Shimizu, T.; Tomita, H.; Nishizawa, K.; Matsuura, T. *Photochem. Photobiol.* **1978**, *28*, 687. Sawyer, D., private communication.
- (30) Wilshire, J.; Sawyer, D. *Acc. Chem. Res.* **1979**, *12*, 105.
- (31) As an example see: Nishinaga, A.; Shimizu, T.; Matsuura, T. *J. Org. Chem.* **1979**, *44*, 2983.