

to this quadrupole moment due to HCl depends on the effective dipole moment of HCl in the complex. The Q_{cc} components will be less dependent on the vibrational angle of acetylene. Using this second method, we get $Q(\text{acetylene}) = 7.6 \times 10^{-26}$ esu cm². These results are in reasonably good agreement with calculated values^{15,17} of $Q(\text{acetylene}) = 7.34$ or 7.2×10^{-26} esu cm².

(17) A. D. McLean and M. Yoshimine, *Natl. Bur. Stand. (U.S.), Tech. Note*, **438**, 67 (1967).

Acknowledgment. The support of the National Science Foundation and the University of Illinois School of Chemical Sciences is gratefully acknowledged. Acknowledgment is made to the donors of the Petroleum Research Fund, administered by the American Chemical Society, for partial support of this research. One of us (S.G.K.) thanks the University of Illinois Research Board for support. We wish to thank Peter Aldrich for the use of his computer programs for part of the data analysis.

Registry No. Acetylene-hydrochloric acid, 52218-20-9.

Rotational Barriers in Diphenylmethyl Anions Stabilized by Trimethylsilyl and Tricarbonylchromium(0) Moieties


Siden Top,[†] Gérard Jaouen,^{*†} Brian G. Sayer,[‡] and Michael J. McGlinchey^{*†}

Contribution from the *École Nationale Supérieure de Chimie, 75231 Paris Cedex 05, France, and the Department of Chemistry, McMaster University, Hamilton, Ontario, Canada L8S 4M1. Received March 15, 1983*


Abstract: A series of diphenylmethyl anions has been prepared in which the phenyl rings bear π -bonded $\text{Cr}(\text{CO})_3$ groups and/or a trimethylsilyl moiety. Variable-temperature ¹³C NMR studies on these compounds have allowed the evaluation of the barriers to phenyl rotation. It is shown that a π -bonded $\text{Cr}(\text{CO})_3$ group stabilizes the negative charge much more efficiently than does a p - SiMe_3 functionality. However, when the Me_3Si substituent is bonded directly to the α -carbon atom, the charge preferentially resides on the silicon, allowing the (π - $\text{Cr}(\text{CO})_3$ - C_6H_5) groups to rotate apparently unhindered.

The delocalization of negative charge from carbanionic centers is of paramount importance in many reaction sequences.^{1,2} It is well established that adjacent carbonyl functions,³ cyano⁴ or nitro⁵ groups, etc., very significantly stabilize carbanions such that the carbon formally bearing the negative charge is probably best regarded as being in an sp^2 -hybridized, trigonal-planar environment. In contrast, carbanions lacking neighboring groups that facilitate charge dispersal are presumably pyramidal and mimic amines in their behavior.

The interactions of trimethylsilyl groups with carbanions have been the subject of some debate;⁶ however, it was quickly recognized that such moieties, when directly bonded to the carbanionic center, lead to marked stabilization of the anion.⁷ A quantitative assessment of the stabilizing effect of a remote trimethylsilyl substituent was attempted via an ingenious NMR experiment described recently.⁸ In this study, the authors used DNMR to show conclusively that in the 4-methyl-4'-(trimethylsilyl)diphenylmethyl anion, **1**, the ortho carbons (and



protons) of the two phenyl rings are equilibrated via phenyl rotation at quite different rates. Indeed, the 4-methylphenyl ring is rotating faster than the 4'-(trimethylsilyl)phenyl moiety by a factor of 200 at 255 K. These data allowed the evaluation of the rotational barriers, which are ~ 10 kcal mol⁻¹ for the methyl-substituted ring and ~ 12.7 kcal mol⁻¹ for the silylated ring. The conclusions drawn were that the anion was essentially coplanar and that, assuming that the barrier was determined exclusively by the strength of π bonding to the benzylic carbon, the π interaction with the silylated ring was enhanced relative to that in the other ring. One possible rationale would involve $p\pi$ - $d\pi$ overlap, as in **2**, between the arene ring and a vacant d orbital



on silicon.⁶ Others prefer to attribute this stabilization to a polarization effect or to hyperconjugation.⁷

For some years now we have concerned ourselves with the capacity of the tricarbonylchromium(0) moiety to stabilize anions⁹ or cations¹⁰ at the benzylic position. Indeed, we have demonstrated the viability of a number of reactions for the (arene) $\text{Cr}(\text{CO})_3$ complexes not readily achieved for the noncomplexed arenes.^{11,12} We now report the syntheses and variable-temperature NMR spectra of diphenylmethyl anions bearing trimethylsilyl and/or tricarbonylchromium(0) substituents; this allows a direct comparison of the relative stabilizing effects of the two moieties.

(1) "Comprehensive Carbanion Chemistry", Buncl, E.; Durst, T., Eds.; Elsevier: Amsterdam, 1980.

(2) Cram, D. J. "Fundamentals of Carbanion Chemistry"; Academic Press: New York, 1965.

(3) Bartlett, P. D.; Woods, G. F. *J. Am. Chem. Soc.* **1940**, *62*, 2933.

(4) Bugg, C.; Desiderato, R.; Sass, R. L. *J. Am. Chem. Soc.* **1964**, *86*, 3157.

(5) Reference 2, p 52.

(6) (a) Peterson, P. J. *Organomet. Chem. Rev., Sect. A* **1972**, *7*, 295. (b) Seyferth, D.; Mammarella, R. E. *J. Organomet. Chem.* **1978**, *156*, 279.

(7) Adcock, W.; Aldous, G. L. *Tetrahedron Lett.* **1978**, 3387 and references therein.

(8) Bank, S.; Sturgess, J. S.; Heyer, D.; Bushweller, C. H. *J. Am. Chem. Soc.* **1980**, *102*, 3982.

(9) Jaouen, G.; Top, S.; McGlinchey, M. J. *J. Organomet. Chem.* **1980**, *195*, C5.

(10) (a) Top, S.; Jaouen, G.; McGlinchey, M. J. *J. Chem. Soc., Chem. Commun.* **1980**, 1110. (b) Top, S.; Jaouen, G. *J. Organomet. Chem.* **1980**, *197*, 199.

(11) Jaouen, G. *Ann. N.Y. Acad. Sci.* **1977**, *295*, 59.

(12) Jaouen, G. In "Transition Metal Organometallics in Organic Synthesis"; Alper, H., Ed.; Academic Press: New York, 1978; Vol. 2, Chapter 2.

[†]École Nationale Supérieure de Chimie.

[‡]McMaster University.





Figure 1. Section of the 100.6-MHz ^{13}C NMR spectrum of **4**, at various temperatures.

Results and Discussion

The barrier to phenyl rotation in the diphenylmethyl anion **3** has been measured as 11.2 kcal mol $^{-1}$.¹³ We have previously reported⁹ the synthesis of **4**, the bis(chromium(0)tricarbonyl)



complex of **3**, and demonstrated that there is extensive delocalization of the carbanionic charge onto the $\text{Cr}(\text{CO})_3$ group. In contrast to the stabilization of benzylic cations by the $\text{Cr}(\text{CO})_3$ group—a phenomenon¹⁴ that seems to involve direct overlap of a filled chromium 3d orbital with the vacant 2p orbital at the α -carbon as in **5**—the stabilization of the benzylic anion is indirect; indeed, such an overlap between the Cr d orbital and the orbital on the α -carbon would be destabilizing since both orbitals are filled. Taking a simple valence bond viewpoint, one could visualize a resonance structure wherein the $\text{Cr}(\text{CO})_3$ is bonded to the ring in a pentahapto manner and there is essentially a double bond between the C_1 phenyl carbon and the benzylic carbon. Such a picture gains some support from X-ray crystallographic data on (fluorene) $\text{Fe}(\text{C}_5\text{H}_5)$.¹⁵ In any event, the occurrence of charge delocalization into the metal carbonyls is unquestionable. The decrease in metal–carbonyl stretching frequencies and deshielding of the carbonyl ^{13}C NMR peaks in **4** relative to the diphenylmethane precursor are comparable, but of opposite sign, to those observed for the corresponding cation.^{9,14a} Interestingly, a recent X-ray crystallographic study of (*N,N*-diethylaniline)tricarbonylchromium(0) revealed a planar geometry for the exocyclic diethylamino group. Furthermore, the N–C(arene) bond was short, supporting the picture of delocalization of the nitrogen lone pair onto the arene and thence to the $\text{Cr}(\text{CO})_3$ group.¹⁶ Now, since the noncomplexed anions **1** and **3** are postulated to be planar,⁸ the enhanced charge delocalization onto the ring and thence to




Figure 2. Section of the 100.6-MHz ^{13}C NMR spectrum of **7**, at -40°C .

the $\text{Cr}(\text{CO})_3$ groups, together with the bulk of the π -complexed moieties, would be expected to considerably hinder phenyl group rotation; we thus expected an increased ΔG^\ddagger relative to the 11.2 kcal mol $^{-1}$ barrier previously reported¹³ for **3**.

The ^{13}C NMR spectra of **4** in THF at various temperatures are shown in Figure 1. The particular features to focus on are the resonances at δ 82 and 99, which are assigned to the ortho (2, 6) and meta (3, 5) carbons, respectively.¹⁷ We see that it is necessary to cool the anion **4** to -110°C at 9.4 T (100.614 MHz for ^{13}C) to obtain the limiting spectrum; these data yield a barrier ΔG^\ddagger_{178} of 8.1 ± 0.5 kcal mol $^{-1}$. It is difficult to rationalize a diminution of the barrier by such a large amount as 3 kcal mol $^{-1}$ purely on the basis of an electronic effect; indeed it would suggest that the $\text{Cr}(\text{CO})_3$ groups destabilize the anion—an interpretation that would contradict all previous evidence on these systems! A more plausible explanation requires a major portion of the barrier in the noncomplexed anions to be steric in origin.¹⁸ Clearly, to maximize p-orbital overlap and minimize the steric interactions between the bulky $\text{Cr}(\text{CO})_3$ groups, the anion should be planar, but such considerations suggest that the assumption of planarity for the noncomplexed anion **3** is invalid. Clearly, in the absence of X-ray crystallographic data on the anion **3** and its bis(chromium(0)tricarbonyl) analogue **4**, it is difficult to separate the steric and electronic components of the barrier to rotation. Furthermore, at some point one reaches a limit to the ability of the α -carbon atom to delocalize its charge, and it is conceivable that the presence of two $\text{Cr}(\text{CO})_3$ units each competing for back-donated electron density actually has a bond-weakening effect on the other. Such a trans influence is frequently encountered with strong π -acid ligands such as CO or PF_3 .¹⁹ Nevertheless, the results of Bank et al.⁸ unequivocally demonstrated the preference for electronic delocalization into a trimethylsilyl-substituted aryl ring and so, to make a meaningful comparison, the bis(chromium(0)tricarbonyl) complex of the 4-(trimethylsilyl)diphenylmethyl anion, viz. **6**, was synthesized. In this case, as for **4**, it was necessary to use a very high-field spectrometer; at -120°C the silylated aryl ring resonances are separated while the phenyl ring rotation is still rapid on the NMR time scale. The barrier to rotation is 8.5 ± 0.5 kcal mol $^{-1}$ and so, while acknowledging the limits of experimental error, the introduction of a 4-trimethylsilyl group stabilizes the negative charge but only to a minimal extent. Apparently, the Me_3Si group exerts only a second-order effect and merely perturbs the much stronger influence of the $\text{Cr}(\text{CO})_3$ group.

To test this hypothesis, the anion of (4'-(trimethylsilyl)diphenylmethane)tricarbonylchromium(0), **7**, was prepared to allow a direct comparison between a *p*- Me_3Si group and a π -complexed

(13) Bank, S.; Sturges, J.; Bushweller, C. H. In "Stereodynamics of Molecular Systems"; Sarma, R. H., Ed.; Pergamon Press: New York, 1979; p 147.

(14) (a) Seyferth, D.; Merola, J. S.; Eschbach, C. S. *J. Am. Chem. Soc.* **1978**, *100*, 4124. (b) Acampora, M.; Ceccon, A.; Dal Farra, M.; Giacometti, G.; Rigatti, G. *J. Chem. Soc., Perkin Trans. 2* **1977**, 483.


(15) Johnson, J. W.; Treichel, P. M. *J. Am. Chem. Soc.* **1977**, *99*, 1427.

(16) Saillard, J. Y.; Grandjean, D.; LeMaux, P.; Jaouen, G. *Nouv. J. Chim.* **1981**, *5*, 153.

(17) In a preliminary communication (ref 9) these assignments were reversed. However, in the present case the data were remeasured on a 400-MHz spectrometer and the assignments confirmed by single-frequency off-resonance decoupling.


(18) While it is true that the data in ref 8 are for the lithium salt and the present results are for the potassium salt, this does not invalidate the conclusions. Indeed, comparisons between analogous Li and K salts show that aryl rotation in the lithium salts is actually faster: Staley, S. W. In "Pericyclic Reactions"; Marchand, A. P., Lehr, R. E., Eds.; Academic Press: New York, 1977; Vol. 1, p 202.

(19) Typically, in $\text{W}(\text{CO})_6$ it is relatively easy to displace three facially disposed carbonyl groups since each is trans to another carbonyl. However, further replacement of carbonyls is much more difficult: King, R. B. "Transition-Metal Organometallic Chemistry"; Academic Press: New York, 1969; pp 73–75.



Cr(CO)₃ substituent. The results were a striking confirmation of the previous observation. A typical spectrum appears as Figure 2 but, in fact, the spectrum scarcely changes over the temperature range -100 °C to room temperature, above which thermal decomposition is observed. In an attempt to observe coalescence of the peaks due to the ortho and meta carbons of the chromium-complexed ring, the spectra were also obtained on 250- and 90-MHz spectrometers (at 62.86 and 22.62 MHz, respectively), but the results were unchanged. As the 4-(trimethylsilyl)phenyl rotation cannot be made slow on the NMR time scale, it is not possible to deduce a rotational barrier but, assuming similar chemical shift differences to those seen in the other anions studied, Δ*G*[‡] is probably somewhat less than 8 kcal mol⁻¹. In contrast, the -C₆H₅Cr(CO)₃ rotation has a barrier of at least 15 kcal mol⁻¹ and may be much higher. Not surprisingly, the same behavior is observed for **8** in which both the Me₃Si and π-Cr(CO)₃ moieties are in the same ring.

The facility with which the anion **4** could be prepared prompted us to treat this ion with chlorotrimethylsilane and then deprotonate a second time to give the anion **9** in which the carbanionic center



was directly bonded to the silicon. This particular anion is thermally stable at room temperature for several hours—even in air! Furthermore, the α-Me₃Si group in **9** provides much more severe competition to the π-complexed rings in its capacity to tolerate negative charge. Indeed, the ¹³C NMR spectrum of **9** shows no separation of the ortho-carbon resonances even at -110 °C. It is thus clear that very extensive delocalization of the carbanionic charge onto silicon has occurred. This is also reflected in the large shielding of the ²⁹Si NMR signal in **9** relative to the other ²⁹Si shifts that appear in Table I.

Another important factor here is the steric crowding around the α-carbon when it is four-coordinate. Thus, the reaction of the anion **4** with Me₃SiCl is a surprisingly slow process, and the ¹³C NMR spectrum of the product, viz. (C₆H₅Cr(CO)₃)₂CHSiMe₃, shows five distinct proton-bearing aromatic carbons demonstrating that aryl rotation is severely hindered even at elevated temperatures. Thus the relief of steric strain favors the formation of the anion **9** in which aryl rotation proceeds without hindrance.

Conclusions

DNMR studies on a series of diphenylmethyl anions reveal that complexation by Cr(CO)₃ groups lowers the barriers to phenyl rotation relative to those found for the noncomplexed anions. It is proposed that the free anions are pyramidal but that complexation renders them planar. Delocalization of the charge on the α-carbon is much better accommodated on the Cr(CO)₃ moiety than on the *p*-Me₃Si group. However, attachment of the Me₃Si grouping directly to the α-carbon yielded an air- and temperature-stable anion in which very extensive delocalization onto silicon has occurred.

Experimental Section

All manipulations involving the organometallic anions were performed under an atmosphere of nitrogen. Solvents were dried over sodium and freshly distilled just prior to use. ¹³C NMR spectra were obtained on a Bruker WM 400 spectrometer operating at 100.614 MHz; ²⁹Si NMR

Table I. ¹³C, ²⁹Si NMR Spectra (ppm) and ν_{CO} Values for Complexes **4** through **8**

	C ₁	C _{2,6}	C ₄	C _α	C ₁	C _{2,6}	C _{3,5}	C ₄	C _{3,5}	C _{3,5}	C ₄	C ₁	C _{2,6}	C _{3,5}	C ₄	CO	CH ₃	²⁹ Si	ν _{CO} , cm ⁻¹	NMR solvent
	112.1 ^a	95.1	93.4	38.6	112.1	95.1	94.9	93.4	94.9	93.4	93.4	112.1	95.1	94.9	93.4	233.8			1975, 1901	THF
	134.2 ^b	81.2 ^c	78.4	76.4	134.2	81.2 ^c	98.7 ^d	78.4	98.7 ^d	78.4	78.4	134.2	81.2 ^c	98.7 ^d	78.4	237.4			1941, 1858	THF
	113.6 ^a	94.7 ^e	93.1	39.8	111.6	92.7	94.5 ^e	93.1	94.5 ^e	99.4	99.4	111.6	92.7	101.1	99.4	234.1, 233.8	-1.9	0.4	1978, 1970, 1900	acetone
	136.4 ^b	82.3	78.3	77.0	133.5	81.1 ^f	98.7	78.3	103.4 ^g	79.4	79.4	133.5	81.1 ^f	103.4 ^g	79.4	237.7, 237.3	-1.5	-2.7	1949, 1936, 1860	THF
	112.9 ^a	93.9 ^e	90.0	41.3	139.9	129.0	93.4 ^e	90.0	99.6/99.0	139.6	139.6	139.9	129.0	134.6	139.6	234.1	-0.6	-4.6	1970, 1896	benzene
	136.0 ^b	86.4/77.0	72.8	71.4	144.4	123.5	99.6/99.0	72.8	132.3	126.4	126.4	144.4	123.5	132.3	126.4	239.4	-1.5	-7.4	1921, 1822	THF
	114.9 ^a	92.1	97.8	41.7	139.6	129.8	100.4	97.8	129.8	127.8	127.8	139.6	129.8	129.8	127.8	234.5	-0.7	0.1	1970, 1896	benzene
	135.9 ^b	87.6/78.8	76.6	73.4	143.1	124.3	103.3/102.5	76.6	127.1	118.4	118.4	143.1	124.3	127.1	118.4	239.9	-1.1	-3.1	1921, 1822	THF
	113.5 ^a	94.6/90.3	89.3	41.1	113.5	94.6/90.3	96.4/95.5	89.3	96.4/95.5	89.3	89.3	113.5	94.6/90.3	96.4/95.5	89.3	233.2	-0.4	-4.0	1955, 1895	chloroform
	135.8 ^b	86.1	80.5	98.9	135.8	86.1	97.5	80.5	97.5	80.5	80.5	135.8	86.1	97.5	80.5	237.2	+1.7	-10.4	1945, 1840	THF

^a Data for the diphenylmethane complexes. ^b Data for the diphenylmethyl anion complexes (in italics). ^c 82.5/79.6 at -110 °C. ^d 99.4/99.2 at -110 °C. ^e Assignments may be interchanged.

^f 83.5/81.3 at -110 °C. ^g 103.6/103.3 at -120 °C.

spectra were obtained on a Bruker WM 250 spectrometer operating at 49.964 MHz; shifts are reported relative to tetramethylsilane and positive shifts are to high frequency.

Synthesis of 4-(Trimethylsilyl)diphenylmethane. 4-Chlorobenzophenone (22 g, 0.1 mol) dissolved in anhydrous ether was added to LiAlH_4 (6.6 g, 0.18 mol) and AlCl_3 (48 g, 0.36 mol) in 100 mL of ether, and the mixture was heated under reflux for 1.5 h according to the method of Blackwell and Hickinbottom.²⁰ Excess of the reagent was destroyed by adding ethyl acetate, and the mixture was poured into 20% aqueous H_2SO_4 . Ether extraction, solvent evaporation, and vacuum distillation (160 °C/11 torr) gave 4-chlorodiphenylmethane (18 g, 0.089 mol; 89%). This latter product (8 g, 0.04 mol) was stirred with magnesium (3.6 g, 0.15 mol) in THF at 50 °C over a 12-h period. Chlorotrimethylsilane (15 g, 0.138 mol) was added and the mixture heated at 90 °C for 48 h. Addition to water, ether extraction, and removal of solvent in vacuo gave 4-(trimethylsilyl)diphenylmethane (9.5 g) as a pale yellow, slightly viscous liquid, which was not further purified. This product gave a ^{29}Si NMR peak at -5.2 ppm.

Syntheses of the (Diphenylmethane)tricarboxylchromium(0) Complexes. (Diphenylmethane)bis(tricarboxylchromium(0)) was prepared as described previously.^{21,22} 4-(Trimethylsilyl)diphenylmethane (3 g, 12.5 mmol), $\text{Cr}(\text{CO})_6$ (2.2 g, 10 mmol), dibutyl ether (100 mL), and THF (10 mL) were heated under reflux in a nitrogen atmosphere for 24 h. After filtration and evaporation of the solvent, the residual oil was chromatographed on a silica gel column by using ether/petroleum ether (1:10) as eluent. The first product was recrystallized from ether/petroleum ether to give yellow crystals of 4-(trimethylsilyl)diphenylmethane)tricarboxylchromium(0) (700 mg, 1.86 mmol; 19%), mp 80 °C. Anal. Calcd for $\text{C}_{19}\text{H}_{20}\text{CrO}_3\text{Si}$: C, 60.6; H, 5.3. Found: C, 60.9; H, 5.4.

The second product eluted was recrystallized from ether/petroleum ether to give yellow crystals of (4'-(trimethylsilyl)diphenylmethane)tricarboxylchromium(0) (150 mg, 0.4 mmol; 4%), mp 123 °C. Anal. Calcd for $\text{C}_{19}\text{H}_{20}\text{CrO}_3\text{Si}$: C, 60.6; H, 5.3. Found: C, 60.4; H, 5.4.

4-(Trimethylsilyl)diphenylmethane (1.7 g, 7.1 mmol), $\text{Cr}(\text{CO})_6$ (4.4 g, 20 mmol), dibutyl ether (100 mL), and THF (10 mL) were heated under reflux in a nitrogen atmosphere for 48 h. After filtration the solution was cooled in a refrigerator overnight. Yellow crystals precip-

itated and were recrystallized from ether/petroleum ether to give 4-(trimethylsilyl)diphenylmethane)bis(tricarboxylchromium(0)) (2 g, 3.9 mmol; 55%), mp 144 °C. Anal. Calcd for $\text{C}_{22}\text{H}_{20}\text{Cr}_2\text{O}_6\text{Si}$: C, 51.6; H, 3.9. Found: C, 51.5; H, 4.1.

Syntheses of (α -(Trimethylsilyl)diphenylmethane)bis(tricarboxylchromium(0)). (Diphenylmethane)bis(tricarboxylchromium(0)) (220 mg, 0.5 mmol) was dissolved in THF (4 mL) and was treated with potassium *tert*-butoxide (280 mg, 2.5 mmol) at 0 °C. After 5 min the solution was filtered and treated with anhydrous hexane to precipitate the anion 4. After decantation and removal of liquid by using a syringe, the anion was redissolved in THF (6 mL). Immediately, chlorotrimethylsilane (1 mL) was added and the initially orange solution gradually became yellow. After 30 min the reaction was quenched with water. After ether extraction and removal of solvent in vacuo, the crude residue was chromatographed on a silica gel column and was eluted with ether/petroleum ether (3:2) to give yellow crystals of (α -(trimethylsilyl)diphenylmethane)bis(tricarboxylchromium(0)) (30 mg, 0.06 mmol), mp 168 °C. Anal. Calcd for $\text{C}_{22}\text{H}_{20}\text{Cr}_2\text{O}_6\text{Si}$: C, 51.6; H, 3.9. Found: C, 51.8; H, 3.8. Some starting material, viz. $(\text{PhCH}_2\text{Ph})[\text{Cr}(\text{CO})_3]_2$, was also recovered (80 mg); the global yield of the α -silylated complex was thus 19%.

Generation of the Anions for the NMR Study. Freshly distilled THF was added to a mixture of the appropriate complex and a 5-fold quantity of potassium *tert*-butoxide at 0 °C. (For the two monocomplexed compounds this was done at -30 °C since their anions are thermally sensitive.) After stirring for 5 min the anion was filtered directly into the NMR tube and sealed in vacuo. The anions were kept frozen in liquid nitrogen until required for the NMR studies. The data pertaining to the anions listed in Table I were obtained at -40 °C except where stated.

Acknowledgment. We thank Dr. R. E. Lenkinski, Southwest Ontario Regional NMR Center, Guelph, Ontario, for recording the high-field ^{13}C spectra. Financial support from the CNRS (France) and NSERC (Canada) is greatly appreciated. G.J. and S.T. thank the France-Canada Scientific Exchange Program for Visiting Fellowships.

Registry No. 4, 32823-67-9; 4⁻, 75983-29-8; 6, 86969-84-8; 6⁻, 86969-78-0; 7, 86969-82-6; 7⁻, 86969-79-1; 8, 86969-83-7; 8⁻, 86969-80-4; 9, 86969-85-9; 9⁻, 86969-81-5; 4-chlorobenzophenone, 134-85-0; 4-chlorodiphenylmethane, 831-81-2; 4-(trimethylsilyl)diphenylmethane, 17964-29-3.

(20) Blackwell, J.; Hickinbottom, W. J. *J. Chem. Soc.* **1961**, 1405.

(21) Mahaffey, C. A. L.; Pauson, P. L. *Inorg. Synth.* **1975**, *19*, 154.

(22) Top, S.; Jaouen, G. *J. Organomet. Chem.* **1979**, *182*, 381.

Sapphyrins: Novel Aromatic Pentapyrrolic Macrocycles

Victor J. Bauer,[†] Derrick L. J. Clive,[‡] David Dolphin,^{*§} John B. Paine III,[§] Francis L. Harris,^{*||} Michael M. King,[⊥] John Loder,[⊙] Shen-Wei Chien Wang,[▽] and Robert Burns Woodward[&]

Contribution from the Departments of Chemistry, Harvard University, Cambridge, Massachusetts 02138, and University of British Columbia, Vancouver, British Columbia, Canada V6T 1Y6. Received August 10, 1982

Abstract: Sapphyrins are pentapyrrolic macrocycles containing one direct link and four bridging methine groups between the five pyrrolic subunits. The syntheses of decamethylsapphyrin, other peripherally alkylated derivatives, and metal complexes are described. The physical and chemical properties of sapphyrins show them to be aromatic like porphyrins and corroles. This aromaticity is reflected in the large shielding of the methine and deshielding of the NH protons in the NMR spectra and by their optical spectra, which exhibit Soret and visible bands similar to porphyrins but all bathochromically shifted.

The chemistry described here was performed under the guidance and encouragement of the late R. B. Woodward and began two decades ago following a report by Johnson and Price¹ claiming the synthesis of metalcorroles (1). These complexes were

supposedly prepared by treating the palladium complex of 5,5'-bi(5'-bromo-3,3'-diethyl-4,4'-dimethylpyrromethene) (2), originally prepared by Fischer and Stachel,² with formaldehyde and hydrochloric acid in refluxing ethanol. However, it was shown that this reaction gave the metallooxocorroles (3) instead.³ Since the original claim for a corrole synthesis had not been fulfilled, Woodward and Bauer directed their efforts toward such a synthesis

[†]Hoechst-Roussel Pharmaceuticals Inc., Somerville, NJ 08876.

[‡]University of Alberta, Edmonton, Alberta, Canada T6G 2G2.

[§]University of British Columbia, Vancouver, British Columbia, Canada V6T 1Y6.

[⊥]California State University, Northridge, CA 91330.

[⊙]The George Washington University, Washington, DC 20052.

[⊙]CSIRO, Melbourne 3001, Australia.

[▽]The Henkel Co., Hawthorne, CA 90250.

[&]Deceased.

(1) Johnson, A. W.; Price, R. *J. Chem. Soc.* **1960**, 1649.

(2) Fischer, H.; Stachel, A. *Hoppe-Seyler's Z. Physiol. Chem.* **1939**, *258*, 121.

(3) Johnson, A. W.; Kay, I. T. *Proc. Chem. Soc., London* **1961**, 168.