


In the first of these (expt 1), 12 g of 48-h old mycelial pellets of the averufin-accumulating mutant ATCC 24551 were suspended in a low-sugar replacement medium¹⁸ (100 mL) in each of 12 500-mL Erlenmeyer flasks. Labeled acid (12 mg) was administered to each flask under sterile conditions, and they were shaken at 28 °C for 24 h. The averufin produced was isolated by extraction of the filtered pellets and purified by chromatography on silica gel. In the second experiment (expt 2), a 100-mL 48-h-old standing culture of the same mutant grown on a low-salts medium¹⁹ was treated with [1-¹³C]hexanoic acid (25 mg) and again at 72 and 96 h. At 120 h the fermentation was terminated and the averufin was isolated.

When compared to a [¹³C]H NMR spectrum of averufin at natural abundance (normalization to C-5, an unlabeled center derived from C-2 of acetate), the corresponding spectra of the ¹³C-enriched averufin from both expt 1 and 2 showed the resonance for C-1' at δ 66.2 to be enhanced more than 3 times the natural abundance (solid circle in 6). Some degradation of the labeled



hexanoate to [1-¹³C]acetate and secondary incorporation (open circles in 6) were observed, being about 0.5%/site in expt 1 and 1%/site in expt 2.

The implications of these findings were further examined by testing the incorporation of *equimolar* amounts of [1-¹³C]acetate, [1-¹³C]butyrate, [1-¹³C]-5-oxohexanoate,²⁰ and [1-¹³C]-3-oxooctanoate²¹ under the suspended-cell conditions above. For each of these substrates the pattern of ¹³C enrichment in averufin, while varying somewhat in overall magnitude from experiment to experiment, was the same as that for the uniform incorporation of [1-¹³C]acetate (about 0.5%/site for the latter). Therefore, all of these experiments to a first approximation label the acetate pool to roughly the same extent as reflected in the incorporation of C-1 throughout averufin (6) but [1-¹³C]hexanoate alone shows a significant level of intact incorporation.

These data suggest an intact hexanoate starter unit arising from a separate synthetase or from β -oxidation. Alternatively, a single polyketide synthetase may produce the initially reduced segment which is able to exchange with free hexanoyl CoA at the C₆ stage. This process would have to be quite efficient given the 3% specific incorporation of label from hexanoate compared with about 0.5% from acetate. Butyrate and 3-oxooctanoate, which might be expected to show analogous exchange, fail detectably to do so or at best with significantly reduced efficiency. Lastly, it is possible that acylation of a preformed anthraquinone may take place. 1,3,6,8-Tetrahydroxyanthraquinone has been isolated from *A. versicolor*,²² and while structurally symmetrical, its labeling pattern from [1,2-¹³C₂]acetate is not. Such a process would require acylation to occur in the enzyme-bound state. Parallel formation

of averufin from 5-oxohexanoate is not supported experimentally.

Of these interpretations we prefer, but cannot strictly prove, the first for its simplicity and direct relation to precedents noted at the outset. It is of interest at this point to recall the observation of Holker and Mulheirn²³ that the bisfuran carbons of sterigmatocystin, from *A. versicolor*, bore about 10% lower specific activity from [1-¹⁴C]acetate than did nuclear carbons. In contrast Büchi's extensive degradations of aflatoxin B₁ (5) from the corresponding incorporation experiment in *A. flavus* revealed no differential labeling.¹⁴ In the event, it is now clear that the seemingly elaborate formation of averufin (4) is necessary given the constraint of an hexanoate primer (Scheme I) and equally necessary one might assume to ultimately construct the singular bisfuran that characterizes this family of mycotoxins.^{24,25}

Acknowledgment. Professor J. M. Schwab (Catholic University of America) is thanked for recording the [1-¹³C]hexanoate incorporation spectra. We are pleased to note helpful conversations with Dr. U. Weiss (NIH) and Professor J. C. Vederas (University of Alberta). Financial support of the National Institutes of Health (ES 01670) and the Alfred P. Sloan Foundation is gratefully acknowledged.

Registry No. 4, 14016-29-6; hexanoic acid, 142-62-1.

(23) Holker, J. S. E.; Mulheirn, L. J. *J. Chem. Soc., Chem. Commun.* **1968**, 1576-1577.

(24) Townsend, C. A.; Christensen, S. B. *Tetrahedron* **1983**, *39*, 3357-3582.

(25) See for a perhaps fortuitous comparison the effects of, inter alia, added unlabeled butyrate, hexanoate, and octanoate on natural fatty acid production in *Bacillus acidocaldaria*.¹⁰

Total Synthesis of (±)-Poitediol and (±)-4-Epipoitediol


Robert C. Gadwood,* René M. Lett, and Jane E. Wissinger

*Department of Chemistry, Northwestern University
Evanston, Illinois 60201*

Received January 9, 1984

Revised Manuscript Received April 23, 1984

Poitediol (1) is an unusual sesquiterpene diol isolated from the red seaweed *Laurencia Poitei*.¹ It is one of the less complex



members of a growing family of cyclooctane-containing natural products, many of which have interesting biological activities. We wish to report the total synthesis of racemic poitediol and 4-epipoitediol.²

Our earlier work³ had shown that the anionic oxy-Cope rearrangement of dialkenyl cyclobutoxides was an efficient method for the synthesis of substituted cyclooctanones, and our strategy for the synthesis of poitediol is based upon this approach. In order to construct the bicyclo[6.3.0]undecane system of poitediol, we required the bicyclo[3.2.0]heptanone 9 whose synthesis is outlined in Scheme I.⁴

(1) Fenical, W.; Schulte, G. R.; Finer, J.; Clardy, J. *J. Org. Chem.* **1978**, *43*, 3628.

(2) Other total and partial synthesis of cyclooctanoid natural products: (a) (pleuromutilin) Gibbons, E. G. *J. Am. Chem. Soc.* **1982**, *104*, 1767. (b) (epiprecapnelladiene) Birch, A. M.; Pattenden, G. *J. Chem. Soc., Chem. Commun.* **1980**, 1195; *J. Chem. Soc., Perkin Trans. 1*, **1983**, 1913. (c) (ophiobolins) Paquette, L. A.; Andres, D. R.; Springer, J. P. *J. Org. Chem.* **1983**, *48*, 1147 and references cited therein.

(3) Gadwood, R. C.; Lett, R. M. *J. Org. Chem.* **1982**, *47*, 2268.


(18) Lin, M. T.; Hsieh, D. P. H.; Yao, R. C.; Donkersloot, J. A. *Biochemistry* **1973**, *12*, 5167-5171.

(19) Reddy, T. V.; Viswanathan, L.; Venkatasubramanian, T. A. *Appl. Microbiol.* **1971**, *22*, 393-396.

(20) Cf.: Fitzell, D. L.; Hsieh, D. P. H.; Reece, C. A.; Seiber, J. N. *J. Labelled Compd.* **1975**, *11*, 135-139 and references cited.

(21) Prepared by the reaction of *tert*-butyldimethylsilyl [1-¹³C]acetate in the presence of lithium diisopropylamide (2.4 equiv) in tetrahydrofuran at -78 °C with hexanoyl chloride. Loquin, R. *Bull. Soc. Chem. Fr.* **1904**, *108*, 595-599.

(22) Berger, Y. *Phytochemistry* **1980**, *19*, 2779-2780.

Scheme I^a

^a (a) NaBH₄, MeOH, 0 °C; (b) Ac₂O, catalytic H₂SO₄, Et₂O; (c) KO-*t*-Bu, THF; (d) DIBAH, toluene, -78 °C; (e) MeI, NaH, THF, 0 °C; (f) (1) O₃, CH₂Cl₂, -78 °C (2) Me₂S, reflux; (g) NaOH, MeOH, reflux; (h) DIBAH, Et₂O, -100 °C; (i) Et₂Zn, CH₂I₂, toluene, 60 °C; (j) PCC, CH₂Cl₂; (k) CH₂=CHMgBr, THF; (l) BF₃·Et₂O, Et₂O.

Reduction of 5-methyl-2-carbomethoxycyclopentanone⁵ followed by acetylation and elimination afforded cyclopentene 4. Further reduction and methylation of the resulting alcohol produced 5. Ozonolysis (Me₂S workup) and cyclization of the resulting keto aldehyde led to cyclohexenone 6. Stereoselective reduction of 6 with DIBAH in Et₂O at -100 °C⁶ gave a diastereomeric mixture, which was cyclopropanated⁷ to afford a separable mixture of norcaranol 7a and 7b (8:1 ratio).⁸

Oxidation of 7a and addition of vinylmagnesium bromide gave norcaranol 8. Attempted rearrangement of 8 using aqueous HBF₄¹⁰ led to formation of two products, apparently due to competitive migration of both cyclopropyl carbon-carbon bonds. However, treatment with 1 equiv of BF₃·Et₂O afforded bicyclo[3.2.0]heptanone 9 in essentially quantitative yield.

Ring expansion of 9 via sequential addition of 1-lithiovinyl phenyl sulfide (or selenide) and anionic oxy-Cope rearrangement was successful, as was the subsequent oxidative elimination. However, this sequence proceeded in disappointingly low overall

(4) All new compounds gave satisfactory NMR, IR, and mass spectral or analytical data.

(5) Sisido, K.; Utimoto, K.; Isida, T. *J. Org. Chem.* **1964**, *29*, 2781; Meyer, W. L.; Lobo, A. P.; Marquis, E. G. *Ibid.* **1965**, *30*, 183.

(6) Heathcock, C. H.; Tice, C. M.; Gemroth, T. C. *J. Am. Chem. Soc.* **1982**, *104*, 6081.

(7) Furukawa, J.; Kawabata, N.; Nishimura, J. *Tetrahedron Lett.* **1966**, 3353; *Tetrahedron* **1968**, *24*, 53.


(8) The relative stereochemistries of the major and minor norcaranol were determined by acid-catalyzed rearrangement to the bicyclo[3.2.0]heptanones



i and ii, respectively, which gave the ¹³C NMR data shown below. Both C7 and C8 of the ketone ii from the minor norcaranol are clearly shifted upfield due to the steric shielding effect of the endo methyl group.⁹

(9) Hudlicky, T.; Koszyk, F. J.; Kutchan, T. M.; Sheth, J. P. *J. Org. Chem.* **1980**, *45*, 5020. Whitesell, J. K.; Matthews, R. S. *Ibid.* **1977**, *42*, 3878. Christl, M.; Reich, H. J.; Roberts, J. D. *J. Am. Chem. Soc.* **1971**, *93*, 3463.

(10) Wenkert, E.; Arrhenius, T. S. *J. Am. Chem. Soc.* **1983**, *105*, 2030. Cohen, T.; Matz, J. R. *Tetrahedron Lett.* **1981**, *22*, 2455. Wenkert, E.; Golob, N. F.; Hatch, R. P.; Wenkert, D. *Helv. Chim. Acta* **1977**, *60*, 1.

Scheme II^a

^a (a) LiC≡CH, THF, -30 °C, 5 min; (b) hexane, 50 °C, 4 h; (c) MeLi, Et₂O, -78 °C; (d) PCC, CH₂Cl₂; (e) LiMe₂Cu, Me₂S, Et₂O; (f) DIBAH, Et₂O, -78 °C; (g) (1) KH, THF, (2) PhCH₂Br; (h) MCPBA, CHCl₃; (i) LiEt₃BH, THF; (j) SEMCl, *i*-Pr₂NEt, THF, 50 °C; (k) Na, NH₃; (l) (1) oxalyl chloride, Me₂SO, -78 °C; (2) *i*-Pr₂NEt; (m) (1) LDA, THF; (2) CH₂O; (3) MsCl, *i*-Pr₂NEt; (n) DIBAH, hexane, -78 °C; (o) 0.1 M HCl in MeOH.

yield. In contrast, as shown in Scheme II,⁴ the more direct procedure of lithium acetylide addition¹¹ followed by neutral oxy-Cope rearrangement^{12,13} was more successful, leading to 10 in overall yields of 50–60%. The oxy-Cope rearrangement of a 1-alkynyl-2-alkenylcyclobutanol is unprecedented and should open up new avenues for the synthesis of functionalized cyclooctanes.

Geminal dimethylation and carbonyl transposition of 10 were accomplished by addition of methylolithium, oxidative rearrangement with PCC,¹⁴ and reaction with lithium dimethyl cuprate¹⁵ to afford enone 11. Reduction and benzylation followed by epoxidation afforded an inseparable 3:2 mixture of epoxides 12a and 12b. Treatment of this mixture with LiEt₃BH produced a mixture of alcohols, from which the desired diastereomer could be easily separated. Protection of this alcohol as a SEM ether¹⁶ followed by debenylation and Swern oxidation¹⁷ produced ketone 13. Introduction of the α-methylene group to produce 14 was accomplished by enolate formation with LDA, quenching with formaldehyde, and elimination of the intermediate alcohol. Reduction with DIBAH produced a separable 1:1 mixture of alcohols, which were deprotected to afford pitoediol and 4-epipitoediol. The synthetic pitoediol gave spectral data identical with an authentic sample.¹⁸

Acknowledgment is made to the donors of the Petroleum Research Fund, administered by the American Chemical Society, and to NSF for support of this work. We are also grateful to NSF for funds used to purchase a 270-MHz NMR spectrometer.

(11) Midland, M. M. *J. Org. Chem.* **1975**, *40*, 2250.

(12) The anionic oxy-Cope rearrangement of the intermediate adduct was complicated by formation of several unidentified byproducts in addition to the desired 13.

(13) For a review of known Cope rearrangements involving alkynes, see: Viola, A.; Collins, J. J.; Filipp, N. *Tetrahedron* **1981**, *37*, 3765.

(14) Dauben, W. G.; Michno, D. M. *J. Org. Chem.* **1977**, *42*, 682.

(15) House, H. O.; Chu, C. V.; Wilkins, J. M.; Umen, M. J. *J. Org. Chem.* **1975**, *40*, 1460.

(16) Lipshutz, B. H.; Pegram, J. J. *Tetrahedron Lett.* **1980**, 3343.

(17) Huang, S. L.; Omura, K.; Swern, D. *Synthesis* **1978**, 297; Omura, K.; Swern, D. *Tetrahedron* **1978**, *34*, 1651.

(18) An authentic sample of pitoediol was graciously provided by Ms. Valerie Paul from Professor William Fenical's laboratory.