


# Rate-Limiting Deprotonation in Tetrahedral Intermediate Breakdown

Robert A. McClelland

Contribution from the Department of Chemistry and Scarborough Campus, University of Toronto, West Hill, Ontario M1C 1A4, Canada. Received April 6, 1984.  
Revised Manuscript Received August 14, 1984

**Abstract:** A kinetic study in acetate and formate buffers is reported of the breakdown to  $\beta$ -hydroxyethyl benzoate of the hemiothoester-type tetrahedral intermediate 2-phenyl-2-hydroxy-1,3-dioxolane. This species is generated as a transient intermediate in acid solutions in the hydrolysis of the amide acetals 2-phenyl-2-(dimethylamino)-1,3-dioxolane and 2-phenyl-2-(1-imidazolyl)-1,3-dioxolane. Plots of observed rate constant vs. buffer concentration exhibit pronounced curvature in dilute buffers suggestive of a change in rate-limiting step with changing buffer concentration. A mechanism is proposed involving deprotonation of the hydroxy group of the hemiothoester to produce a highly reactive anion. In the absence of buffer and in very dilute buffers the deprotonation step is rate limiting, the anionic hemiothoester going on to product at a rate faster than it is reprotonated. In more concentrated buffers the deprotonation step becomes reversible because of the availability of the acid component of the buffer for the reprotonation. A kinetic analysis is presented, providing a rate constant of approximately  $5 \times 10^7 \text{ s}^{-1}$  for breakdown of the anionic hemiothoester. This intermediate is comparable to that which would be encountered during the hydrolysis of a benzoate ester, and some consequences of the large reactivity are discussed. This is suggested to be a contributing factor behind the small amount of carbonyl  $^{18}\text{O}$  exchange observed during the base hydrolysis of esters. It is also pointed out that with leaving groups significantly better than alkoxide, the lifetime may become so short that the tetrahedral intermediate anion could not exist. In this case an acyl transfer reaction proceeding via this anion would have to be regarded as being concerted with the anion as a transition state, not an intermediate.

A number of examples have recently been reported<sup>1,2</sup> of the observation of hemiothoesters  $\text{R}^{\text{C}}(\text{OR})_2\text{OH}$  as transient intermediates in the hydrolysis of some ortho acid derivative. He-



miothoesters are tetrahedral intermediates of [O,O]-acyl transfer reactions, and the generation from the ortho acid derivative has allowed direct kinetic study of their breakdown. This decomposition is invariably general acid-base catalyzed, and a particularly interesting finding has been that of very large rate constants for hydroxide ion catalysis ( $k_{\text{OH}^-}$ ), in many instances numbers approaching and even exceeding the diffusion limit. The commonly accepted mechanism<sup>1-4</sup> for the reaction with hydroxide involves deprotonation of the hydroxyl group of the hemiothoester followed by expulsion of alkoxide ion from the anion. A value for  $k_{\text{OH}^-}$  at the diffusion limit implies the deprotonation step is rate limiting. The situation is clouded, however, by the presence of buffer in the solutions where  $k_{\text{OH}^-}$  is measured, since the buffer acid:base species can participate in the initial proton transfer reaction.<sup>3</sup>

This paper presents a study of the effect of buffers on the decomposition of a cyclic hemiothoester, 2-phenyl-2-hydroxy-1,3-dioxolane (**1**). We have previously reported<sup>5,6</sup> a value of  $k_{\text{OH}^-}$  of  $6 \times 10^{10} \text{ M}^{-1} \text{ s}^{-1}$  for this compound. This is a number perhaps about an order of magnitude too large for a diffusion-limited rate constant. Its determination was based on extrapolation to zero buffer concentration of data obtained in quite concentrated buffers,<sup>5,6</sup> and we suspected that curvature might occur in the buffer dilution plots in dilute buffers. Significant curvature is indeed observed, implicating a change in rate-determining step with changing buffer concentration. This allows a kinetic analysis

to be carried out which furnishes a measure of the lifetime of the anionic tetrahedral intermediate.

## Experimental Section

**Amide Acetals.** 2-Phenyl-2-(dimethylamino)-1,3-dioxolane was from a previous study.<sup>5</sup> 2-Phenyl-2-(1-imidazolyl)-1,3-dioxolane was prepared as follows. 2-Phenyl-1,3-dioxolan-2-ylum tetrafluoroborate<sup>6</sup> (0.05 mol) and the sodium salt of imidazole (0.08–0.10 mol) (prepared from imidazole and sodium hydride in dry tetrahydrofuran) were isolated in their solid forms under anhydrous conditions in a glovebag, and the solids were mixed (no apparent reaction). The mixture was cooled in a dry ice:acetone bath and 20 mL of cooled ( $-30^\circ\text{C}$ ) acetonitrile freshly distilled from  $\text{P}_2\text{O}_5$  added all at once. The mixture was then stirred and allowed to come to room temperature, 20–50 mL of dry ether added, and the mixture filtered (to remove unreacted imidazole salt and  $\text{NaBF}_4$ ). The filtrate was evaporated on a rotary evaporator and the remaining solid purified by dissolving in benzene and precipitation with hexane. The imidazole amide acetal had mp  $85^\circ\text{C}$  and NMR  $\delta$  7.74 (1 H, br s), 7.36 (5 H, s), 7.08 (2 H, br s), 4.24 (4 H, symmetrical m). Anal. Calcd for  $\text{C}_{12}\text{H}_{12}\text{N}_2\text{O}_2$ : C, 66.65; H, 5.59; N, 12.95. Found: C, 66.59; H, 5.71; N, 12.88.

**Kinetics.** These were performed essentially as previously described<sup>5</sup> by using a Durrum Gibson stopped-flow spectrophotometer. The amide acetals were added to an 0.01 M NaOH solution which had been thermostated at  $25^\circ\text{C}$  in an external water bath. This solution was placed in one syringe of the stopped-flow apparatus and mixed with acetate (or formate) buffers containing 0.01 M HCl. In order to ensure that pH was remaining constant upon buffer dilution the pH of all mixtures was immediately recorded with use of solutions mixed on the stopped-flow apparatus. A variation of  $\pm 0.02$  was considered satisfactory; results outside of this range were discarded. Rate constants were evaluated from the absorbance increase at 230 nm. The photomultiplier output of the stopped-flow was digitized to give 118 absorbance points per kinetic run, plus 4 infinity values. Eight-ten kinetic runs were performed for each buffer. Excellent linearity was observed in all plots of  $\ln(A_\infty - A)$  vs. time; observed rate constants were calculated as the slopes of these plots.

## Results

**Hemiothoester Generation.** The approach for generating the hemiothoester with amide acetals as precursors is summarized in eq 2. The key feature of the chemistry is that the protonated forms of these compounds undergo loss of the exocyclic amine or imidazole extremely rapidly. Hydration of the 1,3-dioxolan-2-ylum ion **4** is also very rapid.<sup>6</sup> Thus in acid solutions where the protonated form of the amide acetal is favored in the initial equilibrium, the steps leading to the hemiothoester are faster than its decomposition. This means that the hemiothoester forms in

(1) Capon, B.; Ghosh, A. K.; Grieve, D. M. A. *Acc. Chem. Res.* **1981**, *14*, 306–312.


(2) McClelland, R. A.; Santry, L. J. *Acc. Chem. Res.* **1983**, *16*, 394–399.

(3) Gravitz, N.; Jencks, W. P. *J. Am. Chem. Soc.* **1974**, *96*, 489–499.

(4) Funderburk, L. H.; Aldwin, L.; Jencks, W. P. *J. Am. Chem. Soc.* **1978**, *100*, 5444–5459. McClelland, R. A.; Coc, M. *Ibid.* **1983**, *105*, 2718–2725.

(5) McClelland, R. A.; Ahmad, M. *J. Org. Chem.* **1979**, *44*, 1855–1860.

(6) Ahmad, M.; Bergstrom, R. G.; Cashen, M. J.; Chiang, Y.; Kresge, A. J.; McClelland, R. A.; Powell, M. F. *J. Am. Chem. Soc.* **1979**, *101*, 2669–2677.



nearly quantitative amounts before undergoing further reaction. Moreover, since the kinetics are based upon the appearance of the benzoate ester **5**, they measure directly the rate of the hemiorthoester decomposition process. The detailed analysis has been presented in a previous study<sup>5</sup> for the dimethylamino compound. Investigations on imidazole amide acetals have just recently begun in our laboratories, and the pertinent results obtained with the dioxolane **3** are presented in the next section. What is important to note here is that the dimethylamine and imidazole systems exhibit identical rate behavior in solutions with pH < 5.5. It is in these solutions that hemiorthoester decomposition is proposed to be the slow step in product formation, and the two compounds must, of course, produce the same hemiorthoester.

**Imidazole Amide Acetal Hydrolysis.** Examples of imidazole amide acetals have only recently been prepared by Brown and co-workers,<sup>8</sup> using exchange reactions of ortho esters and imidazoles. This reaction fails with 2-phenyl-2-methoxy-1,3-dioxolane and imidazole. The preparation of **3**, however, was accomplished by combining the 2-phenyl-1,3-dioxolan-2-ylum cation as its borofluorate salt with the imidazole anion as its sodium salt.

Brown has also recently reported a kinetic study of the hydrolysis of some formic acid derived imidazole amide acetals.<sup>9</sup> A single reaction was observed, involving equilibrium protonation of the imidazole followed by loss of this group to produce the dialkoxycarbocation and eventually ester and alcohol products. This is the mechanism outlined in eq 2. Unlike the situation with **3** the kinetic study with the formic acid derivatives was based<sup>9</sup> upon changes in the imidazole chromophore, so that at all pH the rate of the formation of the dialkoxycarbocation is measured. The rate-pH profile is thus quite straightforward, with rate constants at high pH being proportional to H<sup>+</sup> concentration because of the protonation requirement and the rate constants leveling in acid solution when protonation is complete.

The basic chemistry of the hydrolysis of the imidazole amide acetal **3** is the same, with imidazole and the hydroxy ester **5** being the only products at all pH. In base solutions observed rate constants are linear in H<sup>+</sup> concentration with a slope ( $k_H$ ) of  $8 \times 10^9 \text{ M}^{-1} \text{ s}^{-1}$ . This represents rate-limiting formation of the cation **4**, and  $k_H$  can be set equal to  $k_N/K_{XH^+}$  with the latter terms as defined in eq 2. Brown found  $pK_{XH^+} = 5.97$  for imidazole protonation in his study, and a similar value is likely here since the substituent differences are quite remote from the protonation site. Therefore  $k_N$  should be of the order of  $10^4 \text{ s}^{-1}$ . The rate of formation of the cation **4** in acids is equal to  $k_N$ . Rate constants for the hemiorthoester decomposition are, however, considerably smaller,<sup>2,5,6</sup> so that a change in slow step in product formation occurs. The decomposition becomes rate limiting, and the measured rate constants represent directly this decomposition.

**Rate Constants.** Rate constants for this hemiorthoester decomposition have been measured in a series of acetate buffers and one formate buffer. Emphasis was placed on dilute buffer solutions, and particular care was taken to ensure constant pH on buffer dilution. As noted previously the two precursors give identical rate constants in the same solution. The majority of the measurements were made with the imidazole system. This has the advantage of undergoing only the one reaction; the di-




Figure 1. First-order rate constants ( $25^\circ \text{C}$ , ionic strength  $0.1 \text{ s}^{-1}$ ) for the decomposition of the tetrahedral intermediate 2-phenyl-2-hydroxy-1,3-dioxolane in acetate buffers [(A)  $\alpha(\text{HA}) = 0.8$ , pH 3.92; (B)  $\alpha(\text{HA}) = 0.6$ , pH 4.38; (C)  $\alpha(\text{HA}) = 0.4$ , pH 4.72; (D)  $\alpha(\text{HA}) = 0.2$ , pH 5.12] and in formate buffer [(E)  $\alpha(\text{HA}) = 0.2$ , pH 4.09].

methylamino derivative in solutions above pH 4.5 begins to undergo a competing C-O cleavage.<sup>5</sup> First-order rate constants for the hemiorthoester decomposition are listed in Table S1 and are plotted against buffer concentration in Figure 1. Pronounced curvature is found in dilute buffers. This had been missed in our previous study<sup>5</sup> where only more concentrated buffers had been employed. As seen in Figure 1, rate constants in the more concentrated buffers are approximately linear in buffer concentration. The zero intercept of such plots which we had previously used to calculate  $k_{\text{OH}}$  is obviously not the true rate constant in the absence of buffer.

The rate law which has previously been applied<sup>1,2,5,6</sup> to hemiorthoester decomposition is that of eq 3, with catalysis by H<sup>+</sup>,

$$k_{\text{obsd}} = k_H[\text{H}^+] + k_0 + k_{\text{OH}}[\text{OH}^-] + k_{\text{HA}}[\text{HA}] + k_{\text{A}}[\text{A}^-] \quad (3)$$

OH<sup>-</sup>, buffer acids, and bases and a pH-independent term. This equation is obviously not completely satisfactory in the present case. We can note from our previous work<sup>5,6</sup> that the terms in  $k_H$  and  $k_{\text{HA}}$  are not important in the solutions employed in the present study since pH > 4. The pH-independent term  $k_0$  has a value<sup>6,10</sup> of  $1.5 \text{ s}^{-1}$  so that it does make a contribution. In the analysis the assumption will be made that that contribution is constant, so that it is the variation in  $k_{\text{obsd}} - k_0 \equiv k'_{\text{obsd}}$  which is determining the shapes of the buffer profiles in Figure 1. A brief analysis of the  $k_0$  reaction will be presented at the end of the discussion section.

## Discussion


**Kinetic Analysis.** Curved plots of rate vs. buffer concentration usually mean that a change in rate-limiting step occurs with changing buffer concentration. The reaction of the present study is a particularly simple one, since it involves the breaking of one C-O bond and two proton transfers. The occurrence of a change in rate-limiting step in this case signifies that (a) there must be at least two steps in the reaction and (b) a simple proton transfer step must be rate limiting under some conditions. The reaction

(7) Eigen, M. *Angew. Chem., Int. Ed. Engl.* **1964**, *3*, 1-72.

(8) Curtis, N. J.; Brown, R. S. *J. Org. Chem.* **1980**, *45*, 4038-4040.

(9) Brown, R. S.; Ulan, J. G. *J. Am. Chem. Soc.* **1983**, *105*, 2382-2388.

sequence of eq 4 can be proposed, with the hemiothoester being



converted to its conjugate base prior to breakdown. This deprotonation occurs with hydroxide ion as the base or with the conjugate base form of the buffer. The microscopic reverses of these whereby the neutral hemiothoester is regenerated involve protonation by water and protonation by the acid component of the buffer. Decomposition of the anion occurs by a pH-independent (or water catalyzed) reaction and by a reaction catalyzed by the buffer acid. The actual mechanisms of these reactions and the question of the involvement of a second intermediate  $\text{PhCO}_2\text{CH}_2\text{CH}_2\text{O}^-$  will be discussed later. The acidity constant of the hemiothoester will be defined as  $K_a$ , and the following relationship can be noted,

$$K_a = \frac{k_1 K_W}{k_{-1}} = \frac{k_2 K_{\text{HA}}}{k_{-2}} \quad (5)$$

where  $K_{\text{HA}}$  is the acidity constant of the buffer acid ( $\text{p}K_{\text{HA}} = 4.56$  for acetic acid and 3.56 for formic acid<sup>12</sup>) and  $K_W$  is the autoprotolysis constant ( $\text{p}K_W = 13.7$ ).<sup>12</sup> As will be discussed later,  $\text{p}K_a$  is expected to have a value near 11. This means that for the equilibrium  $\text{TOH} + \text{OH}^- \rightleftharpoons \text{TO}^- + \text{H}_2\text{O}$  it is the deprotonation step which is thermodynamically favored, while for  $\text{TOH} + \text{A}^- \rightleftharpoons \text{TO}^- + \text{HA}$  with acetate and formate buffers it is the protonation step. Consequently, the constants  $k_1$  and  $k_{-2}$  are expected<sup>7</sup> to have values near  $10^{10} \text{ M}^{-1} \text{ s}^{-1}$ . The possibility of deprotonation occurring with a water molecule as the base, and its microscopic reverse, protonation of the anion by  $\text{H}_3\text{O}^+$ , is ignored in our analysis. Although the rate constant for the latter is also expected to be near  $10^{10} \text{ M}^{-1} \text{ s}^{-1}$ , in solutions with  $\text{pH} > 4$  its contribution is relatively unimportant compared to the contribution from the buffer acid and even protonation by water.<sup>11</sup>

The stationary state assumption in  $\text{TO}^-$  produces the following rate expression,

$$k'_{\text{obsd}} = \frac{a[\text{OH}^-] + b[\text{A}^-] + c[\text{H}^+][\text{A}^-]^2}{1 + d[\text{H}^+][\text{A}^-]} \quad (6)$$

where  $a = k_1 k_3 / (k_{-1} + k_3)$ ,  $b = (k_2 k_3 + k_1 k_4 (K_W / K_{\text{HA}})) / (k_{-1} + k_3)$ ,  $c = k_2 k_4 / K_{\text{HA}} (k_{-1} + k_3)$ , and  $d = (k_{-2} + k_4) / K_{\text{HA}} (k_{-1} + k_3)$ . Experimental data for the acetate buffers were fitted to this equation to yield values of the constants:  $a = (5 \pm 2) \times 10^9$ ,  $b = (5.5 \pm 1) \times 10^3$ ,  $c = (1.2 \pm 0.2) \times 10^9$ , and  $d = (8.0 \pm 2) \times 10^{-4}$ . Errors in these coefficients are quite large because of the difficulty of measuring accurately rate constants in very dilute buffer. The constant  $a$  is the  $\text{OH}^-$ -dependent intercept at zero buffer concentration and is particularly difficult to obtain accurately. Although six individual rate constants appear in the kinetic scheme of eq 4, because of the relationships of eq 5, there are only five which are independent. From the experimental fit the four constants  $a$ - $d$  are obtained, so that there is not sufficient information to place a value on all of the rate constants. With the appropriate rearrangement of the expressions for  $a$ - $d$ ,<sup>13</sup> it can be seen that the two constants  $k_1$  and  $k_2$  stand on their own and thus can be assigned a value while the other four can be expressed as ratios,<sup>14</sup> for example, as  $k_{-1}/k_3$  and  $k_{-1}/k_4$ . Values of the constants

(10) This value is measured in HCl solutions where no buffer is present.

(11) For example at pH 4, the most acidic pH of this study, protonation of  $\text{TO}^-$  by  $\text{H}_3\text{O}^+$  will occur with a rate constant of  $10^{10} \times 10^{-4} = 10^6 \text{ s}^{-1}$ . Protonation by water occurs with a rate constant of  $5 \times 10^6 \text{ s}^{-1} - 10^7 \text{ s}^{-1}$  (see analysis).


(12) (a) Values for the constants at 0.1 ionic strength. (b) Harned, H. S.; Owen, B. B. "The Physical Chemistry of Electrolytic Solutions", 3rd ed.; Reinhold Publishing Co.: New York, 1958.

(13)  $a = k_1 / (1 + (k_{-1}/k_3))$ ,  $b = (k_2 + k_1(k_4/k_{-1}))(k_{-1}/k_3)(K_W/K_{\text{HA}}) / (1 + (k_{-1}/k_3))$ ,  $c = k_2(k_4/k_{-1})(k_{-1}/k_3) / K_{\text{HA}}(1 + (k_{-1}/k_3))$  and  $d = ((k_2/k_1)(K_{\text{HA}}/K_W)(k_{-1}/k_3) + (k_4/k_{-1})(k_{-1}/k_3)) / K_{\text{HA}}(1 + (k_{-1}/k_3))$ .

**Table I.** Rate Constants at 25 °C, Ionic Strength 0.1 for the Decomposition of 2-Phenyl-2-hydroxy-1,3-dioxolane in Acetate Buffers

constant	values	
$k_1, \text{M}^{-1} \text{s}^{-1}$	$6 \times 10^9$	
$k_2, \text{M}^{-1} \text{s}^{-1}$	$6 \times 10^3$	
$(k_{-1}/k_3)$	1.4	
$(k_{-1}/k_4)$	0.024	
$\text{p}K_a$	10.50	10.80
$k_{-1}, \text{s}^{-1}$	$4 \times 10^6$	$7 \times 10^6$
$k_{-2}, \text{M}^{-1} \text{s}^{-1}$	$5 \times 10^9$	$1.1 \times 10^{10}$
$k_3, \text{s}^{-1}$	$2.5 \times 10^7$	$5 \times 10^7$
$k_4, \text{M}^{-1} \text{s}^{-1}$	$1.5 \times 10^8$ <sup>a</sup>	$3 \times 10^8$ <sup>b</sup>

<sup>a</sup>  $k_4 = 4.5 \times 10^8$  for formic acid as catalyst and  $\text{p}K_a = 10.50$ . <sup>b</sup>  $k_4 = 9.1 \times 10^8$  for formic acid and  $\text{p}K_a = 10.80$ .



**Figure 2.** Schematic representation of the effect of buffer concentration on the decomposition of the tetrahedral intermediate 2-phenyl-2-hydroxy-1,3-dioxolane.

or ratios determined from the experimental values of  $a$ - $d$  are listed in Table I.


A schematic representation of the rate buffer profile is given in Figure 2. The change in rate-limiting step can be explained as follows. Since  $k_{-1}/k_3$  is less than unity or  $k_3$  is greater than  $k_{-1}$ , in the absence of buffer formation of the anionic intermediate is essentially irreversible. That is, the anion once formed goes on to product rather than being reprotonated by water. The deprotonation step is rate limiting, and since this is occurring with hydroxide ion, the intercepts of the rate-buffer curve are equal to  $k_1[\text{OH}^-]$ . (This is actually only approximate since with  $k_{-1} = 0.14k_3$  there is some reversal of the deprotonation.) The addition of a small amount of buffer results in a significant increase in the rate. This is due to the presence of the base component of the buffer and its participation in the deprotonation reaction. Although the rate constant for the reaction with  $\text{A}^-$  is numerically much smaller than that for  $\text{OH}^-$ , deprotonation via the buffer can occur since the hydroxide ion concentration is so low. At this stage, very dilute buffers, the proton transfer step is still rate limiting (or mainly rate limiting), but as the buffer concentration continues to increase, the rate of protonation of  $\text{TO}^-$  is significantly increased

(14) This is typical of a system of this sort, involving a change in rate-determining step for a reaction with an unstable intermediate. The appropriate kinetic analysis is capable of providing absolute rate constants for intermediate formation, but rate constants for the return of this intermediate to reagent and for the further reaction can only be expressed as ratios.

through reaction with the acid component of the buffer. The stage is ultimately reached where  $k_{-2}[\text{HA}]$  becomes greater than  $k_3$ , and a change in rate-limiting step must occur. Formation of the anionic intermediate becomes reversible with its breakdown being rate limiting. Rate constants do continue to increase with increasing buffer concentration due to a buffer acid catalyzed breakdown. The non-catalyzed breakdown process also occurs in this region. Intercepts of plots of  $k_{\text{obsd}}$  vs.  $[\text{A}^-]$  have values of  $k_3K_a[\text{OH}^-]$ . This corresponds to the hypothetical situation where breakdown is rate limiting and occurs with no buffer catalysis, but the buffer maintains the initial equilibrium. The buffer acid catalyzed breakdown appears in the overall kinetics as general base catalysis,  $k'_{\text{obsd}} = k_4K_a[\text{A}^-]/K_{\text{HA}}$ , because of the pre-equilibrium deprotonation. It is in fact now commonly accepted that the general base catalysis which is observed for the decomposition of aldehyde hydrates, hemiacetals, and hemiothoacid derivatives is occurring by the kinetically equivalent specific base-general acid route.<sup>3,4</sup>

Absolute values of the rate constants  $k_{-1}$ , etc., can be obtained if one further piece of information is available, for example, the acidity constant  $K_a$  for acid dissociation of the hemiothoester. Two approaches based on linear-linear free energy relationships have been suggested for estimating tetrahedral intermediate acidities. Guthrie<sup>15</sup> uses an equation originally proposed by Hine<sup>16</sup> for carbonyl hydrates and estimates<sup>17</sup> a  $\text{p}K_a$  of 10.7 for  $\text{C}_6\text{H}_5\text{C}(\text{OH})_2\text{OMe}$ . Fox and Jencks<sup>18</sup> use a simple alcohol as a starting point; with their approach a value of 10.9 is estimated for the same species. We have recently been able to measure a value for the cyclic hemiothoester, 2-(*p*-nitrophenyl)-2-hydroxy-4,4,5,5-tetramethyl-1,3-dioxolane.<sup>19</sup> The  $\text{p}K_a$  value obtained is 10.4, in reasonable agreement with the estimates, particularly considering the presence of the acid-strengthening nitro substituent. The  $\text{p}K_a$  value for the hemiothoester of concern here is predicted therefore to lie in the range 10.5–11.0. In Table I values are listed for the various constants<sup>20</sup> obtained by using  $\text{p}K_a$  values of 10.5 and 10.8. That a  $\text{p}K$  of this order of magnitude is reasonable can be seen in the fact that values of the rate constant  $k_{-2}$  are produced of the correct order of magnitude,<sup>21</sup> that is somewhere near  $10^{10} \text{ M}^{-1} \text{ s}^{-1}$ .

**Tetrahedral Intermediate Lifetimes.** The lifetime of the hemiothoester anion in its breakdown reactions is obviously extremely short, about 10 ns even in the absence of buffer. This finding of enormous reactivity for a tetrahedral intermediate anion has several implications. As shown in eq 7, for example, the base hydrolysis of an ester can take place in two steps, hydroxide addition to the ester carbonyl followed by elimination of alkoxide.



The carbonyl oxygen exchange which accompanies this hydrolysis requires protonation of the initial intermediate (or perhaps a direct proton switch). In terms of eq 7, a stationary state assumption for the tetrahedral intermediates produces the following equation

- (15) Guthrie, J. P. *J. Am. Chem. Soc.* **1978**, *100*, 5892–5904.  
 (16) Hine, J.; Koser, G. F. *J. Org. Chem.* **1971**, *36*, 1348–1351.  
 (17) Guthrie, J. P.; Cullimore, P. A. *Can. J. Chem.* **1980**, *58*, 1281–1294.  
 (18) Fox, J. P.; Jencks, W. P. *J. Am. Chem. Soc.* **1974**, *96*, 1436–1449.  
 (19) McClelland, R. A.; Seaman, N. E.; Cramm, D. *J. Am. Chem. Soc.* **1984**, *106*, 4511–4515.  
 (20)  $k_{-1} = k_1K_{\text{W}}/K_a$ ;  $k_{-2} = k_2K_{\text{HA}}/K_a$ ;  $k_3 = k_{-1}/(k_{-1}/k_3)$ ;  $k_4 = k_{-1}/(k_{-1}/k_4)$ .

(21) An alternative approach is to set the value of  $k_{-2}$  at  $10^{10} \text{ M}^{-1} \text{ s}^{-1}$  and then calculate the other constants, including  $\text{p}K_a$ . This obviously results in a  $\text{p}K_a$  value of the expected order.


for the ratio of hydrolysis to exchange.

$$\frac{k_{\text{H}}}{k_{\text{E}}} = \left( \frac{2k_2}{k_{-1}} \right) \left( \frac{k_{-1} + k_2 + k_3}{k_3} \right) \quad (8)$$


Very little exchange is in fact observed for alkyl benzoates in base,  $k_{\text{H}}/k_{\text{E}}$  values ranging from 20 to 100.<sup>22</sup> As seen in eq 8, there are two factors which could be responsible for this, an unfavorable partitioning ( $k_2 > k_{-1}$ ) and tetrahedral intermediate breakdown faster than proton transfer ( $k_{-1}, k_2 > k_3$ ). The latter factor has been suggested as being important,<sup>22,23</sup> although its contribution is frequently ignored. The results of the present study suggest that it is a very real possibility.

It is also interesting to speculate on the lifetimes of tetrahedral intermediate anions with leaving groups such as aryloxy, carboxylate, or even halide. If the expulsion of alkoxide can occur with a rate constant of the order of  $10^7$ – $10^8 \text{ s}^{-1}$ , the situation could easily be reached with these significantly better leaving groups where the lifetime of the intermediate becomes so short that it does not exist.<sup>24</sup> In such a case the tetrahedral species becomes a transition state and the nucleophilic displacement reaction would be regarded as being concerted. The possibility of a one-step acyl-transfer reaction has recently been raised for several quite different systems.<sup>25–28</sup> In one case<sup>27</sup> arguments were based on tetrahedral intermediate lifetimes. Experiments are currently underway to investigate this possibility.


**Nature of Anion Breakdown.** The acid-catalyzed decomposition of the hemiothoester anion ( $k_4$  process) has a Brønsted  $\alpha$  value of 0.5, calculated by using data for the two buffers—acetic acid and formic acid.<sup>29</sup> This intermediate  $\alpha$  value implies a concerted reaction with proton transfer and C–O bond breaking occurring simultaneously. The rate constant  $k_3$  for the pH-independent



breakdown lies four orders of magnitude above the extrapolated line for these two acids. Although this extrapolation is a lengthy one, the implication is that the reaction is mechanistically different. Possibly in this case the alkoxide anion is formed as a discrete intermediate.




The possibility of catalysis by  $\text{H}_3\text{O}^+$  can also be considered.



This reaction would correspond overall to a pH-independent process in the decomposition of the neutral hemiothoester, and it is thus a possible candidate for the observed noncatalyzed reaction ( $k_0$  term in eq 3). The pH-independent rate constant for eq 11 is equal to  $k_6K_a$  and the observed  $k_0$  is 1.5. Therefore, since  $\text{p}K_a = 10.5$ – $11.0$ ,  $k_6$  is required to be near  $10^{11} \text{ M}^{-1} \text{ s}^{-1}$ , that is slightly greater than diffusion limited. This suggests that the

- (22) Shain, S. A.; Kirsch, J. F. *J. Am. Chem. Soc.* **1968**, *90*, 5848–5854.  
 (23) Bender, M. L.; Thomas, R. J. *J. Am. Chem. Soc.* **1961**, *83*, 4189–4193. Note that some of the experimental numbers in this particular study were later shown to be incorrect.<sup>22</sup>  
 (24) Jencks, W. P. *Acc. Chem. Res.* **1980**, *13*, 161–169.  
 (25) Curran, T. C.; Ferrar, C. R.; Niazzy, O.; Williams, A. *J. Am. Chem. Soc.* **1980**, *102*, 6828–6837.  
 (26) Khim, J. K.; Caserio, M. *J. Am. Chem. Soc.* **1981**, *103*, 2124–2127.  
 (27) Ritchie, C. D.; Van Verth, J. E.; Virtanen, P. O. I. *J. Am. Chem. Soc.* **1982**, *104*, 3491–3497.  
 (28) De Tar, D. F. *J. Am. Chem. Soc.* **1982**, *104*, 7205–7212.  
 (29) For formic acid the straight line segment at high buffer concentration (Figure 1) is used to obtain a value for  $k_4K_a/K_{\text{HA}}$  (see Figure 2). Note that the  $\alpha$  value is not dependent on the value of  $K_a$  chosen, since this same constant is used to calculate  $k_4$  values for each buffer.

reaction of eq 11 does not make a significant contribution to  $k_0$ . The possibility has been raised<sup>1,2</sup> of a concerted process for this reaction, with solvent water molecules acting as acid and base to transfer protons to and from the neutral hemiotherester.



**Acknowledgment.** The continued financial support of the National Sciences and Engineering Research Council of Canada is gratefully acknowledged.

**Registry No.** 1, 4353-38-2; 2, 59700-73-1; 3, 92544-12-2.

**Supplementary Material Available:** Table SI listing observed rate constants for the breakdown of 2-phenyl-2-hydroxy-1,3-dioxolane in acetic and formate buffers (1 page). Ordering information is given on any current masthead page.

## Mechanisms of Elimination Reactions. 38. Why Is the Effect of Successive $\beta$ -Alkyl Substitution on the Rates of Elimination from Quaternary Ammonium Salts Nonadditive?<sup>1</sup>

Shune-Long Wu, Yu-Tai Tao,<sup>2</sup> and William H. Saunders, Jr.\*

Contribution from the Department of Chemistry, University of Rochester, Rochester, New York 14627. Received June 8, 1984

**Abstract:** Possible reasons are examined for the nonadditivity of the effect of successive  $\beta$ -methyl substitution on the rates of elimination reactions of quaternary ammonium salts. The temperature dependences of the deuterium isotope effects in E2 reactions of  $R_1R_2NMe_2^+$  show that tunneling is not a significant source of nonadditivity. Neither is a change of gross mechanism or stereochemistry, for studies with  $C_4H_9CHDCHDNMe_3^+$  and  $C_4H_9(CH_3)CHCHDNMe_3^+$  show the reactions to be very predominantly (>88%) anti-E2 in both cases. Secondary tritium isotope effects with  $R_1R_2CHCHTNMe_3^+$  increase, however, in the order ethyl ( $1.108 \pm 0.002$ ), propyl ( $1.150 \pm 0.015$ ), isobutyl ( $1.216 \pm 0.012$ ). This result suggests increasing rehybridization at the  $\alpha$ -carbon in the transition state and therefore a shift toward less E1cB and more central-E2 character. Since methyl substitution is expected to favor a developing double bond, the much smaller rate-depressing effect of the second  $\beta$ -methyl is accounted for by such a shift in transition-state character.

Substituent effects on rates and equilibria in organic reactions are usually assumed to be approximately additive. This assumption is often valid for remote substituents; e.g., the Hammett  $\sigma$  for 3,4-dichlorophenyl is well represented by  $\sigma_{p-Cl} + \sigma_{m-Cl}$ .<sup>3</sup> When substituents are closer to the reaction center, saturation effects may be observed. Chloroacetic acid is 79 times as acidic as acetic acid, but dichloroacetic acid is only 37 times as acidic as chloroacetic acid.<sup>4</sup> This saturation effect is still modest. When successive substituents affect each other's ability to undergo resonance interaction with the reaction center, dramatic effects often result. Diphenylmethane is  $10^8$  times as acidic as toluene, but triphenylmethane is less than  $10^2$  times as acidic as diphenylmethane.<sup>4</sup>

We have been intrigued for some time by a pattern of non-additive substituent effects for certain E2 reactions. In the Hofmann elimination at 85–150 °C of quaternary ammonium hydroxide,  $R_1R_2NMe_2^+OH^-$ , the ethylene/propylene ratio when  $R_1 = Et$  and  $R_2 = n-Pr$  is 41, but the propylene/isobutylene ratio when  $R_1 = n-Pr$  and  $R_2 = i-Bu$  is only 2.7.<sup>5</sup> In other words, the effect of the second  $\beta$ -methyl group is only one-fifteenth that of the first. A similar, though somewhat less dramatic effect persists in direct rate measurements. The relative rates of the elimination reactions of  $RNMe_3^+$  with ethoxide in ethanol at 104.2 °C run

**Table I.** Temperature Dependence of the Product Ratios for the Reactions of  $R_1R_2NMe_2^+Br^-$  with Sodium 1-Butoxide in 50% 1-Butanol-Dimethyl Sulfoxide

T, °C	ethylene/ propylene		propylene/isobutylene		
	1 <sup>a</sup>	2 <sup>a</sup>	3 <sup>a</sup>	4 <sup>a</sup>	5 <sup>a</sup>
50	68.12				
60	59.80	19.88	2.594	0.7072	11.59
70	49.65	16.20			
80			2.696	0.8458	10.80
90	38.42	13.20	2.764	0.8748	10.65
100		12.50	2.892	0.9059	10.12

<sup>a</sup> See text for structures of these compounds. Each product ratio is the average of two runs. Derivations from mean run usually 1–2% and never above 4%.

$Et/n-Pr = 13.8$  and  $n-Pr/i-Bu = 3.1$ .<sup>6</sup> To get the true chemical effects, the above values should be corrected for symmetry number differences. The corrections can most simply be understood from the fact that there are three indistinguishable anti-E2 transition states for loss of ethyl, two enantiomeric ones for loss of propyl, but only one for loss of isobutyl. The ratios from  $R_1R_2NMe_2^+OH^-$  then become ethylene/propylene 27.3 and propylene/isobutylene 1.35, while the relative rates from  $RNMe_3^+$  become  $Et/n-Pr$  9.20 and  $n-Pr/i-Bu$  1.55.

Even more interesting is the fact that this nonadditivity is almost completely absent in the reactions of the corresponding sulfonium ions,  $RSMe_2^+$ . Here the relative rates of reaction with ethoxide

(1) This work was supported by the National Science Foundation.

(2) Sherman Clarke fellow, 1979–1981.

(3) Hine, J. "Structural Effects on Equilibria in Organic Chemistry"; Wiley: New York, 1975; pp 86–88.

(4) Lowry, T. H.; Richardson, K. L. "Mechanism and Theory in Organic Chemistry", 2nd ed.; Harper and Row: New York, 1981; Chapter 3.

(5) Cope, A. C.; LeBel, N. A.; Lee, H. H.; Moore, W. R. *J. Am. Chem. Soc.* 1957, 79, 4720–4729.

(6) Banthorpe, D. V.; Hughes, E. D.; Ingold, C. K. *J. Chem. Soc.* 1960, 4054–4087.