



Table I. Protonolysis Reactions of Silyl Hydride Anions, 298 K

anion	reference acids ^a					inference ^b
	EtOH (376.1)	MeOH (381.4)	H ₂ O (390.7)	D ₂ O (392.9)	NH ₃ (403.6)	
SiH ₅ ⁻ (1)	+	+	-	-	-	18.0 ≤ HA ≤ 26.9
<i>n</i> -C ₅ H ₁₁ SiH ₄ ⁻ (2)	+	+	+	+	-	5.3 ≤ HA ≤ 16.3
Et ₃ SiH ₂ ⁻ (3)	+	+	+	+	-	4.7 ≤ HA ≤ 15.6

^a $\Delta H_{\text{acid}}(\text{HX})$ shown in parentheses in kcal/mol.¹² A + sign indicates occurrence of protonolysis and formation of the corresponding conjugate base anion. ^b Estimated limits on the hydride affinity (HA) of the corresponding neutral silane in kcal/mol. Correction factors which take into account changes in rotational and translational entropy have been incorporated.

of the thermochemical cycle in Scheme I, where HA(R₄Si) is the hydride affinity of the silane $D[\text{R}_4\text{Si}-\text{H}^-]$ and $\Delta H_{\text{acid}}(\text{HX})$ and $\Delta H_{\text{acid}}(\text{H}_2)$ are known quantities.¹² Thus, by monitoring the occurrence or nonoccurrence of reaction 4 for a particular silyl hydride and a series of acids of varying acidity, one can derive limits on the hydride affinity of the silane. Table I presents the results of such "bracketing" experiments for three reactant ions. Our experimentally derived limits on the hydride binding energies are in good agreement with the computed values for SiH₄ reported by Baybutt (18.6 kcal/mol, HF),¹⁶ Wilhite and Spialter (16.9 kcal/mol, HF),¹⁷ Keil and Ahlrichs (20.3 kcal/mol, PNO CI),¹⁸ and most recently by Schleyer (18.3 kcal/mol, MP4/6-31++G**).¹⁹ Hydride affinities of this magnitude are among the lowest known for any neutral molecule and are comparable to those of NH₃ (8 kcal/mol),²⁰ H₂O (16 kcal/mol),²¹ and benzene (25 kcal/mol).²²

The facility with which the silyl hydride ions undergo binary hydride transfer reactions combined with their ready adaptability to isotope labeling makes them ideal systems for investigating the structures and fluxional behavior of isolated 10-Si-5 negative ions. Small Berry pseudorotational barriers have been computed for SiH₅⁻ (ca. 2.7 kcal/mol)^{17,18} and measured experimentally for 10-Si-5 phenyl siliconates in solution.^{4,23} Therefore, one may anticipate that the hydrogens in 1–3 undergo rapid intramolecular scrambling once the ions are formed. To test this, we have generated in situ monodeuterated alkylsilyl hydride ions using D⁻ (from ND₃) in conjunction with reaction 1. Addition of CO₂ to the system as a convenient hydride acceptor is followed by determination of the primary relative yields of HCO₂⁻ and DCO₂⁻ products. In the absence of a kinetic isotope effect for transfer of H⁻ vs. D⁻, chemical equivalence of the hydrogens in the reactant ions would be indicated by statistical yields of HCO₂⁻ and DCO₂⁻. The results for ions 2-d and 3-d are shown below (eq 5 and 6),



where the indicated branching ratios have been corrected for the natural isotope distributions in each ion and the contribution to the HCO₂⁻ yield from the unavoidable presence of unlabeled reactant ion.²⁴ An isotope effect is immediately evident since the determined yields deviate significantly from the 1:1 and 3:1 statistical values. However, the deviation is a constant $k_{\text{H}}/k_{\text{D}} = 2.1$ for the two systems, so we may conclude that complete

scrambling of H and D has occurred in the silicon ions prior to the probe reaction with CO₂. Further experiments with Et₂SiH₂D⁻, SiH₄D⁻, and other isotopomers are in progress to confirm this finding.

Acknowledgment. We are grateful to Professor P. v. R. Schleyer for disclosure of his computational results for SiH₅⁻ in advance of publication. Support from the National Science Foundation (Grant CHE-8502515) is gratefully acknowledged.

Proton Transfer as a Function of Hydration Number and Temperature: Rate Constants and Product Distributions for OH⁻(H₂O)_{0,1,2,3} + HF at 200–500 K

Peter M. Hierl,^{*1,2} Anton F. Ahrens,^{2,3}
Michael Henchman,^{*4} A. A. Viggiano,⁵ and John F. Paulson

Air Force Geophysics Laboratory
Hanscom AFB, Massachusetts 01731

D. C. Clary

University Chemical Laboratory
Cambridge CB2 1EW, U.K.

Received January 2, 1986

Proton-transfer reactions which are exothermic in the gas phase are exceedingly efficient—most reacting at essentially every collision.^{6–10} Several studies have shown how this extreme reactivity is influenced by solvation in the gas phase at 300 K.^{8,9,11–16} Here we report how hydration influences the rate constants and the product distributions of the reaction



throughout the temperature range 200–500 K. Such studies in the gas phase—where the solvation number is systematically changed—examine the role of solvate in the transfer of protons.¹⁸

- (1) SCEE Fellow/AFOSR Summer Faculty Research Associate, 1984.
- (2) Permanent address: Department of Chemistry, University of Kansas, Lawrence, KS 66045.
- (3) SCEE Fellow/AFOSR Summer Graduate Research Associate, 1984.
- (4) AFSC-URRP Visiting Professor, 1984–1986. Permanent address: Department of Chemistry, Brandeis University, Waltham, MA 02254.
- (5) System Integration Engineering Inc., Lexington, MA 02173: under contract to AFGL.
- (6) Bohme, D. K. In *Interactions between Ions and Molecules*; Ausloos, P., Ed.; Plenum: New York, 1975; p 489.
- (7) Bohme, D. K.; Mackay, G. I.; Schiff, H. I. *J. Chem. Phys.* **1980**, *73*, 4976.
- (8) Bohme, D. K. *Trans. R. Soc. Can., Sect. 3* **1981**, *19*, 265.
- (9) Bohme, D. K. In *Ionic Processes in the Gas Phase*; Almoester Ferreira, M. A., Ed.; Reidel: Dordrecht, 1984; p 111.
- (10) Moylan, C. R.; Brauman, J. I. *Annu. Rev. Phys. Chem.* **1983**, *34*, 187.
- (11) Bohme, D. K.; Mackay, G. I.; Tanner, S. D. *J. Am. Chem. Soc.* **1979**, *101*, 3724.
- (12) Smith, D.; Adams, N. G.; Henchman, M. J. *J. Chem. Phys.* **1980**, *72*, 4951.
- (13) Tanner, S. D.; Mackay, G. I.; Bohme, D. K. *Can. J. Chem.* **1981**, *59*, 1615.
- (14) Bohme, D. K.; Rakshit, A. B.; Mackay, G. I. *J. Am. Chem. Soc.* **1982**, *104*, 1100.
- (15) Mackay, G. I.; Rakshit, A. B.; Bohme, D. K. *Can. J. Chem.* **1982**, *60*, 2594.
- (16) Rakshit, A. B.; Bohme, D. K. *Can. J. Chem.* **1983**, *61*, 1683.
- (17) Bartmess, J. E.; McIver, R. T., Jr. In *Gas Phase Ion Chemistry*; Bowers, M. T., Ed.; Academic Press: New York, 1979; Vol. 2, p 90. The value is unaltered by the redetermination of E. A. [OH]. Schulz, P. A.; Mead, R. D.; Jones, P. L.; Lineberger, W. C. *J. Chem. Phys.* **1982**, *77*, 1153.

- (16) Baybutt, P. *Mol. Phys.* **1975**, *29*, 389.
- (17) Wilhite, D. L.; Spialter, L. *J. Am. Chem. Soc.* **1973**, *95*, 2100.
- (18) Keil, F.; Ahlrichs, R. *Chem. Phys.* **1975**, *8*, 384.
- (19) Schleyer, P. v. R., private communication.
- (20) (a) Kleingeld, J. C.; Ingemann, S.; Jalonen, J. E.; Nibbering, N. M. *J. Am. Chem. Soc.* **1983**, *105*, 2474. (b) Kalcher, J.; Rosmus, P. *Can. J. Phys.* **1984**, *62*, 1323. (c) Cremer, D.; Kraka, E. *J. Phys. Chem.* **1986**, *90*, 33. (d) Coe, J. V.; Snodgrass, J. T.; Freidhoff, C. B.; McHugh, K. M.; Bowen, K. H. *J. Chem. Phys.*, in press.
- (21) (a) Kleingeld, J. C.; Nibbering, N. M. *Int. J. Mass Spectrom. Ion Phys.* **1983**, *49*, 311. (b) Paulson, J. F.; Henchman, M. J. *Bull. Am. Phys. Soc.* **1982**, *27*, 108.
- (22) Lane, K. R.; Sallans, L.; Squires, R. R. *J. Am. Chem. Soc.*, in press.
- (23) Damrauer, R.; Danahey, S. E. *Organometallics*, in press.
- (24) Ions 2-d and 3-d undergo secondary H/D exchange reactions with the excess neutral silane precursor in the flow tube eventually losing the label.




Figure 1. Rate constants for the reactions $\text{OH}^-(\text{H}_2\text{O})_{0,1,2,3} + \text{HF}$ as a function of temperature. For each reaction, data points are compared with theoretical predictions, see text.

Reaction 1—a proton-transfer reaction involving only four atoms—has been studied in a selected ion flow tube (SIFT)¹⁹ in the temperature range 200–500 K. Figure 1 shows how the experimental rate constants (each a mean of at least three determinations) decrease monotonically with increasing temperature. The data agree, within the limits of experimental error, with theoretical predictions for collision rate constants, shown as solid lines.^{20,21} These are derived by using the adiabatic capture model (ACCSA) due to Clary;²² this model has already shown impressive predictive success for proton-transfer reactions in this temperature range.²³ We therefore conclude that the temperature dependence shown in Figure 1 is that of the physical collision efficiency and not that of the chemical reaction efficiency. Proton transfer, efficient at 300 K,^{6–10} is efficient throughout 200–500 K.^{23,24}

The effect of hydration on reaction 1 is also indicated in Figure 1, where rate constants are shown for the hydroxide ion hydrated with one, two, and three water molecules. Agreement between experiment and theory is seen to be similar for the hydrated and for the unhydrated reactants.²⁵ Proton transfer occurs on essentially every collision throughout the range of hydration^{8–16} and temperature²⁶ studied.

Hydrating the reactant gives identical reaction efficiencies: in contrast, it gives different reaction products. In Figure 2, the product distributions²⁷ are reported as branching ratios, and reaction enthalpies²⁸ are also shown for each possible channel.

(18) Grunwald, E.; Eustace, D. In *Proton Transfer Reactions*; Caldin, E. F., Gold, V., Eds.; Chapman and Hall: London, 1975; pp 103–120.

(19) Viggiano, A. A.; Paulson, J. F. *J. Chem. Phys.* **1983**, *79*, 2241. Miller, T. M.; Wetterskog, R. E.; Paulson, J. F. *Ibid.* **1984**, *80*, 4922.

(20) The theory treats only ion–dipole and ion-induced dipole attraction, ignoring the dipole–dipole attraction, estimated to raise the rate constant in this case by up to 20%.

(21) Agreement with other theoretical predictions is within 20%.²² Su, T.; Chesnavich, W. J. *J. Chem. Phys.* **1982**, *76*, 5183.

(22) Clary, D. C. *Mol. Phys.* **1984**, *53*, 3; **1985**, *54*, 605.

(23) Clary, D. C.; Smith, D.; Adams, N. G. *Chem. Phys. Lett.* **1985**, *119*, 320.

(24) For other examples, see: Mautner, M. In *Gas Phase Ion Chemistry*; Bowers, M. T., Ed.; Academic Press: New York, 1979; Vol. 1, pp 207–215. Lindinger, W.; Smith, D. In *Reactions of Small Transient Species*; Fontijn, A., Clyne, M. A. A., Eds.; Academic Press: New York, 1983; p 403.

(25) Rate constants decrease with increasing hydration only according to the dependence of collision frequency on reduced mass.

(26) Fehsenfeld, F. C.; Dotan, I.; Albritton, D. L.; Howard, C. J.; Ferguson, E. E. *J. Geophys. Res.* **1978**, *83* (C3), 1333.

(27) In the SIFT technique, selection of the reactant ion allows accurate measurement of the product distribution. Smith, D.; Adams, N. G. In *Gas Phase Ion Chemistry*; Bowers, M. T., Ed.; Academic Press: New York, 1979; Vol. 1, pp 28–29.





Figure 2. Enthalpy diagram for the reactants (left column) and for the products (right column), identified by the ionic species. Relative enthalpies²⁸ yield reaction enthalpies by subtraction. Experimental product distributions (independent of temperature) are reported with error limits of 0.25% for $\text{OH}^-(\text{H}_2\text{O})$, $\pm 5\%$ for $\text{OH}^-(\text{H}_2\text{O})_2$, and $\pm 10\%$ for $\text{OH}^-(\text{H}_2\text{O})_3$.

Because the reactions are so efficient, they must be exothermic. For example, $\text{OH}^-\cdot\text{H}_2\text{O}$ gives $\text{F}^-\cdot\text{H}_2\text{O}$ ($\Delta H^\circ = -17.8$ kcal/mol) exclusively and no F^- ($\Delta H^\circ = +5.5$ kcal/mol). (Our shorthand notation for the reaction $\text{OH}^-\cdot\text{H}_2\text{O} + \text{HF} \rightarrow \text{F}^-\cdot\text{H}_2\text{O} + \text{H}_2\text{O}$ is 1 \rightarrow 1, indicating ionic solvation numbers.) For $\text{OH}^-(\text{H}_2\text{O})_2$, two reactions are possible: 2 \rightarrow 1, 2, and for $\text{OH}^-(\text{H}_2\text{O})_3$, there is also a choice: 3 \rightarrow 2, 3. In both cases, the preferred channels, 2 \rightarrow 1 and 3 \rightarrow 2, are given by a simple propensity rule: The most efficient channel is the least exothermic, yielding the ionic product with the minimum number of solvate molecules.^{12,15,16}

Up to three molecules of water have been observed to transfer efficiently from ionic reactant to ionic product, in order to drive the proton-transfer reaction. These experiments demonstrate solvate participation in proton transfer in the absence of solvent. They reflect, in the gas phase, what Grunwald's NMR studies demonstrated in solution over 2 decades ago.¹⁸

When a solvate molecule is transferred, a proton comes to the ionic reactant (or base) and a water molecule leaves from it (formally an OH^- transfer).



Within the transition state, the transferred water molecule is at the center of charge. Concomitant transfer of solvate and negative charge (from left to right) is an efficient pathway of low energy because solvate and charge are never separated. This is in marked contrast to the nucleophilic displacement reactions described in the accompanying communication.²⁹

Consider the following mechanism. Initially a substantial solvation energy (~ 15 kcal/mol) binds the HF to the ion within the intermediate,²⁸ then proton transfer within the intermediate liberates reaction exothermicity as excitation energy, and finally, to the maximum extent possible, solvate water molecules are “boiled off”.^{9,11,30} According to this mechanism, channels 2 \rightarrow 2 and 2 \rightarrow 1 result from the sequential loss of one and two waters, respectively, from the intermediate: $\text{F}^-(\text{H}_2\text{O})_3 \rightarrow \text{F}^-(\text{H}_2\text{O})_2 \rightarrow \text{F}^-\cdot\text{H}_2\text{O}$. If, in the loss of the first water, more than a total of 1.4 kcal/mol is released as relative translational energy and excitation energy in the expelled water, loss of a second water becomes endothermic: the experimental result is that 43% lose only one water. The mechanism predicts that the channels 3 \rightarrow 3 and 3 \rightarrow 2 follow the sequence $\text{F}^-(\text{H}_2\text{O})_4 \rightarrow \text{F}^-(\text{H}_2\text{O})_3 \rightarrow \text{F}^-(\text{H}_2\text{O})_2$. Here “loss” of 3.0 kcal/mol in the expulsion of the first water makes the loss of the second water endothermic, and here only 15% lose only one water. The experimental product distributions are therefore qualitatively consistent with the predictions of the “boil-off” model.³⁰

(28) Kebarle, P. *Annu. Rev. Phys. Chem.* **1977**, *28*, 445.

(29) Hierl, P. M.; Ahrens, A. F.; Henchman, M.; Viggiano, A. A.; Paulson, J. F.; Clary, D. C. *J. Am. Chem. Soc.*, following paper in this issue.

Registry No. OH⁻, 14280-30-9; HF, 7664-39-3; OH⁻(H₂O), 23138-14-9; OH⁻(H₂O)₂, 34118-36-0; OH⁻(H₂O)₃, 34118-37-1.

Supplementary Material Available: Table of numerical values of experimental and theoretical rate constants shown in Figure 1 (1 page). Ordering information is given on any current masthead page.

(30) Kebarle, P. *Adv. Chem. Ser.* **1968**, 72, 24. Fehsenfeld, F. C.; Ferguson, E. E. *J. Chem. Phys.* **1973**, 59, 6272.

Nucleophilic Displacement as a Function of Hydration Number and Temperature: Rate Constants and Product Distributions for OD⁻(D₂O)_{0,1,2} + CH₃Cl at 200–500 K

Peter M. Hierl,^{*1,2} Anton F. Ahrens,^{2,3}
Michael Henchman,^{*4} A. A. Viggiano,⁵ and John F. Paulson


*Air Force Geophysics Laboratory
Hanscom AFB, Massachusetts 01731*

D. C. Clary

*University Chemical Laboratory
Cambridge CB2 1EW, U.K.*

Received January 2, 1986

Nucleophilic displacement reactions which are exothermic do not react on every collision in the gas phase.^{6,7} They exhibit a negative temperature dependence, rate constants *decreasing* with *increasing* temperature,⁸ and reaction at 300 K is quenched dramatically by the addition of only one–three solvate molecules.⁹ In each respect nucleophilic displacement differs from proton transfer, as contrasted in the companion paper.¹⁰ Here we report how hydration influences the rate constant *and* the product distribution of the nucleophilic displacement reaction



within the temperature range 200–500 K. Such data invite interpretation using hypersurfaces calculated for hydrated reactants.¹²

Rate constants for reaction 1 have been measured with a selected ion flow tube (SIFT), using techniques similar to those used in the companion study.¹⁰ Because the OH⁻(H₂O) reactant and the ³⁵Cl⁻ product have the same mass-to-charge ratio (*m/e* = 35), perdeuterated anions, produced from D₂O in the ion source, were used throughout. Rate constants, for the process OD⁻(D₂O)_{*n*} +




Figure 1. Reaction efficiencies for the reactions OD⁻(D₂O)_{0,1,2} + CH₃Cl as a function of temperature.




Figure 2. Enthalpy diagram for the reactants (left column) and products (right column), identified by the ionic species. Relative enthalpies are for undeuterated species¹⁶ and yield reaction enthalpies by subtraction. Experimental product distributions (independent of temperature) are reported with error limits of $\pm 3\%$ for OD⁻(D₂O) and $\pm 10\%$ for OD⁻(D₂O)₂.

CH₃Cl → products, were measured for the mono- and dihydrate (*n* = 1, 2) but were too small to measure for *n* = 3.¹³ The data at 300 K agree, within $\pm 30\%$, with data obtained by Bohme and colleagues using the flowing afterglow technique.⁹

Figure 1 shows the dependence of the rate constant on temperature and solvation number. What is represented is the *reaction efficiency*, which is the ratio of the experimental rate constant to a theoretical collision rate constant. Collision rate constants have been calculated by using the ACCSA procedure, developed by Clary.^{13,14}

Reaction 1 shows the *negative* temperature dependence that has been shown experimentally⁸ and has been predicted theoretically^{15,8} for nucleophilic displacement at an sp³ carbon atom.

(1) SCEEE Fellow/AFOSR Summer Faculty Research Associate, 1984.
(2) Permanent address: Department of Chemistry, University of Kansas, Lawrence, KS 66045.

(3) SCEEE Fellow/AFOSR Summer Graduate Research Associate, 1984.
(4) AFSC-URRP Visiting Professor, 1984–1986. Permanent address: Department of Chemistry, Brandeis University, Waltham, MA 02254.

(5) System Integration Engineering Inc., Lexington, MA 02173: under contract to AFGL.

(6) Tanaka, K.; Mackay, G. I.; Payzant, J. D.; Bohme, D. K. *Can. J. Chem.* **1976**, 54, 1643.

(7) Olmstead, W. N.; Brauman, J. I. *J. Am. Chem. Soc.* **1977**, 99, 4219.

(8) (a) Sen Sharma, D. K.; Kebarle, P. *J. Am. Chem. Soc.* **1982**, 104, 19.

(b) Magnera, T. F.; Kebarle, P. In *Ionic Processes in the Gas Phase*, Almoester Ferreira, M. A., Ed.; Reidel: Dordrecht, 1984; p 135. (c) Caldwell, G.; Magnera, T. F.; Kebarle, P. *J. Am. Chem. Soc.* **1984**, 106, 959. (d) Sharma, R. B.; Sen Sharma, D. K.; Hiraoka, K.; Kebarle, P. *J. Am. Chem. Soc.* **1985**, 107, 3747. (e) Grimsrud, E. P.; Chowdhury, S.; Kebarle, P. *J. Chem. Phys.* **1985**, 83, 1059.

(9) (a) Bohme, D. K.; Mackay, G. I. *J. Am. Chem. Soc.* **1981**, 103, 978.

(b) Bohme, D. K. In *Ionic Processes in the Gas Phase*, Almoester Ferreira, M. A., Ed.; Reidel: Dordrecht, 1984; p 111. (c) Bohme, D. K.; Raksit, A. B. *J. Am. Chem. Soc.* **1984**, 106, 3447.

(10) Hierl, P. M.; Ahrens, A. F.; Henchman, M.; Viggiano, A. A.; Paulson, J. F.; Clary, D. C. *J. Am. Chem. Soc.*, preceding paper in this issue.

(11) The range of the estimates, -47.5^7 to -50 kcal/mol,^{6c} indicates the accuracy.

(12) Morokuma, K. *J. Am. Chem. Soc.* **1982**, 104, 3732. Morokuma, K.; Kato, S.; Kitaura, K.; Obara, S.; Ohta, K.; Hanamura, M. In *New Horizons of Quantum Chemistry*; Lowdin, P.-O., Pullman, B., Eds.; Reidel: Dordrecht, 1983; p 221.

(13) Numerical values are given in the supplemental material.

(14) Clary, D. C. *Mol. Phys.* **1984**, 53, 3; **1985**, 54, 605.

(15) Farneth, W. E.; Brauman, J. I. *J. Am. Chem. Soc.* **1976**, 98, 7891. Olmstead, W. P. Ph.D. Thesis, Stanford University, CA, 1977 (University Microfilms, Ann Arbor, MI, no. 77-12679), p 189. Brauman, J. I. In *Kinetics of Ion-Molecule Reactions*; Ausloos, P., Ed.; Plenum: New York, 1979; p 153.