


# Dihydride Transfer. A Bimolecular Mechanism in the Isomerization of *cis*-Dihydridobromo(carbonyl)-(bis(diphenylphosphino)ethane)iridium, IrH<sub>2</sub>Br(CO)(dppe)

Amanda J. Kunin, Curtis E. Johnson,<sup>†</sup> John A. Maguire, William D. Jones,\* and Richard Eisenberg\*


Contribution from the Department of Chemistry, University of Rochester, Rochester, New York 14627. Received August 11, 1986

**Abstract:** The oxidative addition of H<sub>2</sub> to IrBr(CO)(dppe) (**2**) (dppe = 1,2-bis(diphenylphosphino)ethane) yields a kinetic dihydride species, **3**, which then isomerizes to a more stable isomer, **4**. This isomerization of **3** to **4** has been studied kinetically as a function of initial H<sub>2</sub> pressure. Two pathways are operative at ambient temperature. The first is a reductive elimination/oxidative addition sequence which is first order in complex, while the second is a bimolecular pathway involving dihydride transfer from **3** to **2** to produce **4** and regenerate **2**. The dihydride transfer pathway is second order in complex and becomes the dominant isomerization mechanism when less than 1 equiv of H<sub>2</sub> relative to **2** has been added to the system. All of the kinetic data have been fit to a complete rate law which leads to a bimolecular rate constant for dihydride transfer of 3.5 × 10<sup>-3</sup> M<sup>-1</sup> s<sup>-1</sup>. Below -20 °C, the dihydride transfer pathway for isomerization is the only one operating.

The oxidative addition of H<sub>2</sub> to d<sup>8</sup> metal complexes has been extensively studied over the past 20 years because of its relevance to H<sub>2</sub> activation in homogeneous hydrogenation and hydroformylation.<sup>1,2</sup> One of the most thoroughly investigated systems in this context is Vaska's complex, *trans*-IrCl(CO)(PPh<sub>3</sub>)<sub>2</sub> (**1**), which reacts with H<sub>2</sub> according to eq 1.<sup>3</sup> On the basis of kinetic and mechanistic studies,<sup>3-5</sup> H<sub>2</sub> oxidative addition is generally viewed as a concerted process with a triangular MH<sub>2</sub> transition state leading to a *cis* dihydride product.



Recently we began investigating the oxidative addition chemistry of the related set of *cis* phosphine complexes IrX(CO)(dppe), **2**, and have discovered that its concerted addition reactions proceed under kinetic control.<sup>6,7</sup> With complexes **2**, the oxidative addition of H<sub>2</sub> can follow two possible pathways, i and ii, as shown in eq 2, leading to different diastereomers, **3** and **4**, respectively, for



the concerted process. Pathway i corresponds to H<sub>2</sub> approach to the square-planar complex with the molecular axis of H<sub>2</sub> parallel to P-Ir-CO as shown in A. The concerted oxidative addition along i takes place with a bending of the *trans* P-Ir-CO axis so that one hydride of the product becomes *trans* to CO and the other *trans* to P. Pathway ii corresponds to approach with the H<sub>2</sub>

molecular axis parallel to P-Ir-X, as shown in B, and addition occurs with bending of the P-Ir-X axis.

While two diastereomers thus exist for concerted H<sub>2</sub> oxidative addition to IrX(CO)(dppe), we have found that for X = Cl, Br, I, H, and CN the initial oxidative addition takes place diastereoselectively along pathway i.<sup>6</sup> The reaction with H<sub>2</sub> in solution is essentially complete within 1 min under an atmosphere of H<sub>2</sub>, forming isomer **3** to >99%. That the reaction proceeds under kinetic control is illustrated by the fact that for X = Cl, Br, and I, the initially formed diastereomer slowly equilibrates with the more stable diastereomer corresponding to oxidative addition along pathway ii. On the basis of the variation of X, a steric basis for the diastereoselectivity of H<sub>2</sub> oxidative addition was ruled out, leaving ligand electronic effects as the controlling factor in the diastereoselection process.

The isomerization of the kinetic isomer **3** to the thermodynamic isomer **4** for X = Br was also examined by us in detail.<sup>6a</sup> On the basis of the observation that **3** rapidly forms **3-d<sub>2</sub>** when placed under D<sub>2</sub>, it was determined that the initial oxidative addition is rapid and reversible, occurring much faster than isomerization. The isomerization in acetone under H<sub>2</sub> follows clean first-order kinetics with an observed rate constant, *k*<sub>obsd</sub>, at 55 °C of 1.85 × 10<sup>-4</sup> s<sup>-1</sup>, corresponding to a half-life of 62 min. At 25 °C the half-life of the kinetic isomer **3** is about 35 h. Two possible mechanisms for isomerization appeared consistent with the kinetic data. The first was an intramolecular rearrangement while the second corresponded to a reductive elimination/oxidative addition

(1) Collman, J. P.; Hegedus, L. S. *Principles and Applications of Organotransition Metal Chemistry*; University Science Books: Mill Valley, CA, 1980; Chapter 6. Cotton, F. A.; Wilkinson, G. *Advanced Inorganic Chemistry*, 4th ed.; Wiley-Interscience: New York, 1980; Chapter 30.

(2) Parshall, G. W. *Homogeneous Catalysis*; Wiley/Interscience: New York, NY, 1980; Chapters 3 and 5 and references therein. Atwood, J. D. *Inorganic and Organometallic Reaction Mechanisms*; Brooks/Cole Publishing: Monterey, CA, 1985; Chapter 5 and references therein.

(3) Vaska, L. *Acc. Chem. Res.* **1968**, *1*, 335.

(4) (a) Chock, P. B.; Halpern, J. *J. Am. Chem. Soc.* **1966**, *88*, 3511. (b) Vaska, L.; Werneke, M. F. *Trans. N.Y. Acad. Sci.* **1971**, *33*, 70. (c) Ugo, R.; Pasini, A.; Fusi, A.; Cenini, S. *J. Am. Chem. Soc.* **1972**, *94*, 7364. (d) Strohmeier, W.; Onada, T. *Z. Naturforsch.* **1968**, *23b*, 1527. (e) Strohmeier, W.; Muller, F. J. *Z. Naturforsch.* **1969**, *24b*, 931. (f) Deeming, A. J.; Shaw, B. L. *J. Chem. Soc. A* **1966**, 1128.

(5) Two reviews cite the pertinent literature in addition to those given in ref 4: (a) James, B. R. *Homogeneous Hydrogenation*; Wiley: New York, 1973; Chapter 12. (b) James, B. R. In *Comprehensive Organometallic Chemistry*; Wilkinson, G., Stone, F. G. A., Abel, E. W., Eds.; Pergamon Press: New York, 1982, Vol. 8; Chapter 51.

(6) (a) Johnson, C. E.; Eisenberg, R. *J. Am. Chem. Soc.* **1985**, *107*, 3148. (b) Johnson, C. E.; Fisher, B. J.; Eisenberg, R. *J. Am. Chem. Soc.* **1983**, *105*, 7772.

(7) Johnson, C. E.; Eisenberg, R. *J. Am. Chem. Soc.* **1985**, *107*, 6531.

<sup>†</sup> Present address: Chemistry Division, Research Department, Naval Weapons Center, China Lake, CA 93555.

sequence with the formation of **2** as an intermediate. We favored this latter pathway, i.e.,  $3 \rightleftharpoons 2 + \text{H}_2 \rightarrow 4$ , principally because reductive elimination of  $\text{H}_2$  from **3** occurs much more rapidly than isomerization.

The clean first-order kinetics for the isomerization, however, were observed only in acetone solvent, and under an excess of hydrogen. When the reaction was studied in benzene, the isomerization proceeded much more rapidly with an apparent half-life of ca. 2 h at 25 °C, although the kinetics were not found to be reproducible.<sup>6a</sup> The isomerization of **3** to **4** was also found to be inhibited by  $[\text{TBA}]\text{Br}$  and accelerated by added  $\text{AgBF}_4$  in benzene and by added  $\text{O}_2$  in acetone. Perhaps most puzzling was the observation that isomerization proceeded more rapidly in rigorously deoxygenated acetone when less  $\text{H}_2$  was present. Since the proposed reductive elimination/oxidative addition sequence for isomerization possessed no kinetic dependence on  $\text{H}_2$ , our observation suggested that another mechanism for isomerization existed. We have therefore reinvestigated the isomerization of the kinetic isomer **3** to the thermodynamic isomer **4** as a function of  $\text{H}_2$  pressure.

In this paper we describe in detail that investigation, including the observation that a second isomerization mechanism involving dihydride transfer between metal centers competes with the first-order isomerization mechanism at ambient temperature and is the sole mechanism operating at temperatures below -20 °C.

### Experimental Section

All kinetic experiments were carried out in resealable 5-mm NMR tubes fitted with a Teflon valve purchased from Trillium Glass.  $\text{H}_2$  was used as received (Air Products C.P., 99.3%), and acetone- $d_6$  (Aldrich Gold Label) was distilled from 4A molecular sieves.  $^1\text{H}$  NMR spectra were recorded on a Bruker WH-400 spectrometer at 400.13 MHz. The temperature of the probe was regulated with a Bruker BVT-1000 temperature control unit.

The complex  $\text{IrBr}(\text{CO})(\text{dppe})$  was synthesized following the procedure previously reported.<sup>6b</sup>

**General Procedure for Sample Preparation.** A stock solution of  $\text{IrBr}(\text{CO})(\text{dppe})$  ( $9.55 \times 10^{-3}$  M, 0.033 g of complex in 5 mL solvent) was prepared in acetone- $d_6$  and stored under  $\text{N}_2$  in a drybox. For each experiment,  $0.50 \pm 0.02$  mL of the stock solution was transferred to the NMR tube which was then connected to a high vacuum line containing an  $\text{H}_2$  inlet. After 3 freeze-pump-thaw degas cycles, the solution was maintained at 0 °C in an ice bath while the sample was placed under the desired pressure of  $\text{H}_2$  by opening the valve at the top of the NMR tube. The sample was then shaken thoroughly to ensure mixing of  $\text{H}_2$  and placed in the thermostated probe of the NMR spectrometer. The total volume of the NMR tube was determined to be  $2.00 \pm 0.05$  mL with a solution volume for each run of  $0.50 \pm 0.05$  mL.

### Results and Discussion

The kinetics of the isomerization reaction of the cis dihydrides of formula  $\text{IrH}_2\text{Br}(\text{CO})(\text{dppe})$  has been studied over a wide range of  $\text{H}_2$  pressures, from 12 to 670 mm of added  $\text{H}_2$ . The reactions were monitored by  $^1\text{H}$  NMR spectroscopy, using the integrals of the hydride resonance of isomers **3** and **4** to determine the relative amounts of each isomer present. Through comparison of the integral of the entire hydride region to the integral of the entire methylene region, the amount of unreacted  $\text{IrBr}(\text{CO})(\text{dppe})$  was determined. For each NMR tube experiment, 0.5 mL of a  $9.55 \times 10^{-3}$  M stock solution of  $\text{IrBr}(\text{CO})(\text{dppe})$ , prepared and stored under nitrogen, was used.

**Isomerization under 670 mm of Hydrogen.** The kinetic results of the isomerization of **3** to **4** under 670 mm of added  $\text{H}_2$  reveal that the reaction proceeds by a clean first-order process. At 28 °C, the half-life for isomerization is 30 h, and the corresponding  $k_{\text{obsd}}$  is  $6.42 \times 10^{-6} \text{ s}^{-1}$ . A plot of  $\ln [3]$  vs. time is linear, as shown in Figure 1, essentially confirming the earlier results of Johnson and Eisenberg.<sup>6a</sup> As discussed in the introduction, the isomerization mechanism favored by us previously was a reductive elimination/oxidative addition sequence shown as eq 3 based on the fact that the initial oxidative addition was found to be fast and reversible. The rate law for this mechanism, given as eq 4, depends only on the concentration of the kinetic dihydride **3** and shows no dependence on hydrogen pressure. Since the initial oxidative addition is highly stereoselective,  $k_{-1}$  is much greater




Figure 1. First-order plot for the isomerization of  $\text{IrH}_2\text{Br}(\text{CO})(\text{dppe})$  under 670 mm of added  $\text{H}_2$ .





Figure 2. First-order plot for the isomerization of  $\text{IrH}_2\text{Br}(\text{CO})(\text{dppe})$  under 200 mm of added  $\text{H}_2$ .

than  $k_2$  and the rate law (eq 4) corresponds to that of a system with an equilibrium preceding the rate-determining step.




$$\frac{-d[3]}{dt} = k_{\text{obsd}}[3] = \frac{k_1 k_2 [3]}{k_{-1} + k_2} \sim \frac{k_1 k_2 [3]}{k_{-1}} \quad (4)$$

**Isomerization under 200–450 mm of Hydrogen.** The kinetics of isomerization for three experimental runs under 200, 300, and 450 mm of added  $\text{H}_2$  were found to be approximately first order. That is, plots of  $\ln [3]$  vs. time are linear for at least 2 half-lives, although they show a slight deviation from linearity at early reaction times. This deviation is most evident at the lowest of these pressures of  $\text{H}_2$  as shown in Figure 2. A more significant, and initially more puzzling, aspect of the kinetic runs under these pressures was that the rate of isomerization was observed to be

**Table I.** Kinetic Data for the Isomerization of 3 to 4 under High H<sub>2</sub> Pressure

pressure of H <sub>2</sub> , mm	total equiv of added H <sub>2</sub> <sup>a</sup>	equiv of H <sub>2</sub> in soln <sup>b</sup>	1st-order rate constant (×10 <sup>6</sup> ), s <sup>-1</sup>
200	3.36	1.07	11.32
300	5.03	1.11	7.70
450	7.55	1.18	6.87
670	11.24	1.27	6.42

<sup>a</sup> Calculated from  $P_0(\text{H}_2)V_{\text{gas}}/RT(9.55 \times 10^{-3} \text{ M})V_{\text{sol}}$  where  $V_{\text{gas}}$  and  $V_{\text{sol}}$  are the volumes of the gas and solution phases, respectively, of the NMR tube and  $9.55 \times 10^{-3} \text{ M}$  is the concentration of the starting iridium complex in solution. <sup>b</sup> Based on the measured concentrations of H<sub>2</sub>, 3, and total dppe species as described in the text. The value shown is given by the expression  $([\text{H}_2] + [3])/([2] + [3]) \sim ([\text{H}_2] + [3])/[3]$ .

**Figure 3.** Second-order plot,  $1/[\text{Ir}_T]$  vs. time, for the isomerization of  $\text{IrH}_2\text{Br}(\text{CO})(\text{dppe})$  under 22 mm of added H<sub>2</sub>.

faster as the pressure of added H<sub>2</sub> was lowered, as shown in Table I. This variation in rate with H<sub>2</sub> pressure was inconsistent with the reductive elimination/oxidative addition sequence of (3) and its rate law, (4), which shows no [H<sub>2</sub>] dependence. A plot of  $k_{\text{obsd}}$  vs.  $1/[\text{H}_2]$  suggested that a second isomerization pathway was operating in addition to (3), while showing that the inverse dependence of [H<sub>2</sub>] for this pathway was not strictly linear.

**Isomerization under Low Pressures of Hydrogen.** Four experimental runs were carried out under 12, 22, 29, and 41 mm of added H<sub>2</sub>, all of which correspond to amounts of added hydrogen less than 1 equiv of starting complex,  $\text{IrBr}(\text{CO})(\text{dppe})$ . These reactions proceeded much more quickly than those under higher pressures of H<sub>2</sub>—typically, isomerizations were complete in less than 15 h. Attempts were made to fit the data to a first-order equation, but plots of  $\ln [3]$  vs. time showed significant deviations from linearity. Clearly, the isomerization path which was predominant at low pressures of H<sub>2</sub> did not follow first-order kinetics.

A second-order treatment of these experimental data was more successful in that plots of  $1/[\text{Ir}_T]$  vs. time were linear, where  $\text{Ir}_T$  represents the sum of unreacted and isomerized iridium complexes, 2 + 3. A plot of these data for the run under 22 mm of added H<sub>2</sub> is shown in Figure 3. Contrary to expectations, the data in Table II show that the observed second-order rate constant,  $k_{\text{obsd}}$ , decreases with decreasing [H<sub>2</sub>], but a plot of  $k_{\text{obsd}}$  vs. [H<sub>2</sub>] was found to be distinctly nonlinear. Surprisingly, a linear correlation was obtained when a plot of  $k_{\text{obsd}}$  vs.  $1/[\text{H}_2]$  was constructed as shown in Figure 4. The origin of this linear dependence on  $1/[\text{H}_2]$  will become apparent below.


We thus conclude that an isomerization mechanism which is second order in complex predominates under low pressures of H<sub>2</sub> and possesses an inverse [H<sub>2</sub>] dependence.

**Mechanism for the Second-Order Isomerization Pathway.** Under conditions in which less than 1 equiv of H<sub>2</sub> is added to the


**Table II.** Kinetic Data for the Isomerization of 3 to 4 under Low H<sub>2</sub> Pressure

initial press of H <sub>2</sub> , mm	initial ratio of [3]/[Ir <sub>T</sub> ] <sup>a</sup>	2nd-order rate constant (×10 <sup>3</sup> ), M <sup>-1</sup> s <sup>-1</sup>
12	0.43	3.52
22	0.61	6.63
29	0.76	7.42
41	0.92	8.23


<sup>a</sup> Based on the measured concentrations of 3 and total dppe species as described in the text.

**Figure 4.** Plot of the observed second-order rate constants,  $k_{\text{obsd}}$  vs.  $1/P_0(\text{H}_2)$  for the isomerization of  $\text{IrH}_2\text{Br}(\text{CO})(\text{dppe})$  with less than 1 equiv of added H<sub>2</sub>.

reaction system, both  $\text{IrBr}(\text{CO})(\text{dppe})$  and the kinetic isomer of  $\text{IrH}_2\text{Br}(\text{CO})(\text{dppe})$ , 3, are present in observable concentrations. We propose a bimolecular mechanism involving these two species to explain the isomerization process under these conditions. This mechanism, which is consistent with the kinetic data, involves dihydride transfer between Ir species via a binuclear intermediate, 5, as shown in eq 5.




The proposal of a dihydride-bridged binuclear intermediate has precedents in other, closely related studies. In the investigation of the stereoselective oxidative addition of H<sub>2</sub> to various Ir(I) complexes of type 2, Johnson and Eisenberg described chemistry involving the reactive intermediate  $\text{IrH}(\text{CO})(\text{dppe})$  generated by dehydrohalogenation of  $\text{IrH}_2\text{Br}(\text{CO})(\text{dppe})$ , 4.<sup>6a</sup> When the reaction was carried out with the base DBU (1,8-diazabicyclo[5.4.0]undec-7-ene) under H<sub>2</sub>, the products were the trihydride  $\text{IrH}_3(\text{CO})(\text{dppe})$ , 6, and the kinetic dihydride 3, as shown in eq 6. Use of D<sub>2</sub> showed that whereas most of 6 corresponding to 6-d<sub>2</sub> formed by the oxidative addition of D<sub>2</sub> to  $\text{IrH}(\text{CO})(\text{dppe})$ , an amount of undeuterated 6 roughly equivalent to the amount of 3-d<sub>2</sub> was produced via a different pathway. Since the thermodynamic dihydride 4 does not reductively eliminate H<sub>2</sub> on the time scale of the experiment, the formation of  $\text{IrBr}(\text{CO})(\text{dppe})$  which gives 3-d<sub>2</sub> upon reaction with D<sub>2</sub> was proposed to occur by




**Figure 5.** Second-order plot,  $1/[\text{Ir}_T]$  vs. time, for the isomerization of  $\text{IrH}_2\text{Br}(\text{CO})(\text{dppe})$  under 120 mm of added  $\text{H}_2$ .


direct dihydride transfer from  $\text{IrH}_2\text{Br}(\text{CO})(\text{dppe})$  to  $\text{IrH}(\text{CO})(\text{dppe})$  through a bridged intermediate with concomitant formation of  $\text{IrH}_3(\text{CO})(\text{dppe})$ .



In an earlier report, Drouin and Harrod<sup>8</sup> proposed a dihydride-bridged dimer species in their attempts to convert the monodentate phosphine complex  $\text{IrH}_3(\text{CO})(\text{P}(p\text{-ClC}_6\text{H}_5)_3)_2$ , **7**, to the unsaturated complex  $\text{IrH}(\text{CO})(\text{P}(p\text{-ClC}_6\text{H}_5)_3)_2$ , **8**. When the trihydride **7** was placed under a stream of nitrogen to displace  $\text{H}_2$ ,  $^1\text{H}$  NMR evidence revealed the occurrence of an equilibrium proposed to involve **9** as shown in eq 7.



Recently Jones and Maguire described the direct intermolecular transfer of  $\text{H}_2$  between a rhenium complex and  $\text{IrBr}(\text{CO})(\text{dppe})$ .<sup>9</sup> Treatment of  $(\eta^4\text{-C}_5\text{H}_6)\text{Re}(\text{PPh}_3)_2\text{H}_3$ , **10**, with 1 equiv of  $\text{IrBr}(\text{CO})(\text{dppe})$ , **2**, resulted in the formation of  $(\eta^5\text{-C}_5\text{H}_5)\text{Re}(\text{PPh}_3)_2\text{H}_2$  and the thermodynamic dihydride isomer of  $\text{IrH}_2\text{Br}(\text{CO})(\text{dppe})$ , **4**, as shown in eq 8. If the reaction had proceeded by elimination of  $\text{H}_2$  from **10** followed by  $\text{H}_2$  oxidative addition of **2**, then the kinetic isomer **3** would have been formed. A control experiment ruled out this possibility, and the results thus strongly support the notion that the reaction between **10** and **2** goes via a dihydride-bridged binuclear intermediate.



**Rate Law for the Bimolecular Isomerization Pathway.** The rate law for the bimolecular isomerization mechanism shown above in eq 5 is derived as follows beginning with

$$\text{rate via bimolecular path} = k_3[\mathbf{3}][\mathbf{2}] \quad (9)$$

The concentration of **2** and **3** are related by an equilibrium constant expression where  $[\text{H}_2]$  corresponds to the concentration of dissolved  $\text{H}_2$ . Therefore

$$\frac{[\mathbf{3}]}{[\mathbf{2}][\text{H}_2]} = \frac{k_{-1}}{k_1} = K_{\text{eq}} \quad (10)$$

and

$$[\mathbf{3}] = K_{\text{eq}}[\mathbf{2}][\text{H}_2] \quad (11)$$

Substituting for **3** into eq 9 yields

$$\text{rate via bimolecular path} = k_3K_{\text{eq}}[\mathbf{2}]^2[\text{H}_2] \quad (12)$$

We next express the rate in terms of  $[\text{Ir}_T]$  corresponding to the sum of unreacted and unisomerized Ir species, **2** + **3**:

$$[\text{Ir}_T] = [\mathbf{2}] + [\mathbf{3}] = [\mathbf{2}](1 + K_{\text{eq}}[\text{H}_2]) \quad (13)$$

$$[\mathbf{2}] = \frac{[\text{Ir}_T]}{(1 + K_{\text{eq}}[\text{H}_2])} \quad (14)$$

Upon substitution of this expression for **2** into eq 12 we obtain

$$\text{rate via bimolecular path} = \frac{k_3K_{\text{eq}}[\text{Ir}_T]^2[\text{H}_2]}{(1 + K_{\text{eq}}[\text{H}_2])^2} \quad (15)$$

The  $[\text{H}_2]$  dependence in rate expression (15) is complex, but it is evident that the value of  $K_{\text{eq}}[\text{H}_2]$  determines the observed hydrogen dependence of the "second-order" or bimolecular pathway. Two limiting regimes can be envisioned, which are as follows: (a) For  $1 \gg K_{\text{eq}}[\text{H}_2]$ , the bimolecular rate =  $k_3K_{\text{eq}}[\text{Ir}_T]^2[\text{H}_2]$ . In this limit, the hydrogen concentration is extremely low, and the rate is proportional to  $[\text{H}_2]$ . (b) For  $K_{\text{eq}}[\text{H}_2] \gg 1$ , the bimolecular rate =  $k_3K_{\text{eq}}[\text{Ir}_T]^2/[\text{H}_2]$ . When the hydrogen concentration is higher, the bimolecular rate becomes proportional to  $1/[\text{H}_2]$ .

The fact that the observed second-order rate constant for isomerization shows an inverse  $[\text{H}_2]$  dependence, as illustrated by the data in Table II and the plot of Figure 4, indicates that even at the low  $\text{H}_2$  pressures used in the present isomerization study,  $K_{\text{eq}}[\text{H}_2] \gg 1$ .

**Isomerization under an Intermediate Pressure of  $\text{H}_2$ .** For an examination of the isomerization in between the high and low  $\text{H}_2$  pressure regimes, an experiment was carried out under 120 mm of added  $\text{H}_2$ . On the basis of  $^1\text{H}$  NMR integrations of the kinetic dihydride **3** and dissolved  $\text{H}_2$ , this initial pressure of  $\text{H}_2$  corresponds to 1.03 equiv of  $\text{H}_2$  in solution. As in the high  $\text{H}_2$  pressure regime, no  $\text{IrBr}(\text{CO})(\text{dppe})$  is observed in solution. The reaction is half complete in 13.5 h, and surprisingly the best fit of the kinetic data is obtained when the reaction is treated as a second-order, bimolecular process. That is, a reasonably straight line results from a plot of  $1/[\text{Ir}_T]$  vs. time, Figure 5, where  $[\text{Ir}_T]$  in this case represents the concentration of the kinetic dihydride **3**.

**The Complete Rate Law.** The kinetic results described above indicate that both isomerization mechanisms operate to differing extents over the range of  $\text{H}_2$  pressures examined. For a system with parallel reaction paths, the complete rate expression is given by the sum of the component rate laws. For the isomerization of  $\text{IrH}_2\text{Br}(\text{CO})(\text{dppe})$ , the complete rate expression is given by eq 16 in which there are terms to account for both the first- and second-order components. If eq 16 is correct, then it should be possible to fit the experimental data to this equation.

$$\text{observed rate} = \frac{-d[\mathbf{3}]}{dt} = \frac{k_2[\mathbf{3}]}{K_{\text{eq}}} + \frac{k_3K_{\text{eq}}[\text{Ir}_T]^2[\text{H}_2]}{(1 + K_{\text{eq}}[\text{H}_2])^2} \quad (16)$$

In order to do this, we analyze the kinetic data in terms of initial rates corresponding to  $(-d[\mathbf{3}]/dt)_{\text{init}}$  since this will give a rate value in  $\text{M}\cdot\text{s}^{-1}$  for each run, unbiased by our perception of whether it is primarily in the first-order or second-order regime. In the initial rate limit, the value of  $[\text{Ir}_T]$  corresponding to **2** + **3** is the same

(8) Drouin, M.; Harrod, J. F. *Inorg. Chem.* **1983**, *22*, 999.

(9) Jones, W. D.; Maguire, J. A. *J. Am. Chem. Soc.* **1985**, *107*, 4544.

**Table III.** Initial Rate Data for the Isomerization of **3** to **4** for all H<sub>2</sub> Pressures

$P_0(\text{H}_2)$ , mm	$[\text{H}_2]_{\text{soln}}$ ( $\times 10^2$ ), M	$[\text{H}_2]/(1 + K_{\text{eq}}[\text{H}_2])^2$ ( $\times 10^7$ ), M	$[\text{3}]$ ( $\times 10^3$ ), M	initial rate ( $\times 10^8$ ), M·s <sup>-1</sup>	reduced rate ( $\times 10^8$ ), M·s <sup>-1</sup>
12	1.02	61.1	4.10	42.0	39.7
22	2.47	86.2	5.83	59.8	56.7
29	4.49	88.1	7.26	53.7	49.7
41	11.1	66.6	8.79	47.5	42.7
120	48.9	22.7	9.55	18.0	12.7
200	81.5	14.4	9.55	14.4	9.1
300	122.0	9.87	9.55	9.6	4.4
450	183.0	6.70	9.55	6.6	1.4
670	272.0	4.56	9.55	6.1	0.9

for each experiment ( $9.55 \times 10^{-3}$  M), whereas the kinetic isomer concentration,  $[\text{3}]$ , is measured throughout the course of each run. Initial rates for all kinetic runs were determined by fitting the experimental data ( $[\text{3}]$  vs. time) to a fifth-order polynomial and extrapolating to  $t \rightarrow 0$ . This allows us to ignore the concentration of the thermodynamic isomer **4**. As seen in Table III, the initial rate first increases with increasing  $[\text{H}_2]$  and then turns over, decreasing as  $[\text{H}_2]$  continues to increase. This is consistent with the functional dependence of  $[\text{H}_2]$  in eq 16 and its two limiting cases described above.

The concentration of H<sub>2</sub> in solution,  $[\text{H}_2]$ , for each experiment is calculated with use of the material balance expression, eq 17, where  $P_0(\text{H}_2)$  is the initial added pressure of H<sub>2</sub> in atmospheres,  $V_{\text{gas}}$  is the volume above the solution in the NMR tube ( $1.5 \pm 0.05$  mL), and  $V_{\text{sol}}$  is the solution volume (0.5 mL):

$$\frac{P_0(\text{H}_2)V_{\text{gas}}}{RT} = \frac{P(\text{H}_2)V_{\text{gas}}}{RT} + [\text{H}_2]V_{\text{sol}} + [\text{3}]V_{\text{sol}} \quad (17)$$

The total amount of added hydrogen,  $P_0(\text{H}_2)V_{\text{gas}}/RT$ , is distributed as the amount in solution,  $[\text{H}_2]V_{\text{sol}}$ , the amount above the solution,  $P(\text{H}_2)V_{\text{gas}}/RT$ , and the amount which is consumed to make dihydride,  $[\text{3}]V_{\text{sol}}$ . Substitutions are made for  $P(\text{H}_2)$  with use of a modified form of Henry's law (i.e.,  $P(\text{H}_2) = [\text{H}_2]/K_h$ ) and for  $[\text{3}]$  recalling the equilibrium of eq 11:

$$\frac{P_0(\text{H}_2)V_{\text{gas}}}{RT} = \frac{[\text{H}_2]V_{\text{gas}}}{RTK_h} + [\text{H}_2]V_{\text{sol}} + K_{\text{eq}}[\text{2}][\text{H}_2]V_{\text{sol}} \quad (18)$$

Substituting  $[\text{Ir}_T] - [\text{3}]$  for  $[\text{2}]$  in (18) and solving for  $[\text{H}_2]$  leads to

$$[\text{H}_2] = \frac{P_0(\text{H}_2)V_{\text{gas}}/RT}{(V_{\text{gas}}/RTK_h) + V_{\text{sol}} + V_{\text{sol}}K_{\text{eq}}([\text{Ir}_T] - [\text{3}])} \quad (19)$$


Values for  $P_0(\text{H}_2)$  are known for each experiment, and a value for Henry's constant,  $K_h$ , was determined experimentally to be  $3.16 \times 10^{-3}$  M/atm by measuring  $[\text{H}_2]$  in acetone-*d*<sub>6</sub> with <sup>1</sup>H NMR spectroscopy as a function of  $P_0(\text{H}_2)$ . Thus the concentration of hydrogen in solution is determined for each kinetic run.

A slight rearrangement of eq 16 yields an expression for a "reduced rate", eq 20, equivalent to subtracting out the contribution of the first-order component from the observed initial rate.

$$\text{reduced rate} = \text{observed initial rate} - \frac{k_2[\text{3}]}{K_{\text{eq}}} = \frac{k_3K_{\text{eq}}[\text{Ir}_T]^2[\text{H}_2]}{(1 + K_{\text{eq}}[\text{H}_2])^2} \quad (20)$$

A plot of reduced rate vs.  $([\text{H}_2]/(1 + K_{\text{eq}}[\text{H}_2])^2)$  should give a straight line with a slope =  $k_3K_{\text{eq}}[\text{Ir}_T]^2$ . As shown in Figure 6, the experimental data fit the derived function reasonably well with two clusters of data points corresponding to the two pressure regimes studied, one with more than 1 equiv of added H<sub>2</sub> and the other with less than 1 equiv. The fit of the line indicates that the derived equation accurately describes the behavior of the system. The data which are plotted, as well as values for  $P_0(\text{H}_2)$ ,  $[\text{3}]$ , and  $[\text{H}_2]$ , are given in Table III.

The very small values for the reduced rate at high  $[\text{H}_2]$  indicate little contribution from the second-order pathway. From runs at the highest added H<sub>2</sub> pressure, we estimate that  $k_{\text{obsd}}$  is approximately  $k_2/K_{\text{eq}}$  and has a value of  $5.5 \times 10^{-6}$  s<sup>-1</sup>. Different values



**Figure 6.** Plot of the reduced rate defined in eq 20 vs. the function describing the  $[\text{H}_2]$  dependence in the complete rate law.

of the equilibrium constant,  $K_{\text{eq}}$ , were employed in plotting the data with the best trial-and-error fit obtained with use of  $2.8 \times 10^4$  M<sup>-1</sup>. On the basis of this value and the fact that the slope of the line in Figure 6 is  $k_3K_{\text{eq}}[\text{Ir}_T]^2$ , we estimate the bimolecular rate constant for isomerization,  $k_3$ , to be  $3.5 \times 10^{-3}$  M<sup>-1</sup> s<sup>-1</sup>.

**Low-Temperature Evidence of the Bimolecular Pathway.** Further confirmation of the bimolecular mechanism for isomerization was demonstrated by a low-temperature experiment which showed that in the H<sub>2</sub> deficient regime isomerization of IrH<sub>2</sub>Br(CO)(dppe) can occur even when the reductive elimination/oxidative addition path is completely shut down. In one NMR tube, a sample of the kinetic dihydride **3** was prepared by addition of 500 mm of H<sub>2</sub> to IrBr(CO)(dppe) and kept below -50 °C to prevent it from isomerizing. The excess H<sub>2</sub> was removed by 2 freeze-pump-thaw cycles, and 200 mm of D<sub>2</sub> were added. After shaking the sample to ensure mixing of D<sub>2</sub>, the tube was maintained at -23 °C for 24 h. A <sup>1</sup>H NMR spectrum taken at -23 °C showed no isomerization to **4** and no incorporation of D<sub>2</sub> into **3** by integration of the hydride resonances relative to the methylene resonances. This confirmed that reductive elimination of H<sub>2</sub> from IrH<sub>2</sub>Br(CO)(dppe) does not occur at -23 °C over a 24-h period. A second NMR sample was prepared by adding 25 mm of H<sub>2</sub> (<1 equiv) to IrBr(CO)(dppe). After this sample was maintained at -23 °C for 24 h, a <sup>1</sup>H NMR spectrum revealed that isomerization had occurred to the extent of 31%. This experiment thus confirmed that isomerization of **3** to **4** can occur independent of the reductive elimination of hydrogen from IrH<sub>2</sub>Br(CO)(dppe) by a bimolecular path.

**Stereoselectivity of Dihydride Transfer.** In light of the stereoselective oxidative addition of H<sub>2</sub> to IrBr(CO)(dppe) to give the kinetic isomer **3**, it is interesting to consider why the dihydride transfer produces the thermodynamic isomer **4**. That is, why does dihydride transfer to **2** proceed with opposite stereoselectivity to that of H<sub>2</sub> oxidative addition? The answer must be electronic in nature since steric factors for the formation of the two isomers of IrH<sub>2</sub>Br(CO)(dppe) by dihydride transfer are similar.

In H<sub>2</sub> oxidative addition, there are two principal interactions between the d<sup>8</sup> metal complex and the H<sub>2</sub> molecule.<sup>10</sup> The first

involves  $\sigma$  donation from the  $\sigma^b(\text{H}_2)$  orbital into a vacant acceptor orbital on the metal center of  $p_z$  or  $p_z-d_z$  hybrid character, while the second is a back-bonding interaction in which electron density is transferred from a filled metal  $d_x$  orbital into the  $\sigma^*$  orbital of  $\text{H}_2$ . In addition to this synergic interaction, a repulsive  $4-e^-$  interaction between the filled  $\sigma^b(\text{H}_2)$  and  $d_z$  orbitals has been invoked as a major contributor to the activation barrier in the  $\text{H}_2$  oxidative addition process.<sup>10c</sup>

The stereoselectivity of  $\text{H}_2$  oxidative addition to **2** arises by a preferred bending of one set of trans ligands in **2** which become cis to each other and trans to the hydride ligands in the product, as shown in A. This preference relates to the  $4-e^-$  repulsive interaction between  $\sigma^b(\text{H}_2)$  and  $d_z$ . As  $\text{H}_2$  approaches the metal complex, one pair of trans ligands bends such that complex + substrate form a trigonal bipyramid as the transition state with the bending ligands and  $\text{H}_2$  occupying the TBP equatorial positions. Preference for bending of the P–Ir–CO axis over the P–Ir–X axis in **2** (i.e., preference for A over B) occurs because this places the better  $\pi$ -acid ligand, CO, in the equatorial plane where it can better stabilize the developing trigonal bipyramid through back-bonding and withdrawal of electron density from  $d_z$ , thereby reducing the repulsive interaction.

In dihydride transfer, the interaction between the square planar Ir(I) complex and the  $\text{MH}_2$  substrate is not a synergic one. If one considers  $\text{MH}_2$  to approach the  $d^8$  complex in a symmetrical manner with equal Ir...H distances, there is no substrate orbital which corresponds to  $\sigma^b(\text{H}_2)$ . Hence, the  $4-e^-$  repulsive interaction does not exist, and the principal reason for addition along the P–Ir–CO axis is removed. In fact, the better  $d_x$  donor orbital in IrBr(CO)(dppe) is the one oriented in the plane defined by P–Ir–Br and the  $z$  axis, and it is in this plane that dihydride transfer occurs. While a detailed theoretical analysis of dihydride transfer remains to be done, we envision that the major orbital interaction takes place between a filled  $d_x$  orbital of Ir(I) and the  $\sigma^*$  function of  $\text{MH}_2$  possessing the same symmetry.

### Conclusions

The kinetic studies which we have described involving the isomerization of the dihydrides of IrH<sub>2</sub>Br(CO)(dppe) show that the reaction proceeds via two different mechanisms. Both mechanisms operate to differing extents throughout the range of  $\text{H}_2$  concentrations examined, but two limiting regimes may be defined as greater than and less than 1 equiv of  $\text{H}_2$  relative to the unsaturated starting complex, IrBr(CO)(dppe).

(10) (a) Saillard, J.-Y.; Hoffmann, R. *J. Am. Chem. Soc.* **1984**, *106*, 2006. (b) Sevin, A. *Nouv. J. Chim.* **1981**, *5*, 233. (c) Dedieu, A.; Strich, A. *Inorg. Chem.* **1979**, *18*, 2940. (d) Noell, J. O.; Hay, P. J. *J. Am. Chem. Soc.* **1982**, *104*, 4578.

In the presence of excess added hydrogen, the isomerization of **3** to **4** occurs primarily by the first-order reductive elimination/ $\text{H}_2$  oxidative addition sequence shown in eq 3. The rate law for this mechanism shows no dependence on  $[\text{H}_2]$ . However, the half-life for isomerization in this regime ( $P_0(\text{H}_2) > 200$  mm) decreases with decreasing  $[\text{H}_2]$ , indicating that the other mechanism which is  $\text{H}_2$  dependent also operates under these conditions.

The reaction path which predominates under hydrogen deficient conditions involves direct dihydride transfer through a binuclear hydride-bridged species. This pathway is dependent on  $[\text{H}_2]$ , is second order with respect to complex, and follows the rate law shown in eq 15.

Through the use of initial rates, the kinetic data have been accommodated into a single rate expression having first- and second-order components. The fit of the data to eq 20 yields an experimental value for the second-order dihydride transfer rate constant,  $k_3$ , of  $3.5 \times 10^{-3} \text{ M}^{-1} \text{ s}^{-1}$ .

While the isomerization of **3** to **4** is thus established to follow two pathways, we note that these results do not explain the accelerated isomerizations which occur when  $\text{AgBF}_4$  is contacted with **3** in benzene or when  $\text{O}_2$  is added to the system in acetone. Additional mechanisms of isomerization, which have not been delineated, are thus possible and indeed followed, under those conditions.

One of the most significant aspects of this study is that the Ir(I) species IrBr(CO)(dppe), **2**, is capable of abstracting dihydrogen very effectively from a metal polyhydride complex by dihydride transfer. While the formation of stable binuclear hydride-bridged complexes is well known,<sup>11</sup> the unusual feature about the present study is that the transfer of  $\text{H}_2$  from one metal center to the other is complete under mild conditions. Since the loss of  $\text{H}_2$  from polyhydride complexes to achieve coordinative unsaturation often requires forcing thermal or photochemical conditions, we think that an alternative approach based on the  $\text{H}_2$ -abstracting ability of **2** might prove attractive. If, indeed, complex **2** does abstract  $\text{H}_2$  from other polyhydride substrates  $L_n\text{MH}_x$ , then dihydride transfer may become an effective method for preparing highly reactive, coordinatively unsaturated species,  $L_n\text{MH}_{x-2}$ , for C–H bond activation studies. This approach is presently under investigation.

**Acknowledgment.** We thank the National Science Foundation (CHE 83-08064 and 86-03055) and the Office of Naval Research for support of this work and Johnson Matthey Co., Inc., for a generous loan of iridium salts. We also thank Ramy Farid for assistance with the figures.

(11) Venanzi, L. M. *Coord. Chem. Rev.* **1982**, *43*, 251.