

**Figure 1.** 270 MHz  $^1\text{H}$  NMR traces of compound **2** (R = OMe, trace a), **1** (R = NH-*i*-Pr, trace b), **3** (R = N(Me)OH, trace c), and **3-Ga** $^{3+}$  (trace d) in  $\text{CDCl}_3$  solution (10 mM), 298 K. Residual signals of **3** in trace d (ca. 20%) indicate slow exchange under these conditions.


[*n*-BuOCH $_2$ CH $_2$ CONHCH(*i*-Bu)CONH-*i*-Pr] showed almost identical chemical shifts for its diastereotopic protons and little solvent dependence ( $\text{CDCl}_3$  versus  $\text{CD}_3\text{OD}$ ). All this indicated restricted conformational freedom due to interchain H bonds in **1**. A priori, H bonds may involve equivalent or nonequivalent amides of adjacent chains. In order to distinguish between these two possibilities compound **2** with only one type of amide group was examined. The IR spectrum of ester **2** showed two NH absorptions (3435 and 3371  $\text{cm}^{-1}$ , 0.65 mM  $\text{CHCl}_3$ ) and its NMR spectrum (Figure 1a) single signals for the diastereotopic protons  $-\text{CH}_2-\text{O}$  and  $-\text{CH}_2-\text{CO}$  in  $\text{CDCl}_3$ , indicating weak H bonds that do not impair the molecule's conformational freedom. The H bonds in amide **1** thus involve nonequivalent amides. This ultimately causes a tilt of the side chain to generate a propellerlike arrangement.<sup>8</sup>

Replacement of the isopropylamine in **1** by methyl hydroxylamine provided the  $\text{Fe}^{3+}$  binder **3**. **3** adopted a H bonded propellerlike conformation like the parent molecule **1** according to its bonded NH (3284  $\text{cm}^{-1}$  in  $\text{CDCl}_3$ ) in the IR spectrum and the nonequivalence of its diastereotopic protons in the NMR (Figure 1c). Titration showed a 1:1 stoichiometry for  $\text{Fe}^{3+}$  binding, and CD ( $\Delta\epsilon = +3.4$  at 450 nm and  $-6.8$  at 365 nm in MeOH) established predominance of the  $\Lambda$ -cis isomer<sup>3</sup> (see Figure 2). This coordination isomer proved to be stabilized by H bonding (NH absorption at 3353  $\text{cm}^{-1}$  in  $\text{CDCl}_3$ ). The  $\text{Fe}^{3+}$  complex of the proline binder **4** which is incapable of forming such H bonds showed significantly smaller Cotton effects ( $\Delta\epsilon = +1.27$  at 465 nm and  $-4.3$  at 378 nm in MeOH).

NMR analysis of the  $\text{Ga}^{3+}$  complex of **3** (which is structurally close to the  $\text{Fe}^{3+}$  complex<sup>9</sup> but amenable to NMR measurement) showed a single set of signals with a pattern close to that of the free ligand (Figure 1d). This establishes the isomeric purity of the complex and its structural similarity to the uncomplexed molecule. In contrast, the  $\text{Ga}^{3+}$  complex of the proline binder **4** showed a complex NMR pattern, indicative of isomeric mixtures.

(8) Tor, Y.; Libman, J.; Shanzer, A.; Felder, C. E.; Lifson, S. *J. Chem. Soc., Chem. Commun.* **1987**, 749-750.

(9) Borgias, B. A.; Barclay, S. J.; Raymond, K. N. *J. Coord. Chem.* **1986**, 15, 109-123.



**Figure 2.** Schematic representation of the  $\Lambda$ -cis  $\text{Fe}^{3+}$  complex of **3**.

These biomimetic tripodlike binders functioned as growth promoters of *Arthrobacter flavescens*.<sup>10</sup> This mutant still contains ferrichrome receptors but lacks the capability to produce ferrichrome. Therefore, its growth depends on externally added siderophores. Although the activity observed for some of these compounds described above was only around 1% of that of ferrichrome, this result is significant and suggestive of specific recognition. No synthetic compound tested so far on this very system, except for synthetic retrohydroxamate ferrichrome, has shown any activity. If and to which extent the chemistry of the natural ferrichromes is governed by some preorientation of conformation that favors certain pathways of complexation, as in the artificial carriers, remains a topic of future research.

**Acknowledgment.** We thank Prof. S. Lifson for stimulating discussions, R. Lazar for skillful technical assistance, and the US-Israel Binational Science Foundation and the National Council for Research and Development for support.

(10) Emery, T., private communication.

## The First [4 + 3] Cycloaddition of a 1,3-Dipole with a 1,3-Diene

Janusz Baran and Herbert Mayr\*

*Institut für Chemie, Medizinische Universität zu Lübeck, Ratzeburger Allee 160, D-2400 Lübeck 1  
Federal Republic of Germany  
Received May 20, 1987*

1,3-Dipolar cycloaddition reactions are usually considered to proceed via concerted [ $\pi 4_s + \pi 2_s$ ] mechanisms<sup>1</sup> though arguments for the stepwise nature have been presented.<sup>2</sup> The nonstereospecific cycloaddition of an electron rich thiocarbonyl ylide with dimethyl dicyanofumarate, an electron deficient dipolarophile, has just been reported as the first unequivocal example of a nonconcerted 1,3-dipolar cycloaddition.<sup>3</sup>

In accordance with the orbital symmetry rules,<sup>4</sup> 1,3-dipoles have been found to react with 1,3-dienes to give vinyl-substituted five-membered rings exclusively.<sup>5</sup> Corresponding [4 + 3] cycloadditions have only been observed in the ozonation of anthracenes<sup>6</sup> and in one case as an intramolecular variant.<sup>7</sup>


(1) (a) Huisgen, R. *Angew. Chem.* **1963**, 75, 742; *Angew. Chem., Int. Ed. Engl.* **1963**, 2, 633. (b) Huisgen, R. In *1,3-Dipolar Cycloaddition Chemistry*; Padwa, A., Ed.; Wiley: New York, 1984; Vol. I, Chapter 1.

(2) Firestone, R. A. *Tetrahedron* **1977**, 33, 3009.

(3) (a) Huisgen, R.; Mloston, G.; Langhals, E. *J. Am. Chem. Soc.* **1986**, 108, 6401. (b) Huisgen, R.; Mloston, G.; Langhals, E. *J. Org. Chem.* **1986**, 51, 4085.


(4) Woodward, R. B.; Hoffmann, R. *Angew. Chem.* **1969**, 81, 797; *Angew. Chem., Int. Ed. Engl.* **1969**, 8, 781.

(5) Crabb, J. N.; Storr, R. C. in ref 1b, Vol. II, p 545.




**Figure 1.** Formation of **3a,b** and **7** from **1** (0.456 mol/L) and **2** (0.453 mol/L) in toluene at 80.0 °C (**5** and **6** could not be determined by the HPLC method used).

### Scheme I




We have reported recently that the steric shielding of the nonterminal positions of the 1,3-diene fragment in **1** is responsible for the unusually high percentage of 1,4-adducts formed from **1** and dihalocarbenes.<sup>8</sup> We report now that the diene **1**, which was prepared by treatment of 1,2-(bisbromomethyl)-3,3,4,4,5,5-hexamethylcyclopentene<sup>9</sup> with magnesium in THF, exhibits 1,4-reactivity also toward a nitrone.

*C,N*-Diphenylnitronone (**2**) and 1,2-bismethylene-3,3,4,4,5,5-hexamethylcyclopentane (**1**) react in benzene at 80 °C to give the diastereomeric [3 + 2] cycloadducts **3a** (28%) and **3b** (4%), the [4 + 3] cycloadduct **7** (18%), and the partially oxidized products **8** (8%) and **9** (2%).<sup>10</sup> Main evidence for the structure of **7** comes from the <sup>13</sup>C NMR spectrum (triplets at  $\delta = 31.46$




**Figure 2.** Schematic energy profiles for concerted and nonconcerted cycloaddition reactions of nitrones with ordinary 1,3-dienes.

### Scheme II



and 74.59) and the catalytic hydrogenation of **7** to give 1-(hydroxymethyl)-3,3,4,4,5,5-hexamethyl-2-(2-phenylethyl)cyclopentene (**10**).



The constant ratio **3a:3b:7**, which was observed at different degrees of conversion, excludes mutual interconversions of these compounds under the conditions of the cycloaddition (Figure 1). This conclusion is in accord with the observation that isolated samples of **3a**, **3b**, and **7** remained unchanged when heated in benzene for 2 days at 80 °C.

As shown in Scheme I, the [3 + 2] cycloadducts **3a,b** may be formed by a concerted cycloaddition reaction or, like the other products, via the intermediate diradical **4**. An alternative pathway for the formation of the [4 + 3] cycloadduct, generation of **12** via [4 + 2] cycloaddition<sup>12</sup> of **1** and **2** and successive Meisenheimer rearrangement<sup>13</sup> to give **7**, has been ruled out by the experiments described in Scheme II. Compound **11** was oxidized to give the amine oxide **12**, which can be assumed to have the same stereochemistry (phenyl groups *trans*) as the potential Diels-Alder adduct of **1** with **2**. When **12** was heated in toluene at 80 °C in presence and absence of **2**, compound **13** was formed, while **7** was not detectable.

If the applicability of the orbital symmetry rules is taken for granted, i.e., if a concerted [<sub>s</sub>4 + <sub>s</sub>4] process cannot account for the formation of **7**, the diradical **4** must be formed as an intermediate. The nonobservance of [4 + 3] cycloadducts in reactions of nitrones with ordinary 1,3-dienes may be explained in two different ways: (a) Nitronone cycloadditions are concerted processes, and only the [3 + 2] cycloaddition is orbital-symmetry allowed (Figure 2a). (b) Nitronone cycloadditions proceed stepwise, but cyclization to five-membered rings is normally preferred (Figure 2b).

If situation (a) were true, the formation of **7** from **1** and **2** would imply that the special steric situation in **1** increased the barrier of the concerted [3 + 2] process to the same height or above the barrier of the stepwise process. In this case, compound **1** should react more slowly than normal dienes. If Figure 2b were true,

(6) (a) Erickson, R. E.; Bailey, P. S.; Davis, J. C., Jr. *Tetrahedron* **1962**, *18*, 388. (b) Bailey, P. S. *Ozonation in Organic Chemistry*; Academic Press: New York, 1982; Vol. II, pp 81.

(7) Böhmer, M.; Heydt, H.; Maas, G.; Gümbel, H.; Regitz, M. *Angew. Chem.* **1985**, *97*, 571; *Angew. Chem., Int. Ed. Engl.* **1985**, *24*, 597.

(8) Mayr, H.; Heigl, U. W. *Angew. Chem.* **1985**, *97*, 567; *Angew. Chem., Int. Ed. Engl.* **1985**, *24*, 579.

(9) Mayr, H.; Will, E.; Heigl, U. W.; Schade, C. *Tetrahedron* **1986**, *42*, 2519.

(10) All new compounds have been completely characterized by <sup>1</sup>H, <sup>13</sup>C NMR, IR, MS, and elemental analysis (see Supplementary Material).


(11) Corresponding intramolecular hydrogen abstractions have been observed in reactions of nitrile oxides with alkynes: Morrocchi, S.; Ricca, A.; Zanarotti, A.; Bianchi, G.; Gandolfi, R.; Grünanger, P. *Tetrahedron Lett.* **1969**, 3329.

(12) Nitrones have previously been postulated to give aminoxides with tetraphenylcyclopentadienone: Brown, C. W.; Marsden, K.; Rogers, M. A. T.; Tylor, C. M. B.; Wright, R. *Proc. Chem. Soc.* **1960**, 254.

(13) Review: Johnstone, R. A. W. In *Mechanisms of Molecular Migrations*; Thyagarajan, B. S., Ed.; Wiley Intersciences: New York, 1969; Vol. II, pp 249.

the steric bulk in **1** might give rise to the formation of the [4 + 3] cycloadduct **7** at the expense of the [3 + 2] cycloadducts without affecting the overall addition rate.

The rate constant for the 1,3-dipolar cycloaddition of **2** with 1-methyl-3-methylene-1-cyclohexene (**14**), a model compound



which reacts at the exocyclic double bond exclusively, is of the same order of magnitude ( $k_2 = 4.4 \times 10^{-5} \text{ L mol}^{-1} \text{ s}^{-1}$ , toluene, 80 °C, determined by HPLC) as the corresponding value reported for the reaction of **2** with ethyl crotonate ( $k_2 = 14 \times 10^{-5}$ ).<sup>14</sup> A slightly smaller value ( $k_2 = 1.1 \times 10^{-5}$ ) is derived for the reaction of **1** with **2** from Figure 1. As a consequence of the discussion related to Figure 2, the close similarity of the reactivities of **1** and **14** implies that in reactions of **2** with normal 1,3-dienes, generally a stepwise mechanism should be accessible, the activation energy of which is at most 2–3 kcal/mol greater than that of the concerted process.

The intermediate diradical **4** profits from allylic resonance, which would not be the case for intermediates formed from nitrones and alkylethylenes. Therefore, the mechanistic considerations on those cycloadditions are not affected by this work.

**Acknowledgment.** We thank the Deutsche Forschungsgemeinschaft and the Fonds der Chemischen Industrie for support of this work.

**Supplementary Material Available:** Analytical data (mp/bp, <sup>1</sup>H NMR, <sup>13</sup>C NMR, IR, mass spectrum) for **3a**, **3b**, and **7–13** (4 pages). Ordering information is given on any current masthead page.

(14) Estimated from  $k_2$  at 100 °C assuming  $\Delta S^\ddagger = -28 \text{ cal mol}^{-1} \text{ K}^{-1}$ ; Huisgen, R.; Seidl, H.; Brünig, I. *Chem. Ber.* **1969**, *102*, 1102.

### Gaseous Negative Ions from Neutral Molecules and Positive Ions: New Information for Neutralization-Reionization Mass Spectrometry

Rong Feng, Chrysostomos Wesdemiotis, and Fred W. McLafferty\*

Chemistry Department, Baker Laboratory  
Cornell University, Ithaca, New York 14853-1301

Received July 6, 1987

Collisional activation of fast gaseous ions can lead to dissociation and/or charge permutation processes.<sup>1,2</sup> Combinations of these are used in neutralization-reionization mass spectrometry (NRMS)<sup>3</sup> to prepare fast gaseous neutrals from mass-selected multikilovolt ions. A further collision can dissociate the neutrals into structurally indicative fragments which are collisionally reionized into positive ions for mass analysis.<sup>4</sup> In this way NRMS has been used recently to study unusual structures such as H<sub>2</sub>C-FH,<sup>4a</sup> H<sub>2</sub>Cl,<sup>4b</sup> and H<sub>3</sub>C(HO)C.<sup>4c</sup> We consider here the alternative reionization into negative ions, which previously has been applied only to D<sub>3</sub><sup>+</sup>, D<sup>+</sup>,<sup>5</sup> H<sub>3</sub><sup>+</sup>,<sup>6a</sup> H<sub>2</sub><sup>+</sup>,<sup>6b</sup> H<sup>+</sup>,<sup>6c</sup> and C<sub>2</sub>H<sub>2</sub><sup>+</sup> molecules. Col-




Figure 1. (A) Hg (70% T)/Xe (70% T), (B) Hg (15% T), and (C) Xe (15% T) NR<sup>-</sup> spectra of CH<sub>2</sub>=CHCH<sub>2</sub><sup>+</sup> (from propene). Absolute abundance of CH<sub>2</sub>=CHCH<sub>2</sub><sup>-</sup> ( $m/z$  41) in percent of the unattenuated CH<sub>2</sub>=CHCH<sub>2</sub><sup>+</sup> main-beam abundance: (A) 0.00012; (B) 0.00095; (C) 0.00038. Hg (70% T) neutralization of CH<sub>2</sub>=CHCH<sub>2</sub><sup>+</sup> produces 1.8% CH<sub>2</sub>=CHCH<sub>2</sub><sup>-</sup>.

lisional charge reversal of positive ion mixtures (not subjected to mass analysis) has been used to prepare negative ions such as Cl<sup>-</sup>, HO<sup>-</sup>, and S<sup>-</sup>.<sup>8</sup> Gaseous anions have also been generated directly by electron impact, flowing afterglow, and chemical ionization techniques.<sup>9</sup> We report here that collisional electron transfer to mass-selected cations or neutrals provides an alternative route to negative ions; the resulting mass spectra contain novel information on the structures and chemistry of the precursor cations and neutrals, as well as the product anions.

**Generation of Negative Ions.** The appropriate multikilovolt precursor ions are generated and mass-selected in the first mass spectrometer (MS-I) of the tandem mass spectrometer described previously<sup>2,3c</sup> to yield neutral molecules by charge-exchange neutralization or dissociation.<sup>3</sup> These fast neutrals are reionized by collision to produce anions which, together with their anionic dissociation products, are measured by scanning MS-II. The spectrum obtained by Hg neutralization of allyl cations in the first collision region followed by deflection of the residual ions and subsequent Xe reionization in the second collision region ("Hg/Xe NR<sup>-</sup> spectrum") is shown in Figure 1A. Alternatively, a single target can effect the two-electron transfer;<sup>5-8</sup> Figure 1B shows the resulting charge-reversal spectrum with Hg ("Hg NR<sup>-</sup> spectrum") in the first collision region at 15% transmittance ( $T$ ), at which the C<sub>3</sub>H<sub>5</sub><sup>-</sup> yield is maximized. This is ~2.5 times greater than the yield from xenon (Figure 1C) under identical conditions. The 15%  $T$  value for maximum efficiency indicates<sup>3c,10</sup> that for

(1) Cooks, R. G., Ed. *Collision Spectroscopy*; Plenum: New York, 1978.

(2) Todd, P. J.; McLafferty, F. W. In *Tandem Mass Spectrometry*; McLafferty, F. W., Ed.; Wiley-Interscience: New York, 1983; pp 149-174.

(3) (a) Danis, P. O.; Wesdemiotis, C.; McLafferty, F. W. *J. Am. Chem. Soc.* **1983**, *105*, 7454-7456. (b) Burgers, P. C.; Holmes, J. L.; Momms, A. A.; Terlouw, J. K. *Chem. Phys. Lett.* **1983**, *102*, 1-3. (c) Danis, P. O.; Feng, R.; McLafferty, F. W. *Anal. Chem.* **1986**, *58*, 348-354. (d) Wesdemiotis, C.; McLafferty, F. W. *Chem. Rev.* **1987**, *87*, 485-500.

(4) (a) Wesdemiotis, C.; Feng, R.; Danis, P. O.; Williams, E. R.; McLafferty, F. W. *J. Am. Chem. Soc.* **1986**, *108*, 5847-5853. (b) Wesdemiotis, C.; Feng, R.; McLafferty, F. W. *Ibid.* **1986**, *108*, 5656-5657. (c) Wesdemiotis, C.; McLafferty, F. W. *Ibid.* **1987**, *109*, 4760-4761.

(5) (a) Cisneros, C.; Alvarez, I.; Barnett, C. F.; Ray, J. A. *Phys. Rev. A* **1976**, *14*, 76-83. (b) Cisneros, C.; Alvarez, I.; Barnett, C. F.; Ray, J. A.; Russek, A. *Ibid.* **1979**, *19*, 631-640.

(6) (a) Devienne, M. F. M. C. R. *Seances Acad. Sci., Ser. B* **1969**, *268*, 1303-1306. (b) Lee, A. R.; Jonathan, P.; Brenton, A. G.; Beynon, J. H. *Int. J. Mass Spectrom. Ion Processes* **1987**, *75*, 329-343. (c) Burgers, P. C.; Terlouw, J. K. *Rapid Commun. Mass Spectrom.* **1987**, *1*, 9-10.

(7) March, R. E.; Young, A. R. *Int. J. Mass Spectrom. Ion Processes* **1987**, *76*, 11-40.

(8) Keough, T.; Beynon, J. H.; Cooks, R. G. *Int. J. Mass Spectrom. Ion Phys.* **1975**, *16*, 417-429; *Chem. Phys.* **1976**, *12*, 191-198.

(9) (a) Melton, C. E. *Principles of Mass Spectrometry and Negative Ions*; Marcel Dekker: New York, 1970. (b) DePuy, C. H.; Bierbaum, V. M. *Acc. Chem. Res.* **1981**, *14*, 146-153. (c) Bowie, J. H. *Mass Spectrom. Rev.* **1984**, *3*, 1-37. (d) Froelicher, S. W.; Freiser, B. S.; Squires, R. R. *J. Am. Chem. Soc.* **1986**, *108*, 2853-2862. (e) Harrison, A. G. *Chemical Ionization Mass Spectrometry*; CRC Press: Boca Raton, FL, 1983.