

bered cobaltacyclic ring, and we will comment later on its electronic structure.¹⁷ The facts of its existence and its ability to take part in catalytic substitution reactions give strong support to the dissociative model for reductively induced electron-transfer catalytic reactions of mononuclear complexes.

Acknowledgment. This research was supported by the National Science Foundation (CHE 86-03728).

Registry No. 1, 51614-82-5; 1⁻, 113353-94-9; 2⁻, 113353-92-7; 2²⁻, 113353-95-0; 3, 113353-93-8; 3⁻, 113378-72-6.

(17) Donovan, B. T.; Rieger, P. H.; Geiger, W. E., work in progress.

Diastereoselective Reactions of Chiral-at-Iron Carbene Complexes C₅H₅(CO)(PR₃)Fe=CHR⁺. Synclinal Isomers Are More Reactive Than Anticinal Isomers

M. Brookhart,* Yumin Liu, and Robert C. Buck

Department of Chemistry
The University of North Carolina
Chapel Hill, North Carolina 27599

Received October 16, 1987

The C₅H₅(NO)(PPh₃)Re- and C₅H₅(CO)(PPh₃)Fe- groups have been used extensively as chiral auxiliaries to carry out diastereo- and enantioselective reactions.¹⁻⁴ Nucleophilic addition at C_a in carbene complexes of the type C₅H₅(NO)(PPh₃)Re=CHR⁺ normally shows high diastereoselection since one face of the carbene ligand is shielded by PPh₃ and barriers to Re=C_a bond rotation are high; consequently, preparation and addition of nucleophiles to a single carbene isomer can be achieved.^{1a-c,5}

We report here a study of addition of simple nucleophiles to chiral-at-iron carbene complexes of the type Cp(CO)(PR₃)Fe=CHR⁺. These studies show that the factors controlling diastereoselectivity are surprisingly complex and involve not only anticinal:synclinal isomer ratios but also the intrinsic differences in reactivity of these isomers, the nature and concentration of the nucleophile, and the ionic strength of the medium.

Iron-carbene complexes 1-5 were prepared by treatment of the α -ether complexes Cp(CO)(L)FeCH(OCH₃)R with TMSOTf

(1) For examples from the Gladysz lab, see: (a) Constable, A. G.; Gladysz, J. A. *J. Organomet. Chem.* 1980, 202, C21. (b) Kiel, W. A.; Lin, G.-Y.; Constable, A. G.; McCormick, F. B.; Strouse, C. E.; Eisenstein, O.; Gladysz, J. A. *J. Am. Chem. Soc.* 1982, 104, 4862. (c) Kiel, W. A.; Lin, G.-Y.; Bodner, G. S.; Gladysz, J. A. *J. Am. Chem. Soc.* 1983, 105, 4958. (d) Crocco, G. L.; Gladysz, J. A. *J. Am. Chem. Soc.* 1985, 107, 4103. (e) O'Connor, E. J.; Kobayashi, M.; Floss, H. G.; Gladysz, J. A. *J. Am. Chem. Soc.* 1987, 109, 4837.


(2) For examples from the Davies lab, see: (a) Baird, G. J.; Davies, S. G. *J. Organomet. Chem.* 1983, 248, C1. (b) Ambler, P. W.; Davies, S. G. *Tetrahedron Lett.* 1985, 26, 2129. (c) Baird, G. J.; Davies, S. G.; Maberly, T. R. *Organometallics* 1984, 3, 1764. (d) Ayscough, A. P.; Davies, S. G. *J. Chem. Soc., Chem. Commun.* 1986, 1648. (e) Davies, S. G.; Dordor-Hedgecock, I. M.; Warner, P. *J. Organomet. Chem.* 1985, 285, 213. (f) Davies, S. G.; Dordor-Hedgecock, I. M.; Warner, P. *Tetrahedron Lett.* 1985, 26, 2125. (g) Brown, S. L.; Davies, S. G.; Warner, P.; Jones, R. H.; Prout, K. *J. Chem. Soc., Chem. Commun.* 1985, 1446. (h) Davies, S. G.; Walker, J. C. *J. Chem. Soc., Chem. Commun.* 1986, 609.

(3) For examples from the Liebeskind lab, see: (a) Liebeskind, L. S.; Welker, M. E.; Feng, R. W. *J. Am. Chem. Soc.* 1986, 108, 6328. (b) Liebeskind, L. S.; Feng, R. W.; Welker, M. E. *Tetrahedron Lett.* 1985, 26, 3075, 3079. (c) Liebeskind, L. S.; Welker, M. E.; Goedken, V. *J. Am. Chem. Soc.* 1984, 106, 441. (d) Liebeskind, L. S.; Welker, M. E. *Organometallics* 1983, 2, 194.

(4) (a) Brookhart, M.; Tucker, J. R.; Husk, G. R. *J. Am. Chem. Soc.* 1981, 103, 979. (b) Brookhart, M.; Tucker, J. R.; Husk, G. R. *J. Am. Chem. Soc.* 1983, 105, 258. (c) Brookhart, M.; Timmers, D.; Tucker, J. R.; Williams, G. D.; Husk, G. R.; Brunner, H.; Hammer, B. *J. Am. Chem. Soc.* 1983, 105, 6721. (d) Buck, R. C.; Brookhart, M. Abstracts 191st National Meeting of the American Chemical Society; New York, New York, April 13-18, 1986; Inorg. Abstract no. 422.

(5) S. G. Davies has reported diastereoselective hydride additions to complexes of the type Cp(CO)(PPh₃)Fe=C(OR)(R').

Scheme I. Free Energy Diagram for the Reactions of Cp(CO)PR₃Fe=CHR⁺ with Nucleophiles



at -78 °C in CH₂Cl₂.⁴ ¹H NMR analysis of CD₂Cl₂ solutions confirmed quantitative generation of 1-5. For ethylidene systems 1-3, the synclinal and anticinal isomers (Scheme I) could be observed by low-temperature ¹H NMR (ca. -114 °C).^{6,7} Equilibrium isomer ratios and ΔG*'s (from line shape analysis) for interconversions are listed in Scheme I. Benzylidene complexes exhibited a single set of resonances which showed no temperature dependence, and we assume a high anticinal:synclinal ratio (>30) based on Cp(NO)(PR₃)Re=CHC₆H₅⁺ as a model.^{1b} The anticinal:synclinal ratio has been independently established as >30:1 for 5.⁸

Reactions of carbene complexes 1-5 with nucleophiles were carried out by quenching into stirred methanol solutions. Large molar excesses (>6 equiv) of nucleophile were present so concentrations changed little during the quench. Two diastereomers were formed in each quench. Results of Gladysz,¹ Davies² and Liebeskind³ clearly establish that the phosphine ligand shields one face of the carbene moiety.⁹ Thus, one diastereomer (S-Nu) should arise from attack on the Si face of the synclinal isomer, while the other (A-Nu) presumably comes from attack on the Re face of the anticinal isomer (Scheme I).

Diastereomer ratios from quenching 1-5 were determined by ¹H NMR analysis of crude products. Complexes 1-A-Nu:1-S-Nu, 2-A-Nu:2-S-Nu, and 3-A-Nu:3-S-Nu were separated by chromatography and structures assigned by ¹H and ¹³C NMR.⁷ In each pair only one diastereomer exhibited a ⁴J_{P-H} (1.3-1.7 Hz); it was assigned the A-Nu configuration based on the P-Fe-C_a-CH₃ "W" geometry and a previous analogous assignment.¹⁰ X-ray structures of 1-S-SC₆H₅ and 3-S-SCOCH₃ confirm these assignments.¹¹ Equilibration of diastereomers of 4-OMe occurs at 25 °C via PPh₃ dissociation; the equilibrium ratio is 70:1.⁸ Isomer 4-A-OMe is expected to be the more stable,^{12,2d} and assignment is made on this basis. Assignments of 5-A-OMe and 5-S-OMe were based on NMR comparisons with 4-A-OMe and

(6) Low-temperature ¹H NMR data of C_a-H: 1 (-114 °C, CD₂Cl₂), 15.9 ppm (synclinal), 17.46 ppm (anticinal); 2 (-114 °C, CD₂Cl₂), 15.69 ppm (synclinal), 17.65 ppm (anticinal); 3 (-126 °C, CD₂Cl₂/SO₂ClF), 17.05 ppm (synclinal), 18.24 ppm (anticinal).

(7) See Supplementary Material for complete spectral and analytical data.

(8) A complete analysis of this diastereomer interconversion has been carried out (Buck, R. C. Ph.D. Dissertation, 1987, University of North Carolina) and will be published elsewhere.

(9) Most of systems studied in ref 1-4 are triphenylphosphine containing systems with the exception of reactions of [(C₅H₅)Re(NO)(PMe₃)-=CHC₆H₅]⁺PF₆⁻ reported by Gladysz.^{1b} The 20:1 product ratio of 1-S-SCOCH₃:1-A-SCOCH₃ in entry 2 of Table I shows that PMe₃ can effectively sterically shield one face of carbene moiety.

(10) Davies, S. G.; Dordor-Hedgecock, I. M.; Sutton, K. H.; Whittaker, M. J. *J. Organomet. Chem.* 1987, 320, C19.

(11) Liu, Y.; Singh, P.; Brookhart, M., unpublished results.

(12) Crocco, G. L.; Gladysz, J. A. *J. Chem. Soc., Chem. Commun.* 1986, 1154.

Table I. Diastereoselective Reactions of $\text{Cp}(\text{CO})\text{PR}_3\text{Fe}=\text{CHR}^+$ with Nucleophiles

entry	carbene	nucleophile	$[\text{Nu}^-]$ (M)	T (°C)	S-Nu:A-Nu	S:A
1	$\text{Cp}(\text{CO})\text{PMe}_3\text{Fe}=\text{CHCH}_3^+$, 1	NaSPh	0.19	-30	4.7:1.0	1.0:2.8 ^h
2	$\text{Cp}(\text{CO})\text{PMe}_3\text{Fe}=\text{CHCH}_3^+$, 1	KSCOCH ₃	0.20	-30	20:1.0	1.0:2.8 ^h
3 ^a	$\text{Cp}(\text{CO})\text{PPh}_3\text{Fe}=\text{CHCH}_3^+$, 3	NaSPh	0.28	-30	4.1:1.0	1.0:2.8 ^h
4	$\text{Cp}(\text{CO})\text{PPh}_3\text{Fe}=\text{CHCH}_3^+$, 3	KSCOCH ₃	0.28	-30	24:1.0	1.0:2.8 ^h
5 ^b	$\text{Cp}(\text{CO})\text{PEt}_3\text{Fe}=\text{CHCH}_3^+$, 2	NaSPh	0.02	-30	6.0:1.0	1.0:5.2 ^h
6	$\text{Cp}(\text{CO})\text{PEt}_3\text{Fe}=\text{CHCH}_3^+$, 2	NaSPh	0.08	-30	2.5:1.0	1.0:5.2 ^h
7	$\text{Cp}(\text{CO})\text{PEt}_3\text{Fe}=\text{CHCH}_3^+$, 2	NaSPh	0.16	-30	1.6:1.0	1.0:5.2 ^h
8	$\text{Cp}(\text{CO})\text{PEt}_3\text{Fe}=\text{CHCH}_3^+$, 2	NaSPh	0.33	-30	1.1:1.0	1.0:5.2 ^h
9 ^c	$\text{Cp}(\text{CO})\text{PEt}_3\text{Fe}=\text{CHCH}_3^+$, 2	NaSPh	0.17	-30	1.4:1.0	1.0:5.2 ^h
10 ^c	$\text{Cp}(\text{CO})\text{PEt}_3\text{Fe}=\text{CHCH}_3^+$, 2	NaSPh/NaBF ₄	0.17/0.55	-30	2.2:1.0	1.0:5.2 ^h
11 ^{d,e}	$\text{Cp}(\text{CO})\text{PEt}_3\text{Fe}=\text{CHCH}_3^+$, 2	NaSPh	0.07	-30	3.1:1.0	1.0:5.2 ^h
12 ^{d,e}	$\text{Cp}(\text{CO})\text{PEt}_3\text{Fe}=\text{CHCH}_3^+$, 2	NaSPh/NaBF ₄	0.07/0.19	-30	4.9:1.0	1.0:5.2 ^h
13 ^f	$\text{Cp}(\text{CO})\text{PEt}_3\text{Fe}=\text{CHCH}_3^+$, 2	KSCOCH ₃	0.04	-30	13:1.0	1.0:5.2 ^h
14 ^g	$\text{Cp}(\text{CO})\text{PEt}_3\text{Fe}=\text{CHCH}_3^+$, 2	KSCOCH ₃	0.08	-30	8.9:1.0	1.0:5.2 ^h
15	$\text{Cp}(\text{CO})\text{PEt}_3\text{Fe}=\text{CHCH}_3^+$, 2	KSCOCH ₃	0.18	-30	3.9:1.0	1.0:5.2 ^h
16	$\text{Cp}(\text{CO})\text{PEt}_3\text{Fe}=\text{CHCH}_3^+$, 2	KSCOCH ₃	0.32	-30	3.0:1.0	1.0:5.2 ^h
17	$\text{Cp}(\text{CO})\text{PPh}_3\text{Fe}=\text{CHC}_6\text{H}_5^+$, 4	NaOCD ₃	0.2	0	1.6:1.0	<1.0:30(est)
18	$\text{Cp}(\text{CO})\text{PPh}_3\text{Fe}=\text{CHC}_6\text{H}_5^+$, 4	NaOCD ₃	1.0	0	1.0:1.1	<1.0:30(est)
19	$\text{Cp}(\text{CO})\text{PPh}_3\text{Fe}=\text{CHC}_6\text{H}_5^+$, 4	NaOCD ₃	2.0	0	1.0:1.9	<1.0:30(est)
20	$\text{Cp}(\text{CO})\text{PPh}_3\text{Fe}=\text{CHC}_6\text{H}_5^+$, 4	NaOCD ₃	4.0	0	1.0:4.2	<1.0:30(est)
21	$\text{Cp}(\text{CO})\text{PEt}_3\text{Fe}=\text{CHC}_6\text{H}_5^+$, 5	NaOCH ₃	0.5	0	1.0:3.0	<1.0:30(est)
22	$\text{Cp}(\text{CO})\text{PEt}_3\text{Fe}=\text{CHC}_6\text{H}_5^+$, 5	NaOCH ₃	1.0	0	1.0:4.0	<1.0:30(est)
23	$\text{Cp}(\text{CO})\text{PEt}_3\text{Fe}=\text{CHC}_6\text{H}_5^+$, 5	NaOCH ₃	2.0	0	1.0:5.0	<1.0:30(est)

^a $[\text{Nu}^-]/\text{TMSOTf}$ = 3.1. ^b $[\text{Nu}^-]/\text{TMSOTf}$ = 3.4. ^c Parallel experiments. ^d $[\text{Nu}^-]/\text{TMSOTf}$ = 4.2. ^e Parallel experiments. ^f $[\text{Nu}^-]/\text{TMSOTf}$ = 2.9. ^g $[\text{Nu}^-]/\text{TMSOTf}$ = 4.0. ^h Isomer equilibrium ratios at -30 °C extrapolated from -104 °C.

4-S-OMe. Control experiments indicate the diastereomers to be stable under reaction conditions and workup procedures.¹³

Table I summarizes our results. Interpretation of diastereomer ratios is most easily discussed in terms of the complete solution of the Curtin–Hammett–Winstein–Holness equation applied to Scheme I.¹⁴

$$\frac{[\text{A-Nu}]}{[\text{S-Nu}]} = K_{\text{eq}} \frac{k_A}{k_S} \frac{k_S[\text{Nu}^-] + k_{SA} + k_{AS}}{k_A[\text{Nu}^-] + k_{SA} + k_{AS}} \quad (1)$$

boundary condition I: $k_A[\text{Nu}^-], k_S[\text{Nu}^-] \gg k_{AS} + k_{SA}$

$$\frac{[\text{A-Nu}]}{[\text{S-Nu}]} = K_{\text{eq}} \quad (2)$$

boundary condition II: $k_{AS}, k_{SA} \gg k_A[\text{Nu}^-], k_S[\text{Nu}^-]$

$$\frac{[\text{A-Nu}]}{[\text{S-Nu}]} = K_{\text{eq}} \frac{k_A}{k_S} \quad (3)$$

It is clear that none of the product ratios represent kinetic quenching (boundary condition I) of the anticlinal:synclinal equilibrium ratios. For the ethylidene systems, the major product always arises from quenching of the minor synclinal isomer. For the benzylidene systems **4** and **5**, the A-Nu/S-Nu ratio varies between 5 and 0.6 and is never as large as the anticlinal:synclinal isomer ratio. These results imply that the synclinal isomers are intrinsically more reactive than the anticlinal isomers (i.e., $k_S > k_A$). If boundary condition II applied to these systems (i.e., isomer interconversion much faster than nucleophile quenching), then the diastereomer ratios should be *independent* of nucleophile concentration and given simply by $(k_A/k_S)K_{\text{eq}}$. This is clearly not the case for **2**, **4**, or **5**. The observed behavior is best rationalized as an intermediate case where the rates of quenching ($k_A[\text{Nu}^-], k_S[\text{Nu}^-]$) are approximately equal to the rates of isomer interconversion (k_{AS}, k_{SA}) as shown in the free energy diagram in Scheme I.¹⁵ Quenching rates are dependent on $[\text{Nu}^-]$, while isomer interconversions are not. Thus, as $[\text{Nu}^-]$ increases boundary condition I is approached, and the fraction of product

arising from the more stable anticlinal isomer will increase.¹⁶ This behavior is observed in each of the three cases and further validates the structural assignments of diastereomers.

A comparison of $\text{C}_6\text{H}_5\text{S}^-$ versus CH_3COS^- at similar concentrations shows that the weaker nucleophile CH_3COS^- is less reactive and thus gives higher S-Nu/A-Nu ratios than the more reactive $\text{C}_6\text{H}_5\text{S}^-$. Extremely reactive nucleophiles should give kinetic or near-kINETIC quenches, and this has been verified in the case of **5** quenched with BD_4^- .⁸ An ionic strength effect is noted in entries 9, 10, 11, and 12. Increasing ionic strength will stabilize the fully charged ions **2-A** and **2-S** relative to partially neutralized transition states **2-A*** and **2-S*** and thus increase ΔG_A^* and ΔG_S^* . As observed, this will lead to higher fractions of synclinal quenching.

The higher intrinsic reactivity of the synclinal isomers relative to the anticlinal isomers can be explained by using the conformational model for $\text{Cp}(\text{CO})(\text{PPh}_3)\text{Fe-CHRR}'$ and $\text{Cp}(\text{NO})(\text{PPh}_3)\text{Re-CHRR}'$ systems.^{12d,17} Attack by Nu^- on the anticlinal isomer results in rehybridization of C_α toward sp^3 and, as shown in transition state **A***, forces R into the least favorable site between CO and PR₃ with initial formation of the high-energy conformation **A'-Nu**. On the other hand, addition of Nu^- to the synclinal isomer places the smallest substituent H between CO and PR₃ and leads directly to the most favorable conformation of S-Nu.

These studies demonstrate that the generalized Curtin–Hammett principle must be invoked in addition of nucleophiles to $\text{Cp}(\text{CO})(\text{PR}_3)\text{Fe=CHR}^+$ complexes. Since nucleophilic attack and isomer interconversion are competitive and since the minor synclinal isomers are intrinsically more reactive than major anticlinal ones, not only may the major products arise from the minor synclinal isomers but also stereoselectivity can be a function of both the strength and concentration of the nucleophile. Clearly this point must be carefully considered in interpreting any diastereoselective reactions of $\text{Cp}(\text{CO})(\text{PR}_3)\text{Fe}$ complexes where isomer interconversion may compete with reactions. For example, enantioselective carbene transfers to alkenes may also proceed mainly via synclinal isomers of $\text{Cp}(\text{CO})(\text{PR}_3)\text{Fe=CHR}^+$ which would suggest cyclopropane ring closure via a backside mechanism.^{4c,18}

(13) For 4-A-OMe, 4-S-OMe quenches were carried out at 0 °C, and samples were immediately cooled to -78 °C prior to analysis to prevent isomer interconversion.

(14) (a) Seeman, J. I. *Chem. Rev.* 1983, 83, 83. (b) Zifirov, N. S. *Tetrahedron* 1977, 33, 2719.

(15) Fitting known rates of interconversion of **2-A** \rightleftharpoons **2-S** and values for $[\text{2-S-SPH}]/[\text{2-A-SPH}]$ at low PhS⁻ concentrations to eq 1 gives approximate values of k_S and k_A of 2.1×10^6 and $4.9 \times 10^4 \text{ s}^{-1} \text{ L-M}^{-1}$ at -30 °C.

(16) Higher ratios of **2-A-SPH**:**2-S-SPH** and **2-A-SCOCH₃**:**2-S-SCOCH₃** were observed when quenching reactions were conducted at lower temperatures.

(17) (a) Seeman, J. I.; Davies, S. G. *J. Chem. Soc., Chem. Commun.* 1984, 1019. (b) Seeman, J. I.; Davies, S. G. *J. Am. Chem. Soc.* 1985, 107, 6522.

(18) Brookhart, M.; Studabaker, W. B. *Chem. Rev.* 1987, 87, 411.

Acknowledgment is made to the National Institutes of Health and the donors of the Petroleum Research Fund, administered by the American Chemical Society, for support of this work. We thank E. L. Eliel for helpful discussions.

Registry No. **1**, S^+ -OTf $^-$, 113109-87-8; **1-A** $^+$ -OTf $^-$, 113214-87-2; **1-S** $^+$ -BF $_4^-$, 113215-02-4; **1-A** $^+$ -BF $_4^-$, 113299-11-9; **1-S-SPh**, 113109-92-5; **1-A-SPh**, 113215-04-6; **1-S-SCOCH₃**, 113109-93-6; **1-A-SCOCH₃**, 113215-07-9; **2**, S^+ -OTf $^-$, 113109-89-0; **2-A** $^+$ -OTf $^-$, 113214-89-4; **2-S** $^+$ -BF $_4^-$, 113215-03-5; **2-A** $^+$ -BF $_4^-$, 113299-12-0; **2-S-SPh**, 113109-96-9; **2-A-SPh**, 113215-06-8; **2-S-SCOCH₃**, 113109-97-0; **2-A-SCOCH₃**, 113215-08-0; **3**, S^+ -OTf $^-$, 113214-91-8; **3-A** $^+$ -OTf $^-$, 113214-93-0; **3-S-SPh**, 113109-94-7; **3-A-SPh**, 113299-13-1; **3-S-SCOCH₃**, 113109-95-8; **3-A-SCOCH₃**, 113215-05-7; **4**, S^+ -OTf $^-$, 113214-95-2; **4-A** $^+$ -OTf $^-$, 113214-97-4; **4-S-OCD₃**, 113109-98-1; **4-A-OCD₃**, 113215-00-2; **4-S-OCH₃**, 113350-82-6; **4-A-OCH₃**, 104832-41-9; **5**, S^+ -OTf $^-$, 113109-91-4; **5-A** $^+$ -OTf $^-$, 113214-99-6; **5-S-OCH₃**, 113109-99-2; **5-A-OCH₃**, 113215-01-3; NaSPh, 930-69-8; KSCOCH₃, 10387-40-3; NaOCD₃, 6552-73-4; NaOCH₃, 124-41-4.

Supplementary Material Available: IR, 1 H, 13 C NMR, and elemental analysis data for **1-S-SPh**, **1-A-SPh**, **1-S-SCOCH₃**, **2-S-SPh**, **2-A-SPh**, **2-S-SCOCH₃**, **2-A-SCOCH₃**, **3-S-SCOCH₃**, **3-A-SCOCH₃**, **5-S-OCH₃**, and **5-A-OCH₃**; IR, 1 H NMR, and elemental analysis data for **1-A-SCOCH₃**; 1 H NMR data for **4-A-OCH₃**, **1**, and **2**; IR, 1 H, and 13 C NMR data for **3-S-SPh** and **3-A-SPh**; 1 H and 13 C NMR data for **4-S-OCH₃** and **5** (12 pages). Ordering information is given on any current masthead page.

Isonitrile Biosynthesis in the Cyanophyte *Hapalosiphon fontinalis*

Volker Bornemann, Gregory M. L. Patterson, and Richard E. Moore*

Department of Chemistry, University of Hawaii
Honolulu, Hawaii 96822

Received January 11, 1988

In 1957 Hagedorn and Tonjes reported the isolation of the first naturally occurring isonitrile, xanthocillin from the fungus *Penicillium notatum*.¹ Since that time isonitriles have been found in other fungi,² bacteria,³ and marine organisms⁴ and more recently in blue-green algae.⁵ The biosynthetic origin of the isonitrile group has intrigued chemists for three decades, but the data obtained to date with fungi, bacteria, and sponges do not present a simple

(1) Hagedorn, I.; Tonjes, H. *Pharmazie* 1957, 12, 567–580; *Chem. Abstr.* 1958, 52, 6362.

(2) (a) Nobuhara, M.; Tazima, H.; Shudo, K.; Itai, A.; Okamoto, T.; Iitaka, Y. *Chem. Pharm. Bull.* 1976, 24, 832–834. (b) Ando, K.; Tamura, G.; Arima, K. *J. Antibiot.* 1968, 21, 587–591. (c) Takatsuki, A.; Suzuki, S.; Ando, K.; Tamura, G.; Arima, K. *Ibid.* 1968, 21, 671–675. (d) Achenbach, H.; Strittmatter, H.; Kohl, W. *Chem. Ber.* 1972, 105, 3061–3066. (e) Marconi, G. G.; Molloy, B. B.; Nagarajan, R.; Martin, J. W.; Deeter, J. B.; Occolowitz, J. L. *J. Antibiot.* 1978, 31, 27–32. (f) Brewer, D.; Taylor, A. J. *Chem. Soc., Chem. Commun.* 1979, 1061–1062. (g) Fujiwara, A.; Okuda, T.; Matsuda, S.; Shiomi, Y.; Miyamoto, C.; Sekini, Y.; Tazoe, M.; Fujiwara, M. *Agric. Biol. Chem.* 1982, 46, 1803–1809.

(3) (a) Marquez, J. A.; Horan, A. C.; Kalyanpur, M.; Lee, B. K.; Loebenberg, D.; Miller, G. H.; Patel, M.; Waitz, J. A. *J. Antibiot.* 1983, 36, 1101–1108. (b) Evans, J. R.; Napier, E. J.; Yates, P. J. *Antibiot.* 1976, 29, 850–852.

(4) See the following reviews: (a) Faulkner, D. J. *Tetrahedron* 1977, 33, 1421–1443. (b) Faulkner, D. J. *Nat. Prod. Rep.* 1984, 1, 552–597. (c) Faulkner, D. J. *Nat. Prod. Rep.* 1986, 3, 1–33.

(5) (a) Moore, R. E.; Cheuk, C.; Patterson, G. M. L. *J. Am. Chem. Soc.* 1984, 106, 6456–6457. (b) Moore, R. E.; Cheuk, C.; Yang, X. G.; Patterson, G. M. L.; Bonjouklian, R.; Smitka, T. A.; Mynderse, J. S.; Foster, R. S.; Jones, N. D.; Swartzendruber, J. K.; Deeter, J. B. *J. Org. Chem.* 1987, 52, 1036–1043. (c) Moore, R. E.; Yang, X. G.; Patterson, G. M. L. *J. Org. Chem.* 1987, 52, 3773–3777. (d) Schwartz, R. E.; Hirsch, C. F.; Springer, J. P.; Pettibone, D. J.; Zink, D. L. *J. Org. Chem.* 1987, 52, 3706–3708.

Table I. Incorporation Experiments for Hapalindole A

precursor	amount fed (μ Ci)	total incorpn (%)	specific incorpn (%)	loss of 14 C upon acid hydrolysis of 1 to 2 (%)
[14 C]cyanide	24.5	0.16	1.21	94.8
[2- 14 C]glycine	45.1	0.33	1.00	99.0
L-[3- 14 C]serine	27.6	0.34	0.87	97.0
[14 C]formate	50.0	0.11	0.26	96.3
L-[methyl- 14 C]methionine	27.3	0.05	0.08	97.6
[1,2- 14 C]acetate	30.5	0.15	0.57	5.7
DL-[methylene- 14 C]tryptophan	49.9	0.13	0.36	0.06




Figure 1. Proton noise-decoupled 13 C NMR spectra of (A) **1** (natural abundance); (B) **1** obtained from feeding [2- 13 C, 15 N]glycine to *H. fontinalis*; (C) **3** (natural abundance); and (D) **3** obtained from hydrolysis of **1** showing spectrum B.

picture. Studies on the biosynthesis of xanthocillin have suggested that L-tyrosine is the primary source of the isonitrile nitrogen,⁶ but C₁ donors linked to tetrahydrofolate metabolism (methionine, formate, C-2 of glycine, and C-3 of serine)⁷ as well as other C₁ donors (e.g., cyanide and carbamoyl phosphate)⁸ are not sources of the isonitrile carbon. Puar et al.⁹ have found, however, that L-[methyl- 13 C]methionine labels the isocyanato group of the haemimicins in the bacterium *Micromonospora echinosporae* var. *challensis*, and Garson¹⁰ has discovered that [14 C]cyanide is incorporated into the isonitrile carbons of the marine sponge metabolite diisocyanoacadiene. Garson's result is most interesting, since sponges frequently possess symbiotic microorganisms and certain bacteria and blue-green algae generate inorganic cyanide from amino acids.¹¹ Our interest in the possible role of symbiotic

(6) (a) Achenbach, H.; König, F. *Chem. Ber.* 1972, 105, 784–793. (b) Achenbach, H.; Grisebach, H. *Z. Naturforsch., B: Anorg. Chem., Org. Chem., Biochem., Biophys. Biol.* 1965, 20, 137–145.

(7) (a) Herbert, R. B.; Mann, J. *Tetrahedron Lett.* 1984, 25, 4263–4266. (b) Herbert, R. B.; Mann, J. *J. Chem. Soc., Chem. Commun.* 1984, 1474–1475.

(8) Cable, K. M.; Herbert, R. B.; Mann, J. *Tetrahedron Lett.* 1987, 28, 3159–3162.

(9) Puar, M. S.; Munayyer, H.; Hedge, V.; Lee, B. K.; Waitz, J. A. *J. Antibiot.* 1985, 38, 530–532.

(10) Garson, M. J. *J. Chem. Soc., Chem. Commun.* 1986, 35–36.