

4.56. Found: C, 50.83; H, 3.05; N, 4.31.) **9** (190 mg, 29%): oil; ^1H NMR (CDCl_3) δ 3.57 (s, 3 H), 6.39 (d, $J = 4$ Hz, 1 H), 7.14 (d, $J = 4$ Hz, 1 H), 7.45–7.53 (m, 3 H), 7.73–7.77 (m, 2 H); ^{13}C NMR (CDCl_3) δ 51.9, 109.4, 114.9, 123.8, 126.3, 128.9, 129.8, 132.5, 133.8, 160.0, 167.1; MS, m/z (relative intensity) 309 (35), 307 (36), 278 (3), 276 (3), 251 (3), 249 (3), 230 (100); precise mass calcd for $\text{C}_{13}\text{H}_{10}\text{BrNO}_3$ 306.9845, found 306.9840.

Reaction of *N*-Benzoyl-(2*R,S*)-alanine Methyl Ester (2a**) with DTBP.** A mixture of *N*-benzoyl-(2*R,S*)-alanine methyl ester (**2a**) (0.3 g, 1.5 mmol) and DTBP (4 mL, 19 mmol) in *tert*-butyl alcohol (30 mL), contained in a quartz tube under nitrogen, was irradiated in the Rayonet photochemical reactor. After 4 days the reaction mixture was concentrated and chromatographed on silica with ethyl acetate-hexane as eluent to give dimethyl 2,3-dibenzamido-2,3-dimethylbutanedioate (**13**) and *N*-benzoyl-2,2-dimethylglycine methyl ester (**12**). **13** (60 mg, 20%): mp 170–177 °C; ^1H NMR ($\text{DMSO}-d_6$) δ 2.00 (s, 6 H), 3.80 (s, 6 H), 6.80 (br, 2 H), 7.53–7.93 (m, 10 H); MS, m/z (relative intensity) 413 (0.4), 381 (2), 353 (7), 231 (22), 207 (38), 175 (8), 105 (100), 77 (50). (Anal. Calcd for $\text{C}_{22}\text{H}_{24}\text{N}_2\text{O}_6$: C, 64.1; H, 5.9; N, 6.8. Found: C, 63.9; H, 5.9; N, 6.6.) **12** (32 mg, 10%) was identical in all respects with an authentic

sample obtained by derivatization of the corresponding amino acid.³⁶

Acknowledgment. We thank Dr. P. J. Steel for helpful discussion and comment and we gratefully acknowledge the assistance of Professor L. C. Leitch in providing us with details of the preparation of 2,5,5-trideuterioprolin. This work was supported by the Australian Research Grants Scheme.

Registry No. **1a**, 1205-08-9; **1b**, 102770-12-7; (2*S*)-**2a**, 38767-73-6; (2*R*)-**2a**, 7260-27-7; (2*RS*)-**2a**, 38767-73-6; (2*S*)-**2b**, 118013-54-0; (2*R*)-**3a**, 1492-13-3; (2*RS*)-**3a**, 14599-03-2; (2*S*)-**3b**, 116297-93-9; **4a**, 71533-21-6; (2*RS*)-**5a**, 54571-66-3; (2*RS*)-**5b**, 117918-31-7; (2*RS*)-**6a**, 114051-14-8; (2*RS*)-**6b**, 117918-32-8; (2*RS*)-**6c**, 117918-33-9; **9**, 117918-26-0; **10**, 117918-27-1; **11**, 117918-28-2; **12**, 65563-98-6; (\pm)-**13** (diastereomer-1), 117918-29-3; (\pm)-**13** (diastereomer-2), 117918-34-0; **14**, 117918-30-6; **15**, 116453-15-7.

(36) Hoefle, G.; Hoehne, G.; Respondek, J.; Schwarz, H. *Org. Magn. Reson.* 1977, 12, 477.

Steric and Electrochemical Effects on Rates of Electron Transfer and $\text{S}_{\text{N}}2$ Reactions of 9-(Dialkylamino)fluorene Ions with Alkyl Halides


F. G. Bordwell* and John A. Harrelson, Jr.

Contribution from the Department of Chemistry, Northwestern University, Evanston, Illinois 60208. Received June 9, 1988

Abstract: Rate ratios for reactions of PhCH_2Cl vs Ph_2CHCl with seven 9-(dialkylamino)fluorene ($9\text{-R}_2\text{NFl}^-$) ions were measured in Me_2SO solution. Although the reaction sites in these ions are known to be highly congested and Ph_2CHCl is more sterically hindered than is PhCH_2Cl , the $k^{\text{PhCH}_2\text{Cl}}/k^{\text{Ph}_2\text{CHCl}}$ rate ratios for reactions with $9\text{-R}_2\text{NFl}^-$ ions in Me_2SO were all much lower (0.20–4.9) than for the less hindered 9-MeFl^- or $p\text{-MeOC}_6\text{H}_4\text{O}^-$ ions (81 and 138, respectively). This suggested that the Ph_2CHCl reactions with $9\text{-R}_2\text{NFl}^-$ ions were occurring by single electron transfer (SET) mechanisms, despite the formation of high yields of $\text{S}_{\text{N}}2$ -type products. This conclusion was supported by the observation of a close correspondence between SET rates ($\log k_{\text{SET}}$), calculated by using the Marcus equation, and $\log k_{\text{obsd}}$ for reactions of $9\text{-R}_2\text{NFl}^-$ ions with both a known single electron acceptor, $\text{F}_3\text{CCH}_2\text{I}$, and with Ph_2CHCl and $(p\text{-ClC}_6\text{H}_4)_2\text{CHCl}$. Similar $\log k_{\text{SET}}$ vs $\log k_{\text{obsd}}$ comparisons for reactions of the $9\text{-R}_2\text{NFl}^-$ ions with PhCH_2Cl , $c\text{-C}_6\text{H}_{11}\text{Br}$, and $n\text{-BuBr}$ revealed greater disparity.

The idea that, in principle, a concerted ("polar") $\text{S}_{\text{N}}2$ reaction can merge with a single electron transfer (SET) mechanism, wherein the product is formed by coupling of a geminate radical pair, has been recognized for many years.¹ In his recent definitive book on electron transfer reactions Ebersson concludes, however, that it takes a very strong electron donor anion to effect a bimolecular aliphatic substitution reaction on an alkyl halide by an outer-sphere SET mechanism.² Nevertheless, he points out that this has been achieved for certain alkyl halides and that there is good reason to believe that this SET mechanism will merge with the concerted single electron shift $\text{S}_{\text{N}}2$ reaction, as has been suggested by several investigators.³ Outer-sphere SET substi-

tutions have been observed by Lund and Lund for reactions of *t*-BuBr with radical anions, $\text{ArH}^{\cdot-}$, and with carbanion **1**, which



was generated electrochemically from 4-(methoxycarbonyl)-*N*-methylpyridinium iodide. Similar substitutions were also observed for reactions of **1** with 1-adamantyl and neopentyl bromides, but the less hindered ethyl, *n*-butyl, and *sec*-butyl bromides appeared to react by borderline mechanisms.³

In earlier papers⁴ we have shown that reactions of 9-substituted fluorene ions, 9-GFl^- , with PhCH_2Cl are subject to rate-retarding steric effects, as G becomes more bulky along the series, Me, Et,

(1) (a) Dessy, R. E.; Pohl, R. L.; King, R. B. *J. Am. Chem. Soc.* 1966, 88, 5121–5124. (b) Dessy, R. E.; Weisman, P. H. *Ibid.* 5124–5129. (c) Bilevitch, K. A.; Pubnov, N. N.; Okhlobystin, O. Yu. *Tetrahedron Lett.* 1968, 3465–3468. (d) Kormer, G. S.; Hall, M. L.; Traylor, T. J. *J. Am. Chem. Soc.* 1972, 94, 7205–7206. (e) Bank, S.; Noyd, D. A. *J. Am. Chem. Soc.* 1973, 95, 8203–8205. (f) Garst, J. F.; Barbas, J. T. *J. Am. Chem. Soc.* 1974, 96, 3239–3249. (g) Zieger, H. E.; Angres, I.; Mathisen, D. *J. Am. Chem. Soc.* 1976, 98, 2580–2585. (h) Russell, G. C.; Jawdoskiuk, M.; Makosza, M. *J. Am. Chem. Soc.* 1979, 101, 2355–2359. (i) Smith, G. F.; Kuivila, H. G.; Simon, R.; Sultan, L. *J. Am. Chem. Soc.* 1981, 103, 833–839. Kuivila, H. G.; Alnajjah, M. S. *Ibid.* 1982, 104, 6146–6147. (j) Ashby, E. C.; DePriest, R. *J. Am. Chem. Soc.* 1982, 104, 6144–6146.

(2) Ebersson, L. *Electron Transfer Reactions in Organic Chemistry*; Springer-Verlag: New York, 1987.

(3) (a) Lund, H.; Kristensen, L. H. *Acta Chem. Scand., Ser. B* 1979, B33, 495–498. (b) Pross, A. *Acc. Chem. Res.* 1985, 18, 212–219. (c) Lund, T.; Lund H. *Acta Chem. Scand. Ser. B* 1986, B40, 470–485.

(4) (a) Bordwell, F. G.; Hughes, D. L. *J. Org. Chem.* 1983, 48, 2206–2215. (b) Bordwell, F. G.; Cheng, J. P.; Seyedrezai, S. E.; Wilson, C. A. *J. Am. Chem. Soc.*, in press.

Table I. Rates of Reactions of 9-Substituted Fluorene Ions, 9-GFl⁻, with Benzyl and Benzhydryl Chlorides

G	r^a	pK_a	$E_{ox}(A^-)^b$	k^{Ph_2CHCl}	$k^{PhCH_2Cl}/k^{Ph_2CHCl}$
Me	(0.0)	22.3	-0.480	0.609	81
PhCH ₂	-0.10	21.4	-0.402	0.30	65
MeO		22.1	-0.574	0.28	51
4-MeC ₆ H ₄		18.3	-0.318	-0.074 ^d	12
Ph	-0.58	17.9	-0.278	0.057 ^d	11
3-ClC ₆ H ₄		16.85	-0.188	0.038 ^d	6.8
4-MeSO ₂ C ₆ H ₄		15.2	-0.097	0.015 ^d	4.3
c-C ₆ H ₁₁		23.0	-0.434	0.130	28
<i>t</i> -Bu	-2.6	24.35	-0.409	0.041	11.5
azetidiny	-0.17	21.8	-0.865	8.19	2.7
pyrrolidiny	-1.2	22.2	-0.805	3.48	0.83
Me ₂ N	-1.5	22.5	-0.660	0.267	4.9
piperidiny	-1.6	22.5	-0.643	0.270	4.8
2-Me-piperidiny		21.5	-0.598	0.062	1.2
2,2,6,6-Me ₄ -piperidiny		18.3	-0.438	0.028	0.12
<i>i</i> -Pr ₂ N	-3.4	20.8	-0.485	0.0349	0.20

^aIn log units. ^bMeasured by cyclic voltammetry in Me₂SO with a Ag/AgI reference electrode; referenced to the aqueous standard hydrogen electrode (SHE) by adding -0.125 V. ^cM⁻¹ s⁻¹; measured spectrophotometrically by the method described previously.^{4a} ^dReference 8a. ^eThe rate data with PhCH₂Cl are from ref 4, 5, 16, and 21.

i-Pr, *t*-Bu, and *i*-Pr₂N, that cause progressive rate decreases over a range of nearly 10⁴. In contrast, a low sensitivity to the steric bulk of G was observed for reactions of 9-GFl⁻ ions with two single-electron acceptors, 1,1-c-C₆H₁₀(NO₂)Ts⁵ and F₃CCH₂I,⁶ in authentic SET reactions. With each of these acceptors a plot of log k_{obsd} vs the oxidation potentials of the anions, $E_{ox}(A^-)$, was linear for reactions where the steric demands of G varied appreciably, e.g., G = Me, MeO, MeS, *t*-Bu, and R₂N. Furthermore, we have presented evidence in a preliminary account that rates of reactions of a series of 9-R₂NFl⁻ ions with F₃CCH₂I or Ph₂CHCl plot linearly with $E_{ox}(A^-)$. This led to the conclusion that both of these latter reactions were occurring by single electron transfer (SET) radical pair mechanisms, F₃CCH₂I reacting to give radical products and Ph₂CHCl reacting to give substitution products.⁷ We now present a more complete account of these results, including an extension of the study to (*p*-ClC₆H₄)₂CHCl, *n*-BuBr, and c-C₆H₁₁Br.

Results and Discussion

Steric Effects in Reactions of 9-GFl⁻ Ions with PhCH₂Cl and Ph₂CHCl. The rates of reactions of 9-R₂NFl⁻ and other 9-GFl⁻ ions with Ph₂CHCl are presented in Table I and are compared with rates of reactions with PhCH₂Cl.

The presence of the second Ph group in Ph₂CHCl should markedly increase steric hindrance to nucleophilic attack, relative to that with PhCH₂Cl. One would expect on steric grounds, therefore, to see large $k^{PhCH_2Cl}/k^{Ph_2CHCl}$ rate ratios in Table I. Indeed, the ratio with 9-MeFl⁻ ion is 81, and a ratio of 138 has been observed for reactions of these chlorides with the less sterically hindered *p*-MeOC₆H₄O⁻ ion.⁸ For the first nine (miscellaneous) groups (G) in Table I the steric retardation factor, r (eq 2), for

$$r = \beta(\Delta pK_{HA}) - \log(k^{Me}/k^G) \quad (2)$$

reactions with PhCH₂Cl has been shown to increase from 0.0, for the reference group Me, to -2.6 (log units) for *t*-Bu (the Brønsted $\beta = 0.31$); for the second series of seven 9-R₂NFl⁻ ions the r values change from -0.17 for azetidiny to -3.4 for *i*-Pr₂N.^{4b} One would expect to see, for each of these series, a general increase in $k^{PhCH_2Cl}/k^{Ph_2CHCl}$ rate ratios and perhaps a precipitous increase

when the r values are large, as for *t*-Bu and *i*-Pr₂N. Instead, there is a general tendency for the ratios to *decrease* for the first series; for the second series all the ratios are small, those for *t*-Bu and *i*-Pr₂N being 11.5 and 0.20, respectively. The reactions of 9-GFl⁻ ions with PhCH₂Cl and Ph₂CHCl gave good to high yields of S_N2-type substitution products in all cases tested.

In an earlier paper the trio of 9-GFl⁻ ions, where G = Me, MeO, Me₂N, which have nearly the same basicity but progressively more negative $E_{ox}(A^-)$ values, was shown to give progressively faster rates with known SET acceptors. The trio was selected for this reason to serve as a test for SET characteristics in reactions with alkyl halides.⁶ The test failed to reveal SET characteristics in reactions of the trio with PhCH₂Cl, however, the rate constants decreasing progressively along the series instead of increasing. Since the size of the rate constant decreases corresponded closely to rate constant decreases along the series G = CH₃, MeCH₂, Me₂CH, it was concluded that the order of rate constants for the S_N2 reactions with PhCH₂Cl was determined by steric effects with no indication of a SET component. One would then expect the rate constant differences for S_N2 reactions of the trio with the more sterically hindered Ph₂CHCl to be exaggerated. Instead, the differences essentially disappear, the relative rates being 2.3, 1.1, and 1.0, respectively.

Evidence for Single Electron Transfer (SET) in Reactions of 9-R₂NFl⁻ Ions with Ph₂CHCl and PhCH₂Cl. In view of the evidence obtained earlier for a relatively low sensitivity to steric effects with single electron acceptors,^{5,6} the results described above and in Table I suggest that PhCH₂Cl and Ph₂CHCl are reacting by different mechanisms. As a first approximation, we assume that for PhCH₂Cl the mechanism is essentially a concerted S_N2 reaction and as such is sensitive to steric effects. For Ph₂CHCl, we assume that the reactions with 9-GFl⁻ ions are occurring by stepwise SET (radical pair) mechanisms that are relatively insensitive to steric effects. (An S_N1 mechanism is ruled out since the rates are first order in nucleophile as well as electrophile.)

The observation that the rates of reactions (k_{obsd}) of SET acceptors with 9-GFl⁻, where G varies in size and structure, plot linearly with their $E_{ox}(A^-)$ values^{5,6} and the results shown in Table I led us to examine reactions for a series of 9-R₂NFl⁻ ions as a possible test for SET in reactions with electrophiles. The 9-R₂NFl⁻ ions in this series do not differ appreciably in basicity ($pK_{HA} = 22 \pm 1.2$) but have oxidation potentials that cover a considerable range (0.427 V or 9.8 kcal/mol). A Marcus-type plot of log k_{obsd} vs $E_{ox}(A^-)$ for a series of 9-R₂NFl⁻ ions reacting with F₃CCH₂I, a known SET acceptor, was indeed found to be reasonably linear,⁷ and better linearity was observed for a plot for 9-R₂NFl⁻ ions reacting with Ph₂CHCl.⁷ The fit was better for the latter plot because the kinetics were better. The rates were monitored in each instance by following the decrease in absorbance of the 9-R₂NFl⁻ ions. For F₃CCH₂I, the ln(abs₀/abs_{*t*}) vs time plots began to deviate from linearity after 1 half-life, probably because of interference from a strongly absorbing product species (9-R₂NFl⁻ or the like). On the other hand, the kinetics for the reactions with Ph₂CHCl remained linear for several half-lives, as is typical of S_N2 reactions of 9-GFl⁻ ions with alkyl halides.

Additional support for a stepwise SET pathway for the reactions of this series of 9-R₂NFl⁻ ions with F₃CCH₂I and Ph₂CHCl was obtained by a comparison of the rate constants observed for these reactions with those calculated for SET by Ebersson's method using the Marcus equation (eq 3).⁹

$$\log k_{SET} = \log k_d - \log \left\{ 1 + 0.2 \exp \left[\frac{\lambda}{4} \left(1 + \frac{\Delta G^\circ}{\lambda} \right)^2 / RT \right] \right\} \quad (3)$$

The value of k_d in eq 3, the diffusion-controlled rate in Me₂SO, is taken as 3.3×10^9 ,¹⁰ $\Delta G^\circ = 23.06[E_{rd} - E_{ox}(A^-)]$, where E_{rd}

(5) Bordwell, F. G.; Bausch, M. J.; Wilson, C. A. *J. Am. Chem. Soc.* **1987**, *109*, 5465-5470.

(6) Bordwell, F. G.; Wilson, C. A. *J. Am. Chem. Soc.* **1987**, *109*, 5470-5474.

(7) Bordwell, F. G.; Harrelson, J. A., Jr. *J. Am. Chem. Soc.* **1987**, *109*, 8112-8113.

(8) (a) Hughes, D. L. Ph.D. Dissertation, Northwestern University, 1981.

(b) Harrelson, J. A., Jr., unpublished results.

(9) (a) Ebersson, L. *Acta Chem. Scand. Ser. B* **1984**, *B38*, 439-459. (b) Ebersson, L. *Adv. Phys. Org. Chem.* **1982**, *18*, 79-185. (c) Ebersson, L. *Acta Chem. Scand. Ser. B* **1982**, *B36*, 533-543.

(10) Gordon, A. J.; Ford, R. A. In *The Chemist's Companion, A Handbook of Practical Data, Techniques, and References*, Wiley: New York, 1972; pp 67 and 137-138.

Table II. Comparison of Rates (k_{obsd}) with Expected Rates of Single Electron Transfer (k_{SET}) for Reactions of 9-(Dialkylamino)fluorene Ions, $9\text{-R}_2\text{NFl}^-$, with $\text{F}_3\text{CCH}_2\text{I}$ and Ph_2CHCl

NR ²	$\text{F}_3\text{CCH}_2\text{I}^b$			Ph_2CHCl^d		
	ΔG_0^a	$\log k_{\text{obsd}}$	$k_{\text{obsd}}/k_{\text{SET}}^c$	ΔG_0^a	$\log k_{\text{obsd}}$	$k_{\text{obsd}}/k_{\text{SET}}^c$
	-6.8	fast		-9.0	+0.91	0.34
	-5.4	fast		-7.6	+0.54	0.53
NMe ₂	-2.1	+0.75	6.0	-4.3	-0.57	0.40
	-1.7	-0.28	8.3	-3.9	-0.57	0.55
	-0.6	-1.43	0.14	-2.9	-1.21	0.28
N(<i>i</i> -Pr) ₂	+2.0	-1.50	1.1	-0.3	-1.45	1.4
	+3.0	-2.25	1.22	+0.8	-1.55	0.45

^aIn kcal/mol. ^b $E_{\text{rd}} = 0.4 \times (1.425) = -0.57$; ²³ $\lambda = 60$. Radical-reaction-type products are formed. ^cCalculated from eq 2. ^d $E_{\text{rd}} = 0.4 \times (1.185) = -0.474$; ²³ $\lambda = 65$. S_N2 products were obtained.

is the reduction potential of the electrophile and $E_{\text{ox}}(\text{A}^-)$ is the oxidation potential of the anion; both are determined for Me₂SO solution and referenced to the aqueous standard hydrogen electrode (SHE_{aq}). The E_{rd} values are taken from Ebersson^{9a} or calculated as described earlier.⁷ The solvent and bond reorganization parameter, λ , is estimated from those for similar reactions reported elsewhere.^{1,9a} Uncertainties of estimated λ and E_{rd} values cause k_{SET} calculations to be accurate to only 1 order of magnitude.

Ebersson has used eq 3 to obtain $k_{\text{obsd}}/k_{\text{SET}}$ ratios for 33 reactions of anionic nucleophiles with organic electron acceptors in order to estimate whether or not stepwise SET is feasible.^{9a} The ratios vary from about 10⁻⁵ to 10²⁸ or more; for 24 of these reactions the ratios were above 10⁵, which places them in the "SET not feasible" class. Several organometallic reagents containing strongly basic carbanion or nitranion moieties were included, but examples of reactions of anions free of counterion effects were absent. Lund has filled this gap by carrying out an extensive study of reactions of aromatic radical anions, ArH^{•-}, with simple alkyl halides and has found good fits for Marcus-type plots of $\log k$ vs $E_{\text{ox}}(\text{A}^-)$. Four ArH^{•-} ions with $E_{\text{ox}}(\text{A}^-)$ values (vs SHE_{aq}) ranging from -0.611 to -1.488 V reacting with *t*-BuBr were found to give progressively greater yields of substitution products as $E_{\text{ox}}(\text{A}^-)$ became more negative.^{3a} Good yields of substitution products were also obtained from reactions of carbanion **1** with *t*-BuBr or neopentyl bromide, and it was concluded that the rate-limiting step in these reactions, as well as that of **1** with 1-adamantyl bromide, was an outer-sphere single electron transfer. The $k_{\text{obsd}}/k_{\text{SET}}$ ratios for reactions of **1** with these bromides were 2.5, 1.3, and 0.8, respectively, compared to 170, 400, and 2500 for *sec*-BuBr, *n*-BuBr, and EtBr, respectively. In a later paper the reaction of **1** with C₆H₅C(Me)(Et)Cl was reported to give $k_{\text{obsd}}/k_{\text{SET}} = 1.3$. The reaction is thus pure outer-sphere SET; PhCH(Me)Cl gave $k_{\text{obsd}}/k_{\text{SET}} = 8.2$, and the reaction was classified as borderline.¹¹

The results of the application of Ebersson-type calculations to the reactions of the series of $9\text{-R}_2\text{NFl}^-$ ions with $\text{F}_3\text{CCH}_2\text{I}$ and Ph_2CHCl are shown in Table II. The value of λ was modeled after that used by Ebersson for PhCH₂Cl and Ph₃CCl,⁷ and that for $\text{F}_3\text{CCH}_2\text{I}$ was chosen by trial and error to give the best fit for the data. Examination of Table II shows that the $k_{\text{obsd}}/k_{\text{SET}}$ ratios are near unity for the two reactions, indicating that both are reacting by SET mechanisms involving radical-pair intermediates, $\text{F}_3\text{CCH}_2\text{I}$ giving radical-type products,⁶ and Ph_2CHCl giving S_N2-type products. The results with $\text{F}_3\text{CCH}_2\text{I}$ and Ph_2CHCl provide strong support for Ebersson's method of SET calculation in that, for the first time, a family of anions has been used in a

Table III. Comparison of Rates (k_{obsd}) with Expected Rates of Single Electron Transfer (k_{SET}) for Reactions of $9\text{-R}_2\text{NFl}^-$ with (*p*-ClC₆H₄)₂CHCl and PhCH₂Cl

NR ²	$(p\text{-ClC}_6\text{H}_4)_2\text{CHCl}^b$			PhCH ₂ Cl ^d		
	ΔG_0^a	$\log k_{\text{obsd}}$	$k_{\text{obsd}}/k_{\text{SET}}^c$	ΔG_0^a	$\log k_{\text{obsd}}$	$k_{\text{obsd}}/k_{\text{SET}}^c$
	-8.3	+1.36	2.0	-3.3	+1.35	71
	-7.0	+0.67	0.89	-2.0	+0.46	28
NMe ₂	-3.6	+0.25	4.6	+1.8	+0.11	320
	-3.2	+0.39	8.7	+1.4	+0.27	300
	-2.2	-0.45	2.9	+2.8	-1.14	46
N(<i>i</i> -Pr) ₂	+0.4	-1.20	4.4	+5.4	-2.16	44
	+1.5	-1.20	11	+6.5	-2.49	47

^aIn kcal/mol. ^b $E_{\text{rd}} = 0.4 \times (-1.259) = -0.503$; ²³ $\lambda = 65$. S_N2 products were obtained. ^cCalculated from eq 2. ^d $E_{\text{rd}} = 0.52 + 0.20 = 0.72$ (Ebersson's E_{rd} ⁹ referenced to SHE_{aq}). S_N2 products were formed.

reaction known to occur by a SET mechanism, and the correspondence between k_{obsd} and k_{SET} is good over a considerable $E_{\text{ox}}(\text{A}^-)$ range. In reactions of this family, factors such as the strength of the bonds being formed and broken and the nature of the solvent and geometric reorganization are kept essentially constant.






There appears to be little prospect of trapping the $\text{Ph}_2\text{CHCl}^{\cdot-}$ anion radical, since its lifetime has been shown to be less than 20 ps,¹² corresponding to a rate constant of $>5 \times 10^{10} \text{ s}^{-1}$. It has been suggested that, in SET to Ph_2CHCl , the $\text{Ph}_2\text{CH}^{\cdot}$ radical and Cl⁻ ion are formed directly without the $\text{Ph}_2\text{CHCl}^{\cdot-}$ radical ion as an intermediary. The PhCH_2^{\cdot} radical will also be difficult to trap since it is expected to dimerize at a diffusion-controlled rate. On the other hand, the $9\text{-}i\text{-Pr}_2\text{NFl}^{\cdot}$ radical is persistent when formed in the reaction of $9\text{-}i\text{-Pr}_2\text{NFl}^-$ with $\text{F}_3\text{CCH}_2\text{I}$, presumably because it is relatively stable and dimerization is inhibited by steric hindrance. Additional support for the SET mechanism was obtained by observation of a peak at 450 nm, characteristic of the $9\text{-}i\text{-Pr}_2\text{NFl}^{\cdot}$ radical,⁶ in a reaction with Ph_2CHCl . The peak is not persistent under these conditions, however, presumably because of rapid coupling with the $\text{Ph}_2\text{CH}^{\cdot}$ radical. No indication of the presence of $(\text{Ph}_2\text{CH})_2$ or $(9\text{-R}_2\text{NFl})_2$ dimers was evident in the product NMR. This is not surprising since even if the $\text{Ph}_2\text{CH}^{\cdot}$ and $9\text{-R}_2\text{NFl}^{\cdot}$ radicals escape from the cage, theory tells us that, if dimerization of one of the radicals is slow (as is expected for $9\text{-R}_2\text{NFl}^{\cdot}$), the cross-coupling product will be overwhelmingly favored.¹³

The results for $9\text{-R}_2\text{NFl}^-$ ions reacting with Ph_2CHCl were confirmed by repeating the experiments with (*p*-ClC₆H₄)₂CHCl. Since the Hammett σ for *p*-Cl is +0.24 we expected E_{rd} with this chloride to be less negative than that of the parent, but the E_{rd} proved to be slightly more negative. (Evidently the π -donor properties of the *p*-Cl substituent are more decisive in determining the size of E_{rd} than are its field-inductive acceptor properties.) The electron-transfer reactions with this acceptor are therefore a little less exergonic than with Ph_2CHCl . The $k_{\text{obsd}}/k_{\text{SET}}$ ratios with this acceptor are generally larger than unity for this acceptor, instead of being smaller (Table III).


The results with the $9\text{-R}_2\text{NFl}^-$ ion family reacting with Ph_2CHCl and (*p*-ClC₆H₄)₂CHCl agree well with those reported for carbanion **1** reacting with *t*-BuBr, 1-AdBr, *t*-BuCH₂Br,³ and C₆H₄C(Me)(Et)Cl.¹¹ Carbanion **1** has an oxidation potential of -1.48 V (vs SHE_{aq}), which means that its potential is 0.615 V (14.2 kcal/mol) more negative than the best $9\text{-R}_2\text{NFl}^-$ ion donor and 1.042 V (24 kcal/mol) more negative than the worst. In

(11) Lund, T.; Lund, H. *Acta Chem. Scand. Ser. B* **1987**, *B41*, 93-102.(12) Simon, J. D.; Peters, K. S. *Organometallics* **1983**, *2*, 1867-1869.(13) Fisher, H. J. *Am. Chem. Soc.* **1986**, *108*, 3925-3927.

Table IV. Rate Constants for Reactions of 9-R₂NF⁻ Ions with *n*-Butyl Bromide and Cyclohexyl Bromide

R ₂ N	<i>n</i> -BuBr ^b			<i>c</i> -C ₆ H ₁₁ Br ^d		
	ΔG ⁰ ^a	log k _{obsd}	k _{obsd} /k _{SET} ^c	ΔG ⁰ ^a	log k _{obsd}	k _{obsd} /k _{SET} ^c
	+3.3	+0.80	1600	-1.03	-0.250	3.1
	+4.7	+0.33	1700	+0.35	-0.777 ^e	3.4
NMe ₂	+8.1	-0.04	22000			
	+8.5	+0.01	32000	+4.08	-1.03	4.7
	+9.5	-1.24	4600	+5.11	-1.82	17
N(<i>i</i> -Pr) ₂	+12.1	-2.62	2200	+7.7	-2.12 ^e	95
	+13.2	-2.80	3100			

^aIn kcal/mol. ^bE_{rd} = -0.82 - 0.20 = -1.01 (Ebersson's E_{rd}⁹ referenced to SHE_{aq}); λ = 62 (Lund's λ = 73).¹¹ ^cCalculated from eq 1. ^dE_{rd} = 0.4 × (-2.05) = -0.82 vs SHE_{aq}.²³ λ = 62; S_N2 and E2 products were formed. ^eResults of C. A. Wilson.

Scheme I

reactions of **1** with PhCH₂Cl and *p*-MeOC₆H₄CH₂Cl, *k*_{obsd}/*k*_{SET} ratios of 66 and 93, respectively, were reported,¹¹ which agree well with ratios obtained for reactions of PhCH₂Cl with four of the 9-R₂NF⁻ ions shown in Table III.

Our study of reactions of 9-R₂NF⁻ ions was next extended to *n*-butyl and cyclohexyl bromides, which have E_{rd} potentials at least 100–300 mV more negative than that of PhCH₂Cl (Table IV). The *k*_{obsd}/*k*_{SET} ratios with *n*-BuBr were 2–4 orders of magnitude higher than Lund's ratio for the reaction of **1** with *n*-BuBr. On the other hand, the *k*_{obsd}/*k*_{SET} ratios observed for the reactions of 9-*c*-C₃H₆NF⁻, 9-*c*-C₄H₈NF⁻, and 9-*c*-C₅H₁₀NF⁻ ions with *c*-C₆H₁₁Br were similar to those found for reactions with (*p*-ClC₆H₄)₂CHCl (Table III). It seems likely that the greater steric hindrance in the *c*-C₆H₁₁Br electrophile may be favoring the SET reaction, a result that would be similar to that of Lund's in comparisons of *k*_{obsd}/*k*_{SET} ratios for *t*-BuBr, 1-AdBr, and Me₃CCH₂Br vs EtBr, *n*-BuBr, and *sec*-BuBr.^{3a} It is noteworthy in this regard that, as will be brought out shortly, our ratios with *c*-C₆H₁₁Br are 1 or 2 orders of magnitude smaller than Lund's with the less hindered *sec*-BuBr. The reactions of 9-R₂NF⁻ ions with *n*-BuBr give high yields of substitution products, whereas those with *c*-C₆H₁₁Br give a mixture of substitution and elimination products.

The *k*_{obsd} values for the reactions of 9-R₂NF⁻ ions with Ph₂CHCl and (*p*-ClC₆H₄)₂CHCl, which are occurring by SET mechanisms, are compared graphically with those with PhCH₂Cl in Figure 1. The gently sloping line is that calculated for the SET mechanism from eq 3. Figure 2 shows a similar comparison for the reactions with *c*-C₆H₁₁Br and *n*-BuBr.

Mechanisms of Reactions of 9-R₂NF⁻ Ions with Alkyl Halides.

The mechanism for reaction of 9-GF⁻ ions with F₃CCH₂I or Ph₂CHCl is assumed to occur by a rate-determining outer-sphere dissociative SET, as outlined in general terms in Scheme I. In the first step a molecular ion complex is formed. The binding




Figure 1. Marcus plot of log *k*_{obsd} for reaction of 9-(dialkylamino)fluorene ions with Ph₂CHCl, (*p*-ClC₆H₄)₂CHCl, and PhCH₂Cl vs their free energy of electron transfer (ΔG⁰). The solid line represents part of the expected Marcus parabola for λ = 65.




Figure 2. Marcus plot of log *k*_{obsd} for reaction of 9-(dialkylamino)fluorene ions with *c*-C₆H₁₁Br and *n*-BuBr vs their free energy of electron transfer (ΔG⁰). The solid line represents part of the expected Marcus parabola for λ = 62.

of D⁻ to RX is weak, but a long-range SET can take place because ΔG⁰ is in the range of +3 to -6.8 kcal/mol for reactions of 9-R₂NF⁻ ions with F₃CCH₂I and +0.8 to -9 kcal/mol with Ph₂CHCl (Table II). The electrostatic attraction in the transition states for such outer-sphere SETs is of the order of 1 kcal/mol according to the Marcus theory.¹ For reactions of 9-PhCH₂F⁻ with F₃CCH₂I, disproportionation products from the 9-PhCH₂F⁻ radical and products from the attack of F₃CCH₂• radicals on the fluorene ring have been identified.⁶ For reactions of a variety of 9-GF⁻ ions with Ph₂CHCl, good yields of substitution products have been obtained, but the calculations in Table II indicate that with 9-R₂NF⁻ ions the reactions occur by an SET mechanism (Table II). Similar results are obtained in reactions of 9-R₂NF⁻ ions with (*p*-ClC₆H₄)₂CHCl (Table III).

Reaction of PhCH₂Cl with 9-R₂NF⁻ ions give *k*_{obsd}/*k*_{SET} ratios of 30–320. The calculations appear to indicate that these reactions, as well as those with *c*-C₆H₁₁Br (Table IV), are in a borderline region. Since the reactions of 9-R₂NF⁻ ions with PhCH₂Cl show larger steric effects than do the reactions with the more sterically demanding Ph₂CHCl electrophile, it is clear that the former are

more S_N2 in nature. The reactions of 9- R_2NFI^- ions with *n*-BuBr are further in the endergonic region and the k_{obsd}/k_{SET} ratios are an order of magnitude larger. Reactions that are too endergonic to proceed by the SET pathway can be accommodated in terms of Scheme I by assuming formation of a second ion-molecule complex with a stronger electrostatic attraction bringing the reactants into a closer proximity for reaction. This would require desolvation and some molecular reorganization, but the energy required can be more than repaid by the relatively low barrier provided by the synchronous SET process.¹⁴ It seems likely that when k_{obsd}/k_{SET} ratios are near unity, or below, the radical pair mechanism will obtain, and that when the ratio is 10^2 or above, the concerted S_N2 mechanism will obtain. In the borderline region the two mechanisms may be in competition or, as proposed elsewhere,^{2,15} hybrid or merged mechanisms may occur wherein the distance of approach of D^- and RX is intermediate to the stepwise and concerted SET extremes, resulting in partial bonding between D^* and R^* as electron transfer occurs.

Summary and Conclusions

A series of seven 9- R_2NFI^- ions wherein the R_2N groups are known to possess steric retardation rate (*r*) factors as much as 3.4 orders of magnitude *greater* than that of Me were found to give rate ratios, $k^{PhCH_2Cl}/k^{Ph_2CHCl}$, that were all at least 1 order of magnitude *smaller* than that of Me. We conclude that the 9- R_2NFI^- ions are reacting with the more sterically demanding Ph_2CHCl electrophile by a single electron transfer (SET) mechanism. This conclusion was supported by (a) the observation of near unity rate ratios of k_{obsd} values to k_{SET} , the rate constant calculated by using the Marcus equation, and (b) by a reasonable fit to a Marcus plot (Figure 1). High yields of substitution products were formed in reactions of 9-GFI⁻ ions with Ph_2CHCl as well as $PhCH_2Cl$. The k_{obsd}/k_{SET} ratios for 9- R_2NFI^- ions with F_3CCH_2I , a known SET acceptor, and (*p*-ClC₆H₄)₂CHCl were comparable. For substitution reactions of $PhCH_2Cl$ and substitution/elimination reactions of *c*-C₆H₁₁Br with 9- R_2NFI^- ions, the k_{obsd}/k_{SET} ratios were, with few exceptions, larger, and the fit with a Marcus plot were poorer (Figures 1 and 2). The mechanisms for these reactions appear to be in a borderline SET- S_N2 region. The k_{obsd}/k_{SET} ratio for 9- R_2NFI^- ions reacting with *n*-BuBr was 10^3 - 10^4 , which places the mechanism in the time-honored concerted S_N2 category.

Experimental Section

The methods used for determination of rate constants¹⁶ and cyclic voltammetry measurements¹⁷ have been described in earlier publications. ¹H NMR spectra were obtained with a Varian EM-390 spectrometer and chemical shifts are reported relative to tetramethylsilane. Melting points were measured with a Thomas-Hoover capillary apparatus and are uncorrected. Mass spectra were run by H. L. Hung on a HP5984 GC/MS system.

Materials. Benzyl chloride, benzhydryl chloride, 2,2,2-trifluoroethyl iodide, *n*-butyl bromide, and cyclohexyl bromide were obtained commercially and purified by distillation prior to use.

4,4'-Dichlorobenzhydryl chloride was prepared from the reaction of 4,4'-dichlorobenzhydryl with thionyl chloride (1.2 equiv) in refluxing benzene for 1 h. The product was recrystallized from ether and washed with hexane: mp 60-60.5 °C (lit.¹⁸ mp 63 °C); ¹H NMR (CDCl₃) δ 6.0 (s, 1 H), 7.22 (br s, 10 H). It should be noted that use of this compound (despite proper precautions against exposure) caused an allergic response in one researcher (J.A.H.) resembling chloracne, which Sax¹⁹ describes for chlorinated diphenyls. A previous researcher experienced no such problems.

Preparation of the 9-(dialkylamino)fluorenes was described in a preceding paper.^{4b}

Product Studies. Reactions of 9-GFI⁻ ions with Ph_2CHCl , $PhCH_2Cl$, *n*-BuBr, and other simple alkyl halides have been shown to yield S_N2 products in numerous earlier studies.^{4,8,16,20} In particular, 9-*c*-C₆H₈NFI⁻ was shown to give 70% S_N2 product and 30% 9-*c*-C₆H₈NFIH (E2 product) when reacted with *c*-C₆H₁₁Br.²¹ Additional studies were carried out for cases where products other than expected S_N2 products might be possible and are reported below.

Reaction of 9-Me₂NFI⁻ Ion with Ph_2CHCl . 9-(Dimethylamino)-fluorene (112 mg, 5.4×10^{-4} mol) was reacted with 1 equiv of $CH_3SOCH_2^-K^+$ and then with 1 equiv of benzhydryl chloride for 10 min. The reaction was quenched with water and the product was extracted with ether. Residual Me₂SO was removed from the ether layer by multiple washing with water. After drying of the layer with anhydrous magnesium sulfate, the ether was removed under reduced pressure, giving 198 mg (98%) of the crude substitution product: mp 118-126 °C; ¹H NMR (CDCl₃) δ 2.25 (s, 6 H), 5.25 (s, 1 H), 7.0-7.5 (m, 18 H); MS, *m/e* 375 (24.9), 374 (molecular ion, 81.1), 374 (81.1), 371 (20.3), 208 (57.4), 192 (39.4), 165 (100); recrystallization from ethanol gave material with mp 136-136.5 °C. (The ¹H NMR (CDCl₃) of the recrystallized material was identical with that of the crude product.) Anal. Calcd for C₂₈H₂₅N: C, 89.55; H, 6.71; N, 3.74. Found: C, 89.43; H, 6.64; N, 3.77.

Reaction of 9-*c*-C₆H₈NFI⁻ Ion with Ph_2CHCl . A 113-mg (4.8×10^{-4} mol) sample of 9-pyrrolidinylfluorene was treated with 1 equiv of $CH_3SOCH_2^-K^+$ and reacted for 10 min with 1 equiv of benzhydryl chloride. Workup gave 161 mg (83%) of the crude substitution product: mp 129-132 °C; ¹H NMR (CDCl₃) δ 1.5-1.7 (br, 4 H), 2.4-2.7 (br t, 4 H), 5.25 (s, 1 H), 6.9-7.6 (m, 18 H); MS, *m/e* 403 (1.2), 402 (molecular ion, 3.0), 332 (99.2), 234 (100); recrystallization from ethanol gave material with mp 135-136 °C. The ¹H NMR (CDCl₃) of recrystallized material was identical with that of the crude product.

Reaction of 9-*i*-Pr₂NFI⁻ with Ph_2CHCl . A 39-mg (1.5×10^{-4} mol) sample of 9-(diisopropylamino)fluorene was reacted with 1 equiv of $CH_3SOCH_2^-K^+$ and 1 equiv of benzhydryl chloride for 24 h. Workup gave 65 mg (92%) of the crude substitution product as a semisolid: ¹H NMR (CDCl₃) δ 1.0 (d, 12 H), 3.45 (m, 2 H), 5.9 (s, 1 H), 7.1-8.2 (m, 18 H); MS, *m/e* 432 (3.7), 431 (molecular ion, 9.8), 416 (21.4), 331 (100), 165 (49).

Reaction of 9-*p*-MePhFI⁻ Ion with Ph_2CHCl . 9-*p*-Tolylfluorene (208 mg, 8.1×10^{-4} mol) was reacted^{8a} with 1 equiv of $CH_3SOCH_2^-K^+$ and then with 1 equiv of benzhydryl chloride for 2 h. Workup gave 330 mg (97%) of crude substitution product: mp 164-178 °C; ¹H NMR (CDCl₃) δ 2.2 (s, 3 H), 5.6 (s, 1 H), 6.5-7.5 (m, 22 H). Recrystallization from hexane/ethyl acetate gave material with mp 176-177 °C.

Reaction of 9-PhFI⁻ Ion with (*p*-ClC₆H₄)₂CHCl. 9-Phenylfluorene (243 mg, 1.0×10^{-3} mol) was reacted²² with 1 equiv of $CH_3SOCH_2^-K^+$ and then 1 equiv of 4,4'-dichlorobenzhydryl chloride for 1.5 h. Workup gave 472 mg (92%) of crude substitution product: ¹H NMR (CDCl₃) δ 5.5 (s, 1 H), 6.4-7.8 (m, 21 H); MS, *m/e* 477 (1.2), 241 (100), 237 (44.3), 235 (68.5); recrystallization (ethanol/hexane) gave a material with mp 176.5-177.5 °C.

Reaction of 9-*i*-Pr₂NFI⁻ Ion with *n*-BuBr. 9-(Diisopropylamino)-fluorene (94 mg, 3.5×10^{-4} mol) was reacted with 1 equiv of $CH_3SOCH_2^-K^+$ and then with 1 equiv of *n*-butyl bromide for 24 h. Workup gave 94 mg (82%) of the substitution product as a viscous oil: ¹H NMR (CDCl₃) δ 0.9 (d, 12 H), 1.2-2.3 (br, 9 H), 3.6 (m, 2 H), 7.1-7.8 (m, 8 H); MS, *m/e* 265 (22.8), 264 (100), 222 (49.6), 165 (33.9).

Reaction of 9-*c*-C₆H₈NFI⁻ Ion with *n*-BuBr. 9-Pyrrolidinylfluorene (32.2 mg, 1.4×10^{-4} mol) was reacted with 1 equiv of $CH_3SOCH_2^-K^+$ and then with 1 equiv of *n*-butyl bromide for 18 h. Workup gave 39.8 mg (85%) of substitution product as a viscous oil: ¹H NMR (CDCl₃) δ 0.6-0.9 (m, 5 H), 1.0-1.3 (m, 2 H), 1.5-1.7 (m, 4 H), 2.15-2.35 (m, 2 H), 2.4-2.6 (m, 4 H), 7.15-7.8 (m, 8 H); MS, *m/e* 291 (molecular ion, 3.0), 234 (100), 165 (13.7).

Reaction of 9-*i*-Pr₂NFI⁻ Ion with *c*-C₆H₁₁Br. 9-(Diisopropylamino)-fluorene (80.1 mg, 3.0×10^{-4} mol) was reacted with 1 equiv of $CH_3SOCH_2^-K^+$ and then with 1 equiv of cyclohexyl bromide for 24 h. Workup gave 90.9 mg of product which was assigned by NMR as 50%

(14) For a recent discussion of the importance of "critical distance" in chemical reactions, see: (a) Menger, F. M. In *Nucleophilicity*; Harris, J. M., McManus, S. P., Eds.; Advances in Chemistry 215, American Chemical Society: Washington, DC, 1987; Chapter 14. (b) Menger, F. M. *Acc. Chem. Res.* **1985**, *18*, 128-134.

(15) Pross, A.; Shaik, S. S. *Acc. Chem. Res.* **1983**, *16*, 363-370.

(16) Bordwell, F. G.; Hughes, D. L. *J. Org. Chem.* **1980**, *45*, 3314-3320.

(17) Bordwell, F. G.; Bausch, M. J. *J. Am. Chem. Soc.* **1986**, *108*, 1979-1985.

(18) Norris, J. F.; Banta, C. *J. Am. Chem. Soc.* **1928**, *50*, 1808-1812.

(19) Sax, N. I. *Dangerous Properties of Industrial Materials*, 6th ed; Van Nostrand-Reinhold: New York, 1984.

(20) Bordwell, F. G.; Mrozack, S. R. *J. Org. Chem.* **1982**, *47*, 3802-3803.

(21) Wilson, C. A., unpublished results.

(22) Mrozack, S. R. Ph.D. Dissertation, Northwestern University, 1982.

(23) The observed E_{red} , measured as a peak potential (E_p), has been multiplied by 0.4 to correct for the anionic shift caused by the rapid follow reaction. This fraction is patterned after the estimates of Ebersohn, which indicate that $E_{1/2}$ is reduced by 30-40% in Me₂SO or DMF for active halides, including $CH_2=CHCH_2Cl$, $CH_2=CHCH_2Br$, $PhCH_2Cl$, $PhCH_2Br$, and *t*-BuBr.^{9c}

S_N2 -type substitution product and 50% 9-*i*-Pr₂NFl⁻H (from an elimination reaction): ¹H NMR (CDCl₃) δ 1.0–2.0 (m, 18.2 H), 2.9 (br, 2 H), 4.95 (s, 0.5 H), 7.2–7.8 (m, 8 H); MS, *m/e* 348 (1.3), 347 (substitution molecular ion, 3.1), 265 (19.2), 265 (elimination molecular ion, 61.7), 247 (30.6), 222 (36.0), 165 (100).

UV/Vis Spectra of 9-*i*-Pr₂NFl⁻ + Ph₂CHCl Reaction. The anion of 9-(diisopropylamino)fluorene has been shown to have three peaks (λ_{\max} = 479, 510, 556 nm) in Me₂SO while the corresponding radical has a peak at 450 nm.²¹ UV/vis spectra taken at various points during the reaction of *i*-Pr₂NFl⁻ ion with benzhydryl chloride showed the appearance of a small peak at 450 nm in addition to the three anion peaks. However, unlike reactions of this anion with electron acceptors (PhSO₂CH₂Cl, *c*-C₆H₁₀(NO₂)Ts, and F₃CCH₂I), the radical peak is not persistent and a colorless solution results when all of the anion absorbance has dissipated, presumably because of coupling of the *i*-Pr₂NFl⁻ radical with the Ph₂CH[•] radical.

Acknowledgment. This work was supported by the National Science Foundation. We are indebted to C. A. Wilson for some of the rate and product studies with benzyl chloride and cyclohexyl

bromide and to T.-Y. Lynch for the rate data for F₃CCH₂I and for checking for the appearance of 9-*i*-Pr₂NFl⁻ in the product study of 9-*i*-Pr₂NFl⁻ with Ph₂CHCl.

Registry No. 9-MeFl⁻, 31468-21-0; 9-PhCH₂Fl⁻, 53629-11-1; 9-MeOFl⁻, 71805-70-4; 9-(4-MeC₆H₄)Fl⁻, 42730-14-3; 9-PhFl⁻, 31468-22-1; 9-(3-ClC₆H₄)Fl⁻, 73872-45-4; 9-(4-MeSO₂C₆H₄)Fl⁻, 73872-44-3; 9-*c*-C₆H₁₁Fl⁻, 117959-61-2; 9-*t*-BuFl⁻, 73838-69-4; 9-*c*-C₃H₆NFl⁻, 111933-70-1; 9-*c*-C₄H₈NFl⁻, 111933-71-2; 9-Me₂NFl⁻, 83936-70-3; 9-*c*-C₅H₁₀NFl⁻, 111933-72-3; 9-(2-Me-*c*-C₅H₉N)Fl⁻, 111933-73-4; 9-(2,2,6,6-Me₄-*c*-C₅H₉N)Fl⁻, 111933-74-5; 9-*i*-Pr₂NFl⁻, 109495-02-5; Ph₂CHCl₂, 90-99-3; PhCH₂Cl, 100-44-7; F₃CCH₂I, 353-83-3; (*p*-ClC₆H₄)₂CHCl, 782-08-1; *n*-BuBr, 109-65-9; *c*-C₆H₁₁Br, 108-85-0; 9-Me₂NFl, 53156-46-0; 9-Ph₂CH-9-Me₂NFl, 117959-62-3; 9-*c*-C₄H₈NFl, 7596-59-0; 9-Ph₂CH-9-*c*-C₄H₈NFl, 117959-63-4; 9-*i*-Pr₂NFl, 109495-00-3; 9-Ph₂CH-9-*i*-Pr₂NFl, 117959-64-5; 9-(4-MeC₆H₄)Fl, 18153-43-0; 9-Ph₂CH-9-(4-MeC₆H₄)Fl, 117959-65-6; 9-PhFl, 789-24-2; 9-(*p*-ClC₆H₄)₂CH-9-PhFl, 117959-66-7; 9-Bu-9-*i*-Pr₂NFl, 117959-67-8; 9-*c*-C₆H₁₁-9-*i*-Pr₂NFl, 117959-69-0; 9-*i*-Pr₂NFl⁻, 117959-70-3; 4,4'-dichlorobenzhydryl, 90-97-1.

Synthesis of (2*R*,3*R*)- and (2*S*,3*S*)-[2,3-²H₂]Oxirane and Application of It to the Synthesis of Chirally Labeled Homoserine[†]

John M. Schwab,* Tapan Ray, and Chong-Kei Ho

Contribution from the Department of Medicinal Chemistry and Pharmacognosy, School of Pharmacy and Pharmaceutical Sciences, Purdue University, West Lafayette, Indiana 47907.

Received May 16, 1988

Abstract: (2*R*,3*R*)- and (2*S*,3*S*)-[2,3-²H₂]oxirane have been synthesized from 2-propynol, the key step being the asymmetric epoxidation of (*E*)-3-(triphenylsilyl)-2-propenol. To determine the enantiomeric purities of the oxiranes, they were reacted with phenyllithium, and the resulting 2-phenylethanol samples were converted to esters of (1*S*)-(-)-camphanic acid. ²H NMR analysis (in the presence of Eu(dpm)₃) showed that (*R*)- and (*S*)-oxirane had ee values of 92% and 94%, respectively. The utility of (2*R*,3*R*)- and (2*S*,3*S*)-[2,3-²H₂]oxirane as chiral labeling synthons was demonstrated by a two-step synthesis of chirally labeled homoserine lactone, which was resolved chromatographically as its *N*-(3,5-dinitrobenzoyl) derivative. The diastereomeric purity of the latter was assessed by ¹H NMR. Acid hydrolysis of (2*R*,3*S*,4*S*)-*N*-benzoyl[3,4-²H₂]homoserine lactone resulted in extensive epimerization at C-4 of the lactone. An ¹⁸O-labeling experiment failed to support a mechanism involving amide participation. It is concluded that the lactone was hydrolyzed by an unprecedented A_{AL}2 mechanism.

In many cases, syntheses of chirally labeled compounds are quite long, owing to the limited number of reagents and stereoselective reactions by which isotopic labels can be introduced.¹⁻³ While this has been true in our own work, in one recent instance it was clear that the necessary labeled substrates could be made economically by using a divergent route in which chirally labeled oxirane would serve as a common synthetic intermediate.

In this paper, we report the details of our synthesis and analysis of (2*R*,3*R*)- and (2*S*,3*S*)-[2,3-²H₂]oxirane, including improvements over the methods that we have described in a preliminary communication.⁴ The general utility of chirally labeled oxirane is also demonstrated by the synthesis of homoserine that is chirally labeled at carbons 3 and 4.⁵ In addition, in the course of the homoserine synthesis, a mechanistically unprecedented acid-catalyzed epimerization at homoserine C-4 was observed.

Results

Synthesis and Analysis of Chirally Labeled Oxirane. The synthesis of chirally labeled oxirane (Scheme I) begins with 3-(triphenylsilyl)-2-propynol,^{6,7} which is readily available from

propargyl alcohol.⁶ Reduction of the triple bond with lithium aluminum deuteride,⁸ using deuterium oxide to quench the reaction, leads to (*E*)-3-(triphenylsilyl)-2-[2,3-²H₂]propenol.^{6,9} This labeled allylic alcohol is epoxidized by using a modified Sharpless procedure,^{10,11} with (+)- and (-)-diisopropyl tartrate (DIPT) giving

(1) Parry, R. J. In *Bioorganic Chemistry*; Van Tamelen, E. E., Ed.; Academic: New York, 1978; Vol. 2, pp 247-272.

(2) Young, D. W. In *Tritium in Organic Chemistry*; Buncel, E., and Lee, C. C., Eds.; Elsevier: Amsterdam, 1978; pp 177-294.

(3) Hill, R. K. In *Bioorganic Chemistry*; Van Tamelen, E. E., Ed.; Academic: New York, 1978; Vol. 2, pp 111-151.

(4) Schwab, J. M.; Ho, C.-K. *J. Chem. Soc., Chem. Commun.* **1986**, 872-873.

(5) Schwab, J. M.; Ray, T. *J. Chem. Soc., Chem. Commun.* **1988**, 29-31.

(6) For the synthesis of the trimethylsilyl analogue, see: Denmark, S. E.; Jones, T. K. *J. Org. Chem.* **1982**, *47*, 4595-4597.

(7) For the synthesis of the trimethylsilyl analogue, see: Mironov, V. F.; Maksimova, N. G. *Izv. Akad. Nauk SSSR* **1960**, 2059-2061.

(8) For the synthesis of the trimethylsilyl analogue, see: Grant, B.; Djerassi, C. *J. Org. Chem.* **1974**, *39*, 968-970.

(9) For the synthesis of the trimethylsilyl analogue, see: Stork, G.; Jung, M. E.; Colvin, E.; Noel, Y. *J. Am. Chem. Soc.* **1974**, *96*, 3684-3686.

(10) Wang, Z.-m.; Zhou, W.-s.; Lin, G.-q. *Tetrahedron Lett.* **1985**, *26*, 6221-6224.

(11) Katsuki, T.; Sharpless, K. B. *J. Am. Chem. Soc.* **1980**, *102*, 5974-5976.

[†] C.-K.H.'s portion of these studies was performed in partial fulfillment of the requirements for the Ph.D. degree from the Department of Chemistry, The Catholic University of America, Washington, DC 20064.