

asymmetric dihydroxylation of olefins recently developed.¹¹ We believe that the simple strategy of using a diether of a C₂ symmetric chiral diol as a stereocontrol catalyst provides a basis for further development of asymmetric reactions.¹²

(11) Tomioka, K.; Nakajima, M.; Koga, K. *J. Am. Chem. Soc.* **1987**, *109*, 6213. Lohray, B. B.; Kalantar, T. H.; Kim, B. M.; Park, C. Y.; Shibata, T.; Wai, J. S. M.; Sharpless, K. B. *Tetrahedron Lett.* **1989**, *30*, 2041.

(12) We are grateful to the Japan Foundation for Optically Active Compounds and the Hoansha Foundation and for a Grant-in-Aid for Scientific Research, Ministry of Education, Science and Culture, Japan (No. 01571143), for partial financial support.

Transition-Metal-Mediated Thiosulfinate Ester Synthesis

Marlene E. Raseta, Steven A. Cawood, and Mark E. Welker*

Department of Chemistry, Wake Forest University
Winston-Salem, North Carolina 27109

Arnold L. Rheingold

Department of Chemistry, University of Delaware
Newark, Delaware 19711

Received June 8, 1989

Unlike sulfur dioxide (SO₂), the coordination and organic reaction chemistry of disulfur monoxide (S₂O) has received little attention. A handful of transition-metal S₂O complexes have been synthesized via oxidation of the corresponding disulfur (S₂) complexes.¹ A few Diels-Alder reactions of simple dienes with S₂O have also been reported.² However, there were no reports of direct S₂O complex synthesis prior to our initial work.³

In order to undertake a detailed study of the chemistry of S₂O or S₂O complexes, a reliable source of S₂O or an S₂O equivalent was needed. Previously reported procedures for S₂O generation suitable for use in synthetic studies yielded mixtures of S₂O, SO₂, and SO.^{2,4} We recently published a synthesis of 4,5-diphenyl-3,6-dihydro-1,2-dithiin 1-oxide (**2**) which liberates S₂O via a transition-metal-assisted retro-Diels-Alder reaction.^{3,5} Here we

Scheme I





Table I. Cyclizations of 2-Alkynyl Complexes **12**

	M	n	m	R	yield, %
13a	Fe	5	2	CH ₃	74
13b	Fe	5	2	Ph	81
13c	Fe	0	2	CH ₃	71
13d	Fe	0	2	Ph	72
13e	Mo	0	3	CH ₃	45 ^a
13f	Mo	0	3	Ph	43 ^a

^a Greater than 90% based on recovered starting material.

report further on the unusual reactivity of **2** and its utilization in the synthesis of cyclic thiosulfinate esters.



Thiosulfinate esters have shown biological activity as antibacterials,⁶ antifungals,⁶ antivirals,⁷ plant growth regulators,⁸ platelet aggregation inhibitors,⁹ and tumor growth inhibitors.¹⁰ Compounds containing this functional group have practical applications as alkene autoxidation inhibitors¹¹ and radioprotective agents.¹² Thiosulfinate esters of particular synthetic interest to us were analogues of the asparagusic acid S-oxides (**5**) (potent plant growth regulators)⁸ and the brugerols (**6**) (isolated from mangroves).¹³ One possible route to compounds of this general



type would be a (3 + 2) cycloaddition reaction between transition-metal 2-alkynyl complexes (**7**) and S₂O (**8**) (Scheme I). These complexes (**7**) have been shown to react in this manner with a variety of other small organic electrophiles.¹⁴ If cyclization

(1) (a) Faller, J. W.; Ma, Y. *Organometallics* **1989**, *8*, 609. (b) Schenk, W. A. *Angew. Chem., Int. Ed. Engl.* **1987**, *26*, 98. (c) Herberhold, M.; Schmidkonz, B. *J. Organomet. Chem.* **1986**, *308*, 35. (d) Herberhold, M.; Hill, A. F. *J. Organomet. Chem.* **1986**, *309*, C29. (e) Hoots, J. E.; Rauchfuss, T. B. *Inorg. Chem.* **1984**, *23*, 3130. (f) Hoots, J. E.; Rauchfuss, T. B.; Wilson, S. R. *J. Chem. Soc., Chem. Commun.* **1983**, 1226. (g) Dirand-Colin, J.; Schappacher, M.; Ricard, L.; Weiss, R. *J. Less-Common Met.* **1977**, *54*, 91. (h) Schmid, G.; Ritter, G. *Chem. Ber.* **1975**, *108*, 3008. (i) Schmid, G.; Ritter, G. *Angew. Chem., Int. Ed. Engl.* **1975**, *14*, 645.

(2) Dodson, R. M.; Srinivasan, V.; Sharma, K. S.; Sauer, R. F. *J. Org. Chem.* **1972**, *37*, 2367.

(3) Urove, G. A.; Welker, M. E. *Organometallics* **1988**, *7*, 1013.

(4) (a) Schenk, P. W.; Stuedel, R. *Angew. Chem., Int. Ed. Engl.* **1964**, *3*, 61. (b) For a review of methods used to produce S₂O for spectroscopic studies, see: Vasudeva Murthy, A. R.; Narayanan, T. R.; Sharma, D. K. *Int. J. Sulfur Chem., Part B* **1971**, *6*, 161.

(5) Use of retro-Diels-Alder reactions for the production of singlet oxygen^{5a} and singlet sulfur^{5b} have been reported: (a) Wasserman, H. H.; Scheffer, J. R.; Cooper, J. L. *J. Am. Chem. Soc.* **1972**, *94*, 4991. (b) Ando, W.; Sonlose, H.; Akasaka, T. *Tetrahedron Lett.* **1987**, *28*, 6653. For other reports of transition-metal-assisted retro-Diels-Alder reactions, see: (c) Kawka, D.; Mues, P.; Vogel, E. *Angew. Chem., Int. Ed. Engl.* **1983**, *22*, 1003. (d) Marinetti, A.; Mathey, F. *J. Am. Chem. Soc.* **1982**, *104*, 4484. (e) Marinetti, A.; Mathey, F. *Organometallics* **1982**, *1*, 1488.

(6) Cavallito, C. J.; Bailey, J. H. *J. Am. Chem. Soc.* **1944**, *66*, 1950, 1952. (7) Frolov, A. F.; Mishenkova, E. L. *Mikrobiol. Zh. (Kiev)* **1970**, *32*, 628; *Chem. Abstr.* **1971**, *74*, 74916c.

(8) Yanagawa, H.; Kato, T.; Kitahara, Y. *Tetrahedron Lett.* **1973**, 1073. (9) Block, E.; Ahmad, S.; Catalfamo, J. L.; Jain, M. K.; Apitz-Castro, R. *J. Am. Chem. Soc.* **1986**, *108*, 7045.

(10) (a) Hirsh, A. F.; Piantadosi, C.; Irvin, J. L. *J. Med. Chem.* **1965**, *8*, 431. (b) DePaolo, J. A.; Carruthers, C. *Cancer Res.* **1960**, *20*, 431.

(11) Cunneen, J. I.; Lee, D. F. *J. Appl. Polym. Sci.* **1964**, *8*, 699.

(12) Pihl, A.; Eldjarn, L. *Pharmacol. Rev.* **1958**, *10*, 437.

(13) Kato, A.; Numata, M. *Tetrahedron Lett.* **1972**, 203.





Figure 1. Molecular structure of **13d**. S(2) is disordered; a minor site S(2') is omitted. Fe–C(8), 1.740 (7) Å; Fe–C(10), 2.008 (7) Å; C(8)–C(9), 1.503 (9) Å; C(9)–S(2), 1.823 (9) Å; S(1)–S(2), 2.082 (4) Å; S(1)–C(10), 1.824 (7) Å; C(8)–C(10), 1.330 (10) Å; S(1)–O(1), 1.456 (6) Å. CNT–Fe–C(6), 123.0 (3)°; CNT–Fe–C(7), 124.0 (3)°; CNT–Fe–C(8), 123.1 (3)°; C(9)–S(2)–S(1), 95.7 (3)°; S(2)–S(1)–C(10), 93.0 (2)°; S(2)–S(1)–O(1), 112.2 (3)°; C(10)–S(1)–O(1), 107.9 (3)°; S(1)–C(10)–C(8), 119.4 (5)°.


occurs with S₂O the proposed intermediate cationic allene complex (**9**) could close via sulfur (path a) or oxygen (path b). Control of the mode of cyclization should be effected by varying the electrophilicity of the metal M in **9**.

We have tested this idea by treating a variety of CpMe_nM(CO)_mCH₂C≡CR complexes (**12**) with S₂O equivalent **2** (Table I). Cyclization proceeded cleanly at room temperature in tetrahydrofuran to yield metalthiosulfinate esters (**13**).¹⁵ As expected, increased electron density at the metal center accelerated this reaction.^{14g} Complexes **12a** and **12b** reacted completely with a slight excess of **2** within 2 h at room temperature, whereas it was most convenient to use 2 equiv of **2** to effect complete cyclization of **12c** and **12d** in 2 h. Reactions of molybdenum complexes **12e** and **12f** were only 50% complete after 24 h. There was good mass balance in all reactions, and 2,3-diphenylbutadiene (**4**) was isolated as the organic byproduct in all cases. We see no evidence for the production of the regioisomeric oxygen cyclization product (**11**). Complexes **13a–f** are all air stable, crystalline, and chromatographable. Confirmation of proposed structure **13d** comes from X-ray crystallographic data (see Figure 1). The sulfur–sulfur (2.082 Å) and sulfur–oxygen (1.456 Å) bond lengths are consistent with other limited crystallographic data available on thiosulfinate esters.¹⁶ Due to crowding on the


Scheme II



carbon–carbon double bond, the Fe–C(8)–C(10) and C(16)–C(10)–C(8) bond angles are large, 129.5° and 131.8°, respectively.



When these metalthiosulfinate esters (**13b–d,f**) were treated with ceric ammonium nitrate under 1 atm of CO in ethanol, CO insertion and iron–carbon bond cleavage occurred.¹⁷ This produced the desired five-membered-ring thiosulfinate esters **14** in good yield (from **13c**, R = Me, 50%; from **13b**, **13d**, and **13f**, R = Ph, 66%, 75%, and 59%, respectively).



The most unusual aspect of this chemistry is that cyclization occurred at room temperature where **2** is stable in solution in the absence of the transition-metal 2-alkynyl complexes (**12**). A mechanism that accounts for the observed products can be formulated by using Wojcicki's results with SO₂ as an analogy (Scheme II).^{14c}

This mechanism requires an unprecedented mode of reactivity for thiosulfinate esters which normally undergo S–S bond cleavage when treated with carbon nucleophiles.¹⁸ One possible driving force for this nucleophile-induced ring opening would be the resonance stabilization energy gained upon conversion of **2** to **4**.

Our current efforts are centered around investigation of the reactivity of **2** with other organic as well as organometallic nucleophiles.

Acknowledgment is made to the North Carolina Board of Science and Technology, the donors of the Petroleum Research Fund, administered by the American Chemical Society, and the National Science Foundation (CHE 8817985) for support of this work. The Midwest Center for Mass Spectrometry, a National

(14) (a) Bucheister, A.; Klemarczyk, P.; Rosenblum, M. *Organometallics* **1982**, *1*, 1679. (b) Chen, L. S.; Lichtenberg, D. W.; Robinson, P. W.; Yamamoto, Y.; Wojcicki, A. *Inorg. Chim. Acta* **1977**, *25*, 165. (c) Rosenblum, M. *Acc. Chem. Res.* **1974**, *7*, 122. (d) Roustan, J.-L.; Merour, J.-Y.; Benaim, J.; Charrier, C. *C. R. Seances Acad. Sci., Ser. C* **1972**, *274*, 537. (e) Thomasson, J. E.; Robinson, P. W.; Ross, D. A.; Wojcicki, A. *Inorg. Chem.* **1971**, *10*, 2130. (f) Bannister, W. D.; Booth, B. L.; Haszeldine, R. N.; Loader, P. L. *J. Chem. Soc. A* **1971**, 930. (g) Bell, P.; Wojcicki, A. *Inorg. Chem.* **1981**, *20*, 1585.

(15) General cyclization procedure: Iron propargyl complex **12d**¹⁹ (0.200 g, 0.68 mmol) was dissolved in tetrahydrofuran (15 mL) and stirred under nitrogen at 25 °C. Dithiolenone **2** (0.390 g, 1.37 mmol) was dissolved in tetrahydrofuran (5 mL) and added dropwise to the propargyl complex. After 2 h at 25 °C, TLC analysis (silica gel, 20:1 petroleum ether/diethyl ether) showed complete disappearance of starting material (**12d**). The solvent was removed by rotary evaporation, and the crude product was purified by column chromatography (silica gel). Elution with petroleum ether yields 2,3-diphenylbutadiene (**4**), followed by elution with 10% ethanol in diethyl ether, which yields the metalthiosulfinate ester **13d** as a yellow-brown solid (0.182 g, 72%); mp 148 °C dec (acetone/petroleum ether).

(16) (a) Stuedel, R.; Luger, P.; Bradaczek, H.; Rebsch, M. *Angew. Chem., Int. Ed. Engl.* **1973**, *12*, 423. (b) Wahl, G. H.; Bordner, J.; Harpp, D. N.; Gleason, J. G. *Acta Crystallogr. B* **1973**, *29*, 2272. (c) Kiers, C. T.; Vos, A. *Recl. Trav. Chim. Pays-Bas* **1978**, *97*, 166.

(17) A procedure analogous to that reported by Reger et al. was used: Reger, D. L.; Mintz, E.; Lebiada, L. *J. Am. Chem. Soc.* **1986**, *108*, 1940. Spectral data: (**14d**) (R = Ph) mp 172 °C dec; IR (CDCl₃, cm⁻¹) 3158, 3063, 2985, 2938, 1712, 1560, 1464, 1375, 1311, 1282, 1238, 1078; ¹H NMR (CDCl₃) 7.90–7.60 (m, 5 H), 5.15 (d, J = 18 Hz, 1 H), 4.69 (d, J = 18 Hz, 1 H), 4.09 (q, J = 7 Hz, 2 H), 1.05 (t, J = 7 Hz, 3 H); HRMS FAB HFBA matrix MH⁺ calcd for C₁₂H₁₃O₃S₂ 269.0306, found 269.0312.

(18) (a) Vinkler, E.; Klivenyi, F.; Klivenyi, E. *Acta Chem. Acad. Sci. Hung.* **1958**, *16*, 747. (b) Savige, W. E.; Fava, A. *J. Chem. Soc., Chem. Commun.* **1965**, 417. (c) Kice, J. L.; Large, G. B. *Tetrahedron Lett.* **1965**, 3537. (d) Kice, J. L.; Large, G. B. *J. Am. Chem. Soc.* **1968**, *90*, 4069. (e) Vinkler, E.; Klivenyi, F. *Int. J. Sulfur Chem.* **1973**, *8*, 111. (f) Kishi, M.; Ishihara, S.; Komeno, T. *Tetrahedron* **1974**, *30*, 2135. (g) Mikolaczkyk, M.; Drabowicz, J. *J. Chem. Soc., Chem. Commun.* **1976**, 220.

(19) Roustan, J.-L.; Cadiot, P. *C. R. Seances Acad. Sci., Ser. C* **1969**, *268*, 734.

Science Foundation regional instrumentation facility (CHE 8211164), performed mass spectrometric analyses.

Supplementary Material Available: Spectroscopic and analytical data for all new compounds and tables of thermal parameters, positional parameters, bond distances, and bond angles for **13d** (5 pages). Ordering information is given on any current masthead page.

Characterization of the ESR Spectrum of the Superoxide Anion in the Liquid Phase

R. N. Bagchi,¹ A. M. Bond,^{*1} F. Scholz,² and R. Stösser²

Department of Chemical and Analytical Sciences
Deakin University, Geelong 3217
Victoria, Australia
Sektion Chemie, Humboldt-Universität
Hessische Strasse 1-2, Berlin 1040
German Democratic Republic
Received November 7, 1988

The anion radical O_2^- , which is of considerable importance in living matter,^{3,4} has been shown to be moderately stable in low-temperature aprotic media.⁵⁻⁸ Consequently, ESR studies on O_2^- ⁹⁻²³ generally have been undertaken in frozen media containing aprotic solvent⁹⁻¹⁹ or at low temperatures on various solid oxide surfaces.²⁰⁻²³ For example, Bray et al.⁹ studied the low-temperature ESR spectrum of O_2^- generated in aqueous dimethyl sulfoxide media by using pulse radiolysis followed by a rapid freezing method. The intensity of the two-line spectrum was found to be extremely temperature dependent, and the disappearance of the ESR lines at temperatures prior to melting of the solvent was attributed to line broadening resulting from fast relaxation. In contrast only brief reports at single temperatures are available on the solution phase O_2^- ESR spectrum.^{24,25}

This communication reports that ESR spectra of O_2^- can be




Figure 1. ESR spectra at various temperatures of the superoxide anion O_2^- generated by electrochemical reduction of molecular oxygen in butyronitrile containing 0.1 M Bu_4NClO_4 .

obtained in solution at temperatures well above the melting point of aprotic solvents²⁶⁻²⁸ such as butyronitrile (mp $-112.0^\circ C$) and propionitrile (mp $-92.9^\circ C$) and that frozen or solid-state media are not required for temperature-dependent studies.

Figure 1 shows the ESR spectra of electrochemically generated O_2^- in butyronitrile in the frozen solvent at $-210^\circ C$ and in the liquid phase at -105 and $-51^\circ C$. Obviously it is possible to detect the ESR signal or O_2^- as high as $60^\circ C$ above the melting point ($-112^\circ C$) of butyronitrile. A similar study was undertaken in propionitrile, and the temperature dependence of the ESR signal amplitude in this solvent is shown in Figure 2 over a temperature range encompassing frozen solvent and the liquid phase. The temperature dependence is much greater than expected on the basis of the Curie law. For comparison purposes, Figure 2 also includes the temperature-dependent behavior of the well-known stable 4-hydroxy-2,2',6,6'-tetramethylpiperidine-*N*-oxyl (TEMPO) radical.

TEMPO is a much larger molecule than O_2^- and has a much smaller g -factor anisotropy. Data for TEMPO are in complete agreement with the Curie law.

The ESR spectra of O_2^- in frozen butyronitrile and propionitrile are characterized by $g_{||} = 2.072$ (broad line) and $g_{\perp} = 2.009$ and are similar to the results reported in frozen solutions of dimethyl sulfoxide.⁵ However, there are two unusual features of the solution ($g_{av} = 2.031$) spectra: (a) the signal height decreases sharply with

- (1) Deakin University.
- (2) Humboldt-Universität.
- (3) Michelson, A. M.; McCord, J. M.; Fridovich, I. A. *Superoxide and Superoxide Dismutases*; Academic Press: London, 1977.
- (4) *Oxygen Radicals in Chemistry and Biology*; Bors, W., Saran, M., Tait, D., Eds.; Walter de Gruyter: Berlin, 1984.
- (5) Sawyer, D. T.; Chiercato, G.; Angelis, Ch. T.; Nanni, E. J.; Tsuchiya, T. *Anal. Chem.* **1982**, *54*, 1720.
- (6) Sawyer, D. T.; Yamaguchi, K.; Calder, T. S. ref 4, p 65.
- (7) Bradic, Z.; Wilkins, R. G. *J. Am. Chem. Soc.* **1984**, *106*, 2236.
- (8) *C.R.C. Handbook of Methods for Oxygen Radical Research*; Greenwald, R. W., Ed.; C.R.C. Press, Inc.: Boca Raton, FL, 1985.
- (9) Bray, R. C.; Mautner, G. M.; Fielden, E. M.; Carle, C. I. ref 3, pp 61-75.
- (10) Maricle, D. L.; Hodgson, W. G. *Anal. Chem.* **1965**, *37*, 1562.
- (11) Symons, M. C. R.; Eastland, G. W.; Denny, L. R. *J. Chem. Soc., Faraday Trans. 1* **1980**, *76*, 1868.
- (12) Sawyer, D. T.; Calderwood, T. S.; Yamaguchi, K.; Angelis, C. T. *Inorg. Chem.* **1983**, *22*, 2557.
- (13) Ansari, M. A.; Chandrasekaran, J.; Sarkar, S. *Inorg. Chem.* **1988**, *27*, 763.
- (14) Valentine, J. S.; Tatsuno, Y.; Nappa, M. *J. Am. Chem. Soc.* **1977**, *99*, 3522.
- (15) Eastland, G. W.; Symons, M. C. R. *J. Phys. Chem.* **1977**, *81*, 1502.
- (16) Green, M. R.; Hill, H. A. O.; Turner, D. R. *FEBS Lett.* **1979**, *103*, 176.
- (17) Ozawa, T.; Hanaki, A.; Yamamoto, H. *FEBS Lett.* **1977**, *74*, 99.
- (18) Bennett, J. E.; Ingram, D. J. E.; Symons, M. C. R.; George, P.; Griffith, J. S. *Phil. Mag.* **1955**, *46*, 443.
- (19) Hoare, J. P. *The Encyclopedia of the Electrochemistry of the Elements*; Bard, A. J., Ed.; Marcel Dekker: New York, 1974; Vol. II, Chapter 5, pp 191-382.
- (20) Symons, M. C. R. *Nature* **1987**, *325*, 659.
- (21) Ragai, J. *Nature* **1987**, *325*, 703.
- (22) Naccache, N.; Meriandean, P.; Che, M.; Tench, A. J. *Trans Faraday Soc.* **1971**, *67*, 506.
- (23) Shiotari, M.; Moro, G.; Feed, J. H. *J. Chem. Phys.* **1981**, *74*, 2616.
- (24) Peover, M. E.; White, B. S. *Chem. Commun.* **1965**, 183.
- (25) Slough, W. *Chem. Commun.* **1965**, 184.

(26) The ESR spectra of O_2^- were recorded on samples prepared by electrochemical reduction at platinum electrodes of molecular oxygen, O_2 , dissolved in butyronitrile and propionitrile (purified by standard procedures) containing 0.1 M Bu_4NClO_4 as the supporting electrolyte. A newly developed in situ electrochemical ESR cell²⁷ or a conventional bulk electrolysis cell was used for the reduction and for obtaining voltammetric data. ESR spectra of O_2^- also were obtained by dissolution of KO_2 , supported by dibenzo-18-crown-6 ether, in the same solvents. The ESR spectra were recorded over a wide temperature range ($-210^\circ C$ to $20^\circ C$) in both the frozen and solution phases either with a "home made" ESR spectrometer²⁸ or a Varian E4 spectrometer.

(27) Bagchi, R. N.; Bond, A. M.; Scholz, F. J. *Electroanal. Chem.* **1988**, *252*, 259.

(28) Anderson, J. E.; Bagchi, R. N.; Bond, A. M.; Greenhill, H. B.; Henderson, T. L. E.; Walter, F. L. *Am. Lab. (Fairfield, Conn)* **1981**, *13* (February), 21.