


intermediate,^{12,13} versicolorin B (4) and versicolorin A hemiacetal (10), which is reduced to 4 under cell-free conditions, should partition in ¹³C-labeled form between aflatoxin B₂ (7) and B₁ (9). Versicolorin A (5), however, should incorporate label solely into aflatoxin B₁ (9).



Racemic [4'-¹³C]averufin,¹² [9-¹³C]versicolorin B (C),¹⁴ [9-¹³C]versicolorin A hemiacetal, and [9-¹³C]versicolorin A were administered to mycelial suspensions of wild-type *A. parasiticus* (SU-1) under standard conditions.^{2,13} After quantification by HPLC analysis of the aflatoxin B₁ and B₂ produced,¹⁵ 7 and 9 were separated and further purified by preparative radial chromatography (silica gel, CHCl₃/acetone 20:1). The sites and extent of ¹³C-incorporation were determined by ¹³C{¹H} NMR spectroscopy and mass spectrometry. In each instance the locus of ¹³C-enrichment in both aflatoxin B₁ and B₂ was that expected. The levels of ¹³C-incorporation are summarized in Table I.

In *A. parasiticus* (SU-1) about 20 times more aflatoxin B₁ (9) is generated than B₂ (7). As might be predicted for an earlier pathway intermediate, i.e., averufin (1), steady-state flux through the proposed branch point metabolite versicolorin B (4) would be expected to lead to equal levels of isotopic labeling in 7 and 9, as was observed. Addition of versicolorin B (4) itself, however, perturbs that steady state and a significantly enhanced enrichment of ¹³C was observed in the minor metabolite aflatoxin B₂ (7). Entirely in keeping with the cell-free experiments above, label from the hemiacetal 10 partitioned similarly to versicolorin B, consistent with its reduction to this tetrahydrobisfuran prior to utilization in the pathway. In contrast, no label from versicolorin A (5) was detected in aflatoxin B₂ (7) indicating irreversible desaturation of versicolorin B to versicolorin A.

In conclusion, formation of the critical dihydrobisfuran is a three-step process from versiconal acetate (3) involving an acetate hydrolysis/cyclization sequence to give versicolorin B (4) and an apparent oxidative desaturation step to irreversibly form versicolorin A (5). Both 1'-hydroxyversicolorone (2) and versiconal acetate (3) are isolated as racemates.¹⁶ Cell-free experiments have demonstrated, however, that versiconal acetate (3) can be completely converted to (-)-versicolorin B (4) suggesting that either a single, bifunctional enzyme or two enzymes selectively process one enantiomer of 3 to establish the absolute configuration of the critical bisfuran ring system in 4. The stereoelectronic arguments made earlier,¹⁷ based on the absolute configuration of averufin (1),¹⁸ would give 2 and 3 in the correct configuration for conversion to bisfurans but fail, however, to account for the competing facility of these intermediates to racemize relative to their rates of forward reaction in the biosynthetic pathway. Once formed, separate pathways diverge from versicolorin B (4) to give rise to the tetrahydro- and dihydrobisfuran-containing aflatoxins 7 and 9. The hydroxylated species 10, which might be imagined to be involved in the oxidation of 4 to 5, does not undergo de-

hydration under either cell-free or whole-cell conditions, but is reduced to versicolorin B.¹⁹

Acknowledgment. We are grateful to the National Institutes of Health for financial support of this research (ES 01670). Major analytical instrumentation used was acquired with funds from the NIH and NSF (NMR: RR01934 and PCM 83-03176; MS: RR 02318).

(19) Hydration, enzymic or chemical, of dihydrobisfurans followed by reduction to the tetrahydro series may provide a minor route for conversion between these bisfuran groups.³ The results in Table I indicate that this is not a significant process for versicolorin A or for later dihydrobisfuran-containing intermediates in the pathway under the present experimental conditions.

Regioselectivity and Diastereoselectivity in Free-Radical Macrocyclization

Ned A. Porter,* Brigitte Lacher, Vincent H.-T. Chang, and David R. Magnin

Department of Chemistry, Duke University
Durham, North Carolina 27706

Received July 17, 1989

The development of free-radical methods for the synthesis of complex organic molecules continues to attract much attention.¹ Construction of five-membered rings has been one of the cornerstones of recent synthetic developments, but the synthesis of large rings by free-radical cyclization has also recently been reported.² Macrocyclic yields can be as high as 90% for secondary or tertiary radicals if the alkene is electron deficient. Steric effects have a profound effect on the reaction.^{3,4} We report here on new substrates, 1a-d, for macrocyclization that have X = CH₂ and Z = COOEt or CONR₂ (Scheme 1). Cyclizations of these compounds are highly selective for the endocyclic product. Furthermore, substrates where Z is a chiral amide derived from alanine, 1c and 1d, give products that are highly enriched in one of the four possible diastereomeric cyclic products.

The synthesis of the substrates 1a-d is straightforward, the key step being phosphonate coupling to form the electrophilic alkene.⁵ For compounds 1b-d, for example, the coupling reaction between Br(CH₂)_nCOCH₂PO(OEt)₂⁶ and the α dicarbonyl HCOCONR₂ gave Br(CH₂)_nCOCH=CHCONR₂ (70%), which was converted to the iodide by reaction with NaI (quantitative). The substrate 1a was prepared via the sequence Br(CH₂)₁₂COCHO + EtOOCCH₂PO(OEt)₂ → Br(CH₂)₁₂COCH=CHCOEt (70%) → 1a (quantitative).

Mixtures of cis and trans geometric isomers were sometimes formed in the coupling reaction, and, if formed, the cis isomers were converted to the trans by reaction with I₂. The dicarbonyl HCOCONR₂ was prepared by reaction of the nitrate O₂NOCH₂CONR₂ with NaOAc in DMSO;⁷ Br(CH₂)₁₂COCHO was prepared by Swern oxidation of Br(CH₂)₁₂CHOHCH₂OH.⁸ The pyrrolidine used for the amide of 1c and 1d was prepared by the method of Schlessinger, and the starting material used in this synthesis was the unnatural (R)-alanine since cyclization

(12) Townsend, C. A.; Christensen, S. B.; Davis, S. G. *J. Am. Chem. Soc.* **1982**, *104*, 6152-6153. Simpson, T. J.; deJesus, A. E.; Steyn, P. S.; Vlegaar, R. *J. Chem. Soc., Chem. Commun.* **1982**, 631-632. Townsend, C. A.; Davis, S. G. *J. Chem. Soc., Chem. Commun.* **1983**, 1420-1422.

(13) Townsend, C. A.; Christensen, S. B.; Davis, S. G. *J. Chem. Soc., Perkin Trans. 1* **1988**, 839-861.

(14) Reduction of 11 with LiAlH₄ and protection of the resulting alcohol gave 12, which was converted in 45% overall yield to versicolorin C (the racemate of versicolorin B) essentially as described for versicolorin A (see preceding paper).

(15) Modification of the method of Schmidt et al. (Schmidt, R.; Mondani, J.; Ziegenhagen, E.; Dose, K. *J. Chromatogr.* **1981**, *207* 435-438).

(16) Townsend, C. A.; Placvan, K. A.; Pal, K.; Brobst, S. W.; Irish, M. S.; Ely, E. W., Jr.; Bennett, J. W. *J. Org. Chem.* **1988**, *53*, 2472-2477, and references cited.

(17) Townsend, C. A. *Pure Appl. Chem.* **1986**, *58*, 227-238. Townsend, C. A.; Christensen, S. B. *Tetrahedron* **1983**, *39*, 3575-3582.

(18) Koreeda, M.; Hulin, B.; Yoshihara, M.; Townsend, C. A.; Christensen, S. B. *J. Org. Chem.* **1985**, *50*, 5426-5428.

(1) (a) Hart, D. *J. Science* **1984**, *223*, 883. (b) Curran, D. P. *Synthesis* **1988**, 489. (c) Dowd, P.; Choi, S.-C. *J. Am. Chem. Soc.* **1987**, *109*, 3493. (d) Curran, D. P.; Chang, C.-T. *Tetrahedron Lett.* **1987**, *28*, 2477. (e) Hart, D. J.; Seely, F. L. *J. Am. Chem. Soc.* **1988**, *110*, 1631. (f) RajanBabu, T. V.; Fukunaga, T. *J. Am. Chem. Soc.* **1989**, *111*, 296.

(2) (a) Porter, N. A.; Magnin, D. R.; Wright, B. T. *J. Am. Chem. Soc.* **1986**, *108*, 2787. (b) Porter, N. A.; Chang, V. H.-T.; Magnin, D. R.; Wright, B. T. *J. Am. Chem. Soc.* **1988**, *110*, 3554.

(3) Giese, B. *Radicals in Organic Synthesis: Formation of Carbon-Carbon Bonds*; Pergamon Press: New York, 1986.


(4) Porter, N. A.; Chang, V. H.-T. *J. Am. Chem. Soc.* **1987**, *109*, 4976. (5) See, for example, Roush, W. R.; Coe, J. W. *Tetrahedron Lett.* **1987**, *28*, 921.

(6) Motoyoshiya, J.; Miyajima, M.; Hirakawa, K.; Kakurai, T. *J. Org. Chem.* **1985**, *50*, 1326.

(7) Kornblum, N.; Frazier, H. W. *J. Am. Chem. Soc.* **1966**, *88*, 865.

(8) Ireland, R. E.; Norbeck, D. W. *J. Org. Chem.* **1985**, *50*, 2198.

Scheme I

Table I. Product Ratio of Cyclization Products from Iodides 1a-d^a

iodide	exo, 4	endo, 3
1a	<2 (14)	>98 (15)
1b	1 (13)	10 (14)
1c ^b	exo 1:exo 2 (13)	endo R:endo S (14)
1d ^b	exo 1:exo 2 (14)	endo R:endo S (15)
1e ^c	1 (15)	2.5 (16)

^aRing size shown in parentheses. ^bTwo diastereomeric endocyclic products and two diastereomeric exocyclic products are formed. The product ratio of the four diastereomers is given here. The configuration of the newly formed stereocenter is indicated. ^cReference 4.

reactions with this auxiliary give natural (-)-(*R*)-muscone for comparison.⁹ All new compounds were fully characterized by spectroscopy and elemental analysis.


Cyclization of the primary iodides was carried out under standard tin hydride conditions.² In a typical cyclization, described here for 1a, a 4.5 mM solution of the iodide in refluxing benzene was reacted with 1.1 equiv of Bu₃SnH and 0.1 equiv of the azo initiator AIBN. After 3 h of reaction and purification on silica gel, the mixture yielded 57% cyclic compounds that included endocyclic and exocyclic products. For 1a, only endocyclic product was isolated, and, if formed, the exocyclic product was present in the product mixture in less than 2%. Endocyclic products could be distinguished from the exocyclic products by mass spectral fragmentation that gives a characteristic loss of CH₂COOR or CH₂CONR₂ from the molecular ion for the exocyclic products.¹⁰ Yields of cyclic products for the amide derivatives 1b-d (40-45%) are lower than for the ester.

For the pyrrolidine amides 1c and 1d, four cyclic products are possible: two exocyclic diastereomers and two endocyclic compounds. All four diastereomers can be separated by HPLC on silica gel (20% ethyl acetate/hexane), and the exocyclic and endocyclic compounds can be readily assigned by mass spectral fragmentation, *vide supra*. The absolute stereochemistry of the exocyclic compounds 4c and 4d has not been assigned, since they are minor products and since they form in a 1:1 ratio. The stereochemistry of the two endocyclic compounds was established by conversion to the natural product muscone as described below. The product distribution of cyclic products formed for the four substrates 1a-d is presented in Table I.

The regioselectivity and diastereoselectivity shown in the cyclization reactions of the iodides 1a-d is striking. It seems reasonable to suggest that the balance of polar effects of the ketone vs the ester or amide groups substituting the olefin in 1a-d pro-

motes endocyclization compared to the fumarate derivative, 1e,⁴ but intermolecular additions of radicals to olefins that are models for the substrates 1c and 1d show a 1:1 ratio of regioisomers.¹¹ The preference for endocyclization in these substrates must therefore arise from the intramolecular nature of these reactions, and substrates where X = CH₂ appear to have an enhanced endocyclization bias compared to those where X = O.⁴

The diastereoselectivity observed in the endocyclization products of cyclizations of 1c and 1d can be understood by analysis of the preferred conformation of the amide auxiliary and the vector of approach of the radical in the addition reaction.¹¹ The preferred conformation of the amide requires that the nucleophilic radical approach the olefin on a vector over the pyrrolidine, thus protecting one face of the olefin from addition. In support of the model, we note that addition to the end of the olefin remote from the amide occurs without diastereoselectivity.



The keto ester 3a, formed by endocyclization of 1a, has been converted to racemic muscone. Thus, the ketone was converted to the ketal by reaction with 2,2-dimethylpropanediol, the ester was reduced to the alcohol 7 with lithium aluminum hydride, and 7 was deoxygenated under Barton conditions.¹² The ketal resulting from deoxygenation was deprotected, to give material that was identical with muscone in every respect. The configuration of the newly formed stereocenter in the macrocyclizations of 1c and 1d was determined by conversion of the pyrrolidine amide group of 1d to the methyl group of muscone. The pyrrolidine amide was particularly difficult to hydrolyze or reduce, and only by reduction of the ketone and alcohol-assisted acid hydrolysis could it be removed.¹³ The lactone 8 was formed in this sequence, and this lactone was converted to (-)-(*R*)-muscone by standard methodology.¹⁴ A mixture of stereoisomeric alcohols was formed in the borohydride reduction of 3d, and the mixture of stereoisomers was carried through the sequence. The stereochemistries of products derived from 1c are assigned by analogy to those from 1d.

The results of the studies on macrocyclization reported here make it clear that alkene amide auxiliaries provide significant stereoselectivity in radical addition reactions and that the development of models to understand these selectivities is worthwhile. In the accompanying communication, intermolecular reactions are described that are analogous to the intramolecular reactions reported here and a model that accounts for the observed stereoselectivity is presented.

Acknowledgment. This work was supported by grants from the NSF and NIH. N.A.P. gratefully acknowledges the Humboldt Foundation for the award of a Senior Fellowship and Professors B. Giese and R. Polniaszek for helpful discussions.

Supplementary Material Available: Details for the synthesis of 1, 3, and 4 (9 pages). Ordering information is given on any current masthead page.

(11) Porter, N. A.; Scott, D. M.; Lacher, B.; Giese, B.; Zeitz, H. G.; Lindner, H. J. *J. Am. Chem. Soc.*, following paper in this issue.

(12) Barton, D. H. R.; McCombie, S. W. *J. Chem. Soc., Perkin Trans. I* **1975**, 1574.

(13) Helmchen, G.; Nill, G.; Flockerzi, D.; Schuhle, W.; Youssef, M. S. *K. Angew. Chem., Int. Ed. Engl.* **1979**, *18*, 62.

(14) The conversion of 3d to (-)-(*R*)-muscone was achieved by the following steps: (a) NaBH₄; (b) H₂SO₄/H₂O/dioxane; (c) LAH; (d) *t*-BuPh₂SiCl; (e) pyridinium chlorochromate; (f) H⁺/2,2-dimethylpropanediol; (g) Bu₄N⁺F⁻; (h) Cl(C=S)OPh; (i) Bu₃SnH; (j) HCl/H₂O. The yield for each step was in excess of 70%, and all products were fully characterized by spectroscopy and elemental analysis or exact mass spectra.

(9) Schlessinger, R. H.; Iwanowicz, E. J. *Tetrahedron Lett.* **1987**, *28*, 2083.
 (10) Dubay, G. R.; Chang, V.; Porter, N. A., Presented at the ASMS Meeting, June 1988.