

Scheme III. Hydroboration of the exocyclic methylene function (C-16) with 9-BBN occurred stereospecifically from the  $\beta$ -face of the double bond and on oxidative workup (3 N NaOH/H<sub>2</sub>O<sub>2</sub>/40 °C/2 h) provided the triol **11** in 85% yield. Direct hydroboration of the  $\beta$ -dicarbonyl compound **9** with 9-BBN furnished the triol **11**; however, the yield was somewhat lower. As shown in Scheme III (structure **11**), 9-BBN has attacked the  $\beta$ -face of the exocyclic methylene function in order to minimize steric repulsion (1,3) between the axial N<sub>b</sub>-methyl function and the incoming hydroboration reagent. This is opposite to the stereochemical outcome of the hydroboration at C-16 observed during the synthesis of ( $\pm$ )-suaveoline.<sup>4b</sup>

The optically active triol (-)-**11** was regioselectively cyclized to the desired (-)-tetrahydroalstonerine monol **12** on stirring with tosyl chloride (1 equiv) in pyridine followed by the addition of either triethylamine or potassium hydroxide. This process gave (-)-**12** in 60% yield, accompanied by starting triol **11** (33%), which could be recycled to provide additional quantities of (-)-tetrahydroalstonerine **12**. When monol **12** was stirred with pyridinium dichromate, an 86% yield of (-)-dihydroalstonerine (**3b**) was realized; however, treatment of **12** under modified Swern<sup>9</sup> conditions [(COCl)<sub>2</sub>/DMSO/CH<sub>2</sub>Cl<sub>2</sub>/-78 °C  $\rightarrow$  -10 °C/1.5 h; Et<sub>3</sub>N] gave (-)-alstonerine (**2a**) (mp 171–172 °C) in 51% yield, accompanied by dihydroalstonerine (**3b**) (31%). The spectral data for (-)-**2a** (<sup>1</sup>H NMR, <sup>13</sup>C NMR, <sup>10</sup>IR, UV, MS)<sup>2a</sup> were in com-

plete agreement with those published for natural (-)-alstonerine (mp 172–173 °C); moreover, the optical rotation {[ $\alpha$ ]<sub>D</sub><sup>25</sup> -190° (c 0.32, EtOH)} of synthetic **2a** indicates that it has been prepared in at least 98% ee.

The synthesis described above represents the first chirally controlled preparation of a member of the macroline-related alkaloids.<sup>11</sup> The stereospecific preparation of tetracyclic ketone **5a**,<sup>4a,b</sup> coupled with the execution of both the Claisen rearrangement (C-15) and the hydroboration process (C-16) in the desired fashion, provides a route for the enantiospecific synthesis of the macroline/sarpagine alkaloids. Further work is in progress to extend this approach to the synthesis of alstophylline (**2b**), as well as a number of bisindole alkaloids,<sup>1–3</sup> including the hypotensive bisindole alkaloid macralstonine (**1**).<sup>1a,b</sup>

**Note Added in Proof.** Recently, base-catalyzed (NaOMe, CH<sub>3</sub>OH,  $\Delta$ ) epimerization of synthetic **3b** gave the epimeric **3a** which had been previously converted into macroline **4** by LeQuesne et al.<sup>3a</sup> Consequently, the synthesis of (-)-**3b** also constitutes a formal total synthesis of **4**, although the yield of this conversion has not been maximized.

(10) Ratnayake, C.; Lakshmi, S. R.; Arambewela, K. T.; Silva, D.; Rahman, A.; Alvi, K. A. *Phytochemistry* **1987**, *26*, 868.

(11) The optical purity (>97% ee) of **5a** as well as other intermediates was proven by the use of chiral shift reagents as detailed in the following: Campbell, J. *Aldrichimica Acta* **1972**, *5*, 2. All compounds including (-)-**2a** gave satisfactory NMR, IR, and mass spectral data.

(9) Mancuso, A. J.; Huang, S.-L.; Swern, S. *J. Org. Chem.* **1978**, *43*, 2480.

## Additions and Corrections

**Phosphate Ester and Phosphinate Binding to the ( $\mu$ -Oxo)diiron(III) Core: Synthesis and Characterization of [Fe<sub>2</sub>O(O<sub>2</sub>P(OC<sub>6</sub>H<sub>5</sub>)<sub>2</sub>)<sub>2</sub>(HBpz<sub>3</sub>)<sub>2</sub>] and [Fe<sub>2</sub>O(O<sub>2</sub>P(C<sub>6</sub>H<sub>5</sub>)<sub>2</sub>)<sub>2</sub>(HBpz<sub>3</sub>)<sub>2</sub>] [*J. Am. Chem. Soc.* **1990**, *112*, 681–690]. PETRA N. TUROWSKI, WILLIAM H. ARMSTRONG, MARY E. ROTH, and STEPHEN J. LIPPARD\***

Page 687: The minus sign in eq 2 should be a plus sign. This change does not affect any results of the paper, for which the correct equation was used.

**Characterization of (Methylcyclopentadienyl)trimethylplatinum and Low-Temperature Organometallic Chemical Vapor Deposition of Platinum Metal** [*J. Am. Chem. Soc.* **1989**, *111*, 8779]. ZILING XUE, M. JANE STROUSE, DAVID K. SHUH, CAROLYN B. KNOBLER, HERBERT D. KAESZ,\* ROBERT F. HICKS, and R. STANLEY WILLIAMS

Page 8780: We have learned of new evidence from NOESY spectra that suggests that the assignment of H<sub>a</sub> and H<sub>b</sub> (Figure 1) and C<sub>a</sub> and C<sub>b</sub> (Figure 2) in (MeCp)PtMe<sub>3</sub> should be reversed (private communication from Richard A. Newmark, Larry D. Boardman, and Allen R. Siedle, 3M Corporate Research Laboratories, Bldg. 201-BS-05, Box 33221, St. Paul, MN 55144-1000). Arguments and supporting data that involve a series of compounds

including the one mentioned above are being prepared for publication.

**X-ray Structures of Cubylcubane and 2-tert-Butylcubylcubane: Short Cage-Cage Bonds** [*J. Am. Chem. Soc.* **1988**, *110*, 7232]. R. GILARDI,\* M. MAGGINI, and P. E. EATON

Page 7232, footnote 3: the *c* dimension should be 13.431 (1) Å rather than 13.341 (1) Å.

**Mechanism of Grignard Reagent Formation. The Surface Nature of the Reaction** [*J. Am. Chem. Soc.* **1989**, *111*, 1896]. H. M. WALBORSKY\* and JANUSZ RACHON

Page 1896: The label for structure **4** should read (S)-(+)-**4**.

**Evidence for a 1,2-Fluoride Shift in a Gaseous Cation** [*J. Am. Chem. Soc.* **1989**, *111*, 6868]. THOMAS A. SHALER and THOMAS HELLMAN MORTON\*

Page 6869, Table I: The first entry should be -216.227 24 au for ion **5**. Footnote *b* should refer to the following reference: Stams, D. A.; Thomas, T. D.; Maclaren, D. C.; Ji, D.; Morton, T. H. *J. Am. Chem. Soc.* **1990**, *112*, 1427–1434.

## Computer Software Reviews

**Material Safety Data Sheets on CD-Rom.** Sigma/Aldrich: 940 West St. Paul Ave., Milwaukee, WI 53233. \$1300 annual subscription rate.

Material Safety Data Sheets<sup>1</sup> on CD-ROM is a single compact disk that provides access to over 38 000 complete MSDSs including chemical

(1) A Material Safety Data Sheet, MSDS, includes a detailed description of a chemical's physical properties, fire, explosive, and reactivity data, health hazard and toxicity information, required protective equipment, spill procedures, and waste disposal methods.

structure with printing options. This CD-ROM and its companion software run on an IBM PC or Apple Macintosh computer equipped with CD-ROM drive. Hardware necessary for using this software, including CD-ROM Drive, IBM Cable and Interface Kit, or Apple Macintosh Cable Kit, can be purchased through Aldrich. For Macintosh computers Finder 6.1 and System 6.0.2 or later versions must be used. The CD-ROM provides information on the majority of Sigma/Aldrich chemicals (industries or universities with multiple users in need of quick access to chemical handling and safety information can be accommodated by