

Asymmetric Induction via a Catalytic Antibody<sup>1</sup>Shoji Ikeda,<sup>2</sup> Michael I. Weinhouse, Kim D. Janda,\* and Richard A. Lerner\*

The Scripps Research Institute  
Department of Molecular Biology and Chemistry  
10666 North Torrey Pines Road  
La Jolla, California 92032

Samuel J. Danishefsky\*

Department of Chemistry, Yale University  
New Haven, Connecticut 06511  
Received May 21, 1991

Recently it was shown that racemic phosphonate haptens can induce catalytic antibodies with either *R* or *S* substrate specificity, thereby accomplishing kinetic resolution.<sup>3</sup> As the next step, we chose to challenge a catalytic antibody with the possibility of enantioselective perturbation of a meso substrate. It was of particular interest to direct antibody-induced lipase-like action toward meso compound **1** (Figure 1).<sup>4</sup> It will be recalled that the corresponding *S* alcohol **2**, previously generated from the action of the electric eel acetylcholinesterase on **1**,<sup>5</sup> serves as a starting material for a concise stereospecific total synthesis of prostaglandin F<sub>2α</sub><sup>6</sup> and congeners thereof (Figure 1).<sup>7</sup>

In undertaking this application, we would be implementing significant extensions of catalytic antibody methodology.<sup>8</sup> First we would be operating without any anchoring aryl substituents in the hapten **3** (Scheme I). Decreased immunogenicity was thus anticipated.<sup>9</sup> Also, the screening protocols would not benefit from a ready spectrophotometric assay.

We sought to apply compound **3** as a transition-state model. Its synthesis (Scheme I) in enantiomerically pure form posed an interesting question of feasibility. There had been concern that the deprotonation of **2** would be accompanied by racemization resulting from acyl migration.<sup>10</sup> In the event, under the conditions that were developed (Scheme I), no racemization occurred as assayed via <sup>19</sup>F NMR of the crude Mosher ester **4**.

Phosphonate **3** was coupled to carrier proteins BSA (bovine serum albumin) and KLH (keyhole limpet hemocyanin) via EDC (1-(3-(dimethylamino)propyl)-3-ethyl-carbodiimide hydrochloride) coupling methodology. Mice were immunized with the KLH conjugate of **3**, and antibodies were generated by standard protocols.<sup>11</sup> Immunization of **3** produced 33 antibodies of the IgG class. The abundance of acetyl- and butyrylcholinesterases in red blood cells and serum<sup>12</sup> necessitated extra caution during antibody purification. Ascites fluid was purified via anion-exchange chromatography (DEAE) followed by affinity chromatography (protein G) and then again by anion-exchange chromatography (Mono Q, Pharmacia). As a control, authentic acetyl- and bu-





Figure 1. Meso substrate **1**; **2**, product from abzyme or enzyme reaction described and starting material for the synthesis of prostaglandin F<sub>2α</sub>.

Scheme I<sup>a</sup>

<sup>a</sup> (a) P(OCH<sub>3</sub>)<sub>3</sub>, *p*-cymene, 170 °C, 17 h, 62%. (b) (COCl)<sub>2</sub>, CH<sub>2</sub>Cl<sub>2</sub>, room temperature, 6 h, 100%. (c) (1*R*,4*S*)-(+)-4-Hydroxy-2-cyclopentenyl acetate, LDA, THF, -78 to 0 °C, 2 h, 55%. (d) TFA, CH<sub>2</sub>Cl<sub>2</sub>, room temperature, 1 h, -90%. (e) *tert*-Butylamine, 50 °C, 2 weeks, 60%. (f) NaOMe, MeOH, 0 °C, 3 h, 100%. (g) (*R*)-(+)- $\alpha$ -(Trifluoromethyl)phenylacetyl chloride, pyridine, CH<sub>2</sub>Cl<sub>2</sub>, -30 °C to room temperature, 3 h, 85%. (h) *tert*-Butylamine, 50 °C, 2 weeks, 44%.

Table I. Kinetic Parameters for the Hydrolysis of *cis*-3,5-Diacetoxycyclopent-1-ene (**1**) to (1*R*,4*S*)-(+)-4-Hydroxy-2-cyclopentenyl Acetate (**2**)<sup>a</sup>

antibody/enzyme	<i>K</i> <sub>m</sub> , 10 <sup>-6</sup> M	<i>K</i> <sub>cat</sub> , min <sup>-1</sup>	<i>K</i> <sub>i</sub> , <sup>e</sup> 10 <sup>-6</sup> M
37E8 <sup>b</sup>	177	0.007	7.0 <sup>f</sup>
EC 3.1.1.7 (acetyl) <sup>c</sup>	620	250	630
EC 3.1.1.8 (butyryl) <sup>d</sup>	830	7.0	800

<sup>a</sup> Conditions: <sup>b</sup> Assay conditions (pH 8.0, ATE, [0.1 M *N*-(2-acetamido)-2-aminoethanesulfonic acid (ACES), 0.052 M Tris, 0.052 M ethanolamine], 10  $\mu$ M antibody, 37 °C; under these conditions background hydrolytic rate  $8 \times 10^{-5}$  min<sup>-1</sup>. <sup>c</sup> Type V-S, electric eel, Sigma, 1000–2000 units/mg of protein, used 0.059 units. <sup>d</sup> From horse serum, Sigma, 500–1000 units/mg of protein, used 0.0117 units. All assays consisted of determining **2** by GC analysis (Hewlett-Packard 5890 A, Carbowax 30m, 0.53-mm bore, capillary column) with ethyl levulinate as an internal standard. <sup>e</sup> Inhibition data determined with **3** as the inhibitor. <sup>f</sup> Determined by  $S = K_m, K_i = I$ , with velocity equivalent to  $1/3 V_{max}$ .

tylcholinesterases were shown *not* to be retained on our affinity column under the conditions employed for antibody purification. Antibodies were judged to be homogeneous (>98%) by sodium dodecyl sulfate polyacrylamide gel electrophoresis.<sup>13</sup>

Antibody-mediated production of **2** was initially screened by using 20  $\mu$ M antibody and 1 mM diacetate **1** (Table I). One of the 33 antibodies, 37E8, was found to be catalytic. The initial rate of hydrolysis of **1** catalyzed by 37E8 (10  $\mu$ M) followed Michaelis–Menten kinetics (Table I). The antigen **3** was a potent, tight-binding inhibitor of the reaction (Table I). Tight-binding inhibition, common among enzymes,<sup>14</sup> had previously been observed in an abzyme system.<sup>15</sup>

Our conclusion that the antibody is “contaminase” free is based not only on electrophoresis results but also on the large disparity seen between the abzyme and enzyme *K*<sub>i</sub>'s (Table I). If a cholinesterase were present in the antibody solution under the ex-

(1) This research was supported in part by grants from the National Institutes of Health (GM-43858, K.D.J. and HL-25848, S.J.D.). We thank Dr. Ken Chow for providing us with optically pure **2**.

(2) Visiting scientist on leave from Kanebo, Ltd., Osaka, Japan, 1989–1991.

(3) (a) Janda, K. D.; Benkovic, S. J.; Lerner, R. A. *Science (Washington, D.C.)* **1989**, *244*, 437. (b) Napper, A. D.; Benkovic, S. J.; Tramontano, A.; Lerner, R. A. *Science (Washington, D.C.)* **1987**, *237*, 1041. (c) Pollack, S. J.; Hsui, P.; Schultz, P. G. *J. Am. Chem. Soc.* **1989**, *111*, 5961.

(4) (a) Kaneko, C.; Sugimoto, S.; Tanaka, S. *Synthesis* **1974**, 876. (b) Deardorff, D. R.; Matthews, A. J.; McMeekin, D. S.; Craney, C. L. *Tetrahedron Lett.* **1985**, *26*, 5615.

(5) Deardorff, D. R.; Matthews, A. J.; McMeekin, D. S.; Craney, C. L. *Tetrahedron Lett.* **1986**, *27*, 1255.

(6) Danishefsky, S. J.; Cabel, M. P.; Chow, K. *J. Am. Chem. Soc.* **1989**, *111*, 3456.

(7) Chow, K.; Danishefsky, S. J. *J. Org. Chem.* **1989**, *54*, 6016.

(8) For a comprehensive review on catalytic antibodies, see: Lerner, R. A.; Benkovic, S. J.; Schultz, P. G. *Science (Washington, D.C.)* **1991**, *252*, 659.

(9) Tijssen, P. In *Practice and Theory of Enzyme Immunology*; Burdon, R. H., vanKnippenberg, P. H., Eds.; Elsevier: New York, 1987, pp 39–41.

(10) Under some conditions, such acyl migrations were encountered in one of our laboratories, unpublished results, K. Chow, Yale University.

(11) (a) Kohler, G.; Milstein, C. *Nature (London)* **1975**, *256*, 495. (b) Engvall, E. *Methods Enzymol.* **1980**, *70*, 419.

(12) (a) Stedman, E.; Easson, L. H. *Biochem. J.* **1932**, *26*, 2056. (b) Alles, G. A.; Hanes, R. C. *J. Biol. Chem.* **1940**, *133*, 375.

(13) Laemmli, V. *Nature (London)* **1970**, *227*, 680.

(14) Williams, J. W.; Morrison, J. F. *Methods Enzymol.* **1979**, *63*, 437.

(15) Janda, K. D.; Schloeder, D.; Benkovic, S. J.; Lerner, R. A. *Science (Washington, D.C.)* **1988**, *241*, 1199.

perimental conditions employed,<sup>16</sup> our data (Table I) would dictate virtually no inhibition. The finding, where *extensive inhibition* was observed (greater than 60%), further supports the proposition that the lipase-like activity of the "abzyme" is not a consequence of contamination. In addition, the antibody (20 μM) shows minimal inactivation (less than 20% when 2 is 400 μM) in the presence of the powerful acetylcholinesterase inhibitor diisopropyl phosphorofluoridate (DFP) (50 μM).<sup>17</sup>

Enantioselectivity of the antibody-catalyzed reaction was investigated by GC analysis of a large-scale reaction of 37E8 (40 μM) in ATE (ACES, Tris, ethanolamine), pH 8.0, containing 200 μM 1. At selected time intervals an aliquot of the reaction mixture was removed (600 μL), extracted twice with a 50/50 mixture of ethyl ether/ethyl acetate, and injected onto a microcapillary gas chromatographic column (Chrompack, CP-(optically pure)-cyclodextran-B-236-M-19). At 8 h, 60 μM 2 was detected with enantioselectivity in excess of 86% for the (1R,4S)-(+)-4-hydroxy-2-cyclopentenyl acetate (2). After 14 h, 100 μM 2 with an ee of 84% was detected. These results, while gratifying, appear to be limited only by the catalytic activity of abzyme 37E8 rather than by an inherent lack of enantiotopic group differentiation.

A comparison of abzyme 37E8 to acetyl- and butyrylcholinesterase (Table I) provided us with a crude index for an abzyme-enzyme comparison. Moreover, it provides us with alternative methods to check antibody purity. A comparison of  $k_{cat}$  (Table I) shows abzyme 37E8 to be some 3-4 orders of magnitude less potent catalytically than the cholinesterases. However, the inherent enantio group selectivity of 37E8 is excellent (>98% ee).<sup>18</sup>

The abzyme obtained here is admittedly somewhat primitive compared to its enzymatic counterparts. However, these findings suggest that asymmetric synthesis with catalytic antibodies could well provide attractive opportunities in organic synthesis. Further applications along these lines are envisioned.

(16) The following experiment was performed: Catalytic antibody 37E8 (10 μM) was added to a solution of 800 μM 1 and 25 μM 3. These conditions were chosen so that the concentration of substrate would be at saturation for the antibody (4.5  $K_m$ ) and approximately  $K_m$  for either cholinesterase.

(17) Wilson, B. W.; Walker, C. R. *Proc. Natl. Acad. Sci. U.S.A.* 1974, 71, 3194. In contrast, acetylcholinesterase (2 nM) hydrolysis of 2 (400 μM) was completely inhibited by 50 μM DFP.

(18) Calculated values, which was approximated by correcting for the unwanted antipode (1S,4R)-(-)-4-hydroxy-2-cyclopentenyl acetate, which came from the competing background hydrolytic reaction.

## Quinone/Hydroxide Ion Induced Oxygenation of *p*-Benzoquinone to Rhodizonate Dianion (C<sub>6</sub>O<sub>6</sub><sup>2-</sup>) Accompanied by One-Electron Reduction to Semiquinone Radical Anion

Shunichi Fukuzumi\* and Tomohiro Yorissue


Department of Applied Chemistry, Faculty of Engineering  
Osaka University, Suita, Osaka 565, Japan

Received June 21, 1991

Revised Manuscript Received July 26, 1991

Hydroxide ion is regarded as a better electron donor as well as a stronger base in an aprotic solvent such as acetonitrile than in water, since the solvation energy for OH<sup>-</sup> in an aprotic solvent is much less than in water.<sup>1,2</sup> Thus, OH<sup>-</sup> in aprotic solvents has been reported to act as an electron donor toward quinones and other electron acceptors, since the radical anions of electron acceptors are often formed in the presence of OH<sup>-</sup> in aprotic solvents.<sup>2-6</sup> The oxidized species of OH<sup>-</sup> has been presumed to be

Scheme I



H<sub>2</sub>O<sub>2</sub><sup>2-8</sup> when no substitution reaction takes place.<sup>9</sup> However, no oxidized products of OH<sup>-</sup> (H<sub>2</sub>O<sub>2</sub> or O<sub>2</sub>) have so far been definitely identified in the one-electron reduction of electron acceptors in the presence of OH<sup>-</sup> in aprotic solvents, the mechanism of which remains to be solved.<sup>2-6</sup> This study reports that various quinones are readily reduced in the presence of Me<sub>4</sub>N<sup>+</sup>OH<sup>-</sup> to yield the corresponding semiquinone radical anions and that the one-electron reduction of *p*-benzoquinone (Q) to Q<sup>•-</sup> is accompanied by the oxygenation of Q to rhodizonate dianion (C<sub>6</sub>O<sub>6</sub><sup>2-</sup>), which is the 10-electron-oxidized species of Q.

Upon mixing of *p*-benzoquinone (Q) with Me<sub>4</sub>N<sup>+</sup>OH<sup>-</sup> in deaerated MeCN at 298 K, Q is readily reduced to Q<sup>•-</sup> (λ<sub>max</sub> = 422 nm), and the yield based on the amount of OH<sup>-</sup> is 100%.<sup>10</sup> The Q<sup>•-</sup> thus formed is very stable in deaerated MeCN at 298 K. Similarly, the reactions of various quinones with Me<sub>4</sub>N<sup>+</sup>OH<sup>-</sup> yield the corresponding semiquinone radical anions.<sup>11</sup> No oxidized products of OH<sup>-</sup> (H<sub>2</sub>O<sub>2</sub> or O<sub>2</sub>) have, however, been formed in the one-electron reduction of Q in the presence of OH<sup>-</sup> in deaerated MeCN.<sup>12</sup> No oxidized products derived from the solvent (e.g., succinonitrile)<sup>13</sup> have been detected, either.<sup>14</sup> Then, the question

(4) (a) Ballester, M. *Pure Appl. Chem.* 1967, 15, 123. (b) Ballester, M. *Acc. Chem. Res.* 1985, 18, 380. (c) Ballester, M. *Adv. Phys. Org. Chem.* 1989, 25, 267. (d) Ballester, M.; Pascual, I. *J. Org. Chem.* 1991, 56, 841.

(5) (a) Endo, T.; Miyazawa, T.; Shihashi, S.; Okawara, M. *J. Am. Chem. Soc.* 1984, 106, 3877. (b) Arudi, R. L.; Allen, A. O.; Bielski, B. H. *J. FEBS Lett.* 1981, 135, 265. (c) Umemoto, K. *Chem. Lett.* 1985, 1415. (d) Umemoto, K.; Okamura, N. *Bull. Chem. Soc. Jpn.* 1986, 59, 3047.

(6) (a) Fomin, G. V.; Blyumenfel'd, L. A.; Sukhorukov, V. I. *Dokl. Akad. Nauk SSSR* 1964, 157, 1199. (b) Ledwith, A. *Acc. Chem. Res.* 1972, 5, 133.

(7) In methanol, the real reductant has been reported to be CH<sub>3</sub>O<sup>-</sup> formed by the deprotonation of CH<sub>3</sub>OH with OH<sup>-</sup>: Farrington, J. A.; Ledwith, A.; Stam, M. F. *J. Chem. Soc. D* 1969, 259.

(8) In the one-electron reduction of methylviologen in an alkaline aqueous solution, methylviologen itself acts as a reductant to yield methylviologen radical cation and the pyridone derivative: (a) Calderbank, A.; Charlton, D. F.; Farrington, J. A.; James, R. *J. Chem. Soc., Perkin Trans. 1* 1972, 138. (b) Novakovic, V.; Hoffman, M. Z. *J. Am. Chem. Soc.* 1987, 109, 2341.

(9) For electron-transfer pathways in the nucleophilic substitution by OH<sup>-</sup>: (a) Bacaloglu, R.; Bunton, C. A.; Cerichelli, G. *J. Am. Chem. Soc.* 1987, 109, 621. (b) Bacaloglu, R.; Bunton, C. A.; Ortega, F. *J. Am. Chem. Soc.* 1989, 111, 1041. (c) Bacaloglu, R.; Blasko, A.; Bunton, C.; Dorwin, E.; Ortega, F.; Zucco, C. *J. Am. Chem. Soc.* 1991, 113, 238.

(10) The ESR measurement of the deaerated product solution confirmed the formation of Q<sup>•-</sup> (g = 2.0046, a<sub>H</sub> = 2.4 G): (a) Venkataraman, B.; Fraenkel, G. K. *J. Chem. Phys.* 1955, 23, 588. (b) Fukuzumi, S.; Ono, Y.; Keii, T. *Int. J. Chem. Kinet.* 1975, 7, 535.

(11) The yields of semiquinone radical anions were determined from the absorbances at λ<sub>max</sub> by using the reported ε<sub>max</sub> values: Fukuzumi, S.; Nishizawa, N.; Tanaka, T. *J. Org. Chem.* 1984, 49, 3571. The yields of most semiquinone radical anions are high except for those derived from tetrasubstituted quinones (the yields are 7%, 7%, and 6% for chloranil, bromanil, and fluoranil, respectively).

(12) When the product mixture of the reaction of Q (2.0 × 10<sup>-2</sup> M) with Me<sub>4</sub>N<sup>+</sup>OH<sup>-</sup> (2.0 × 10<sup>-2</sup> M) was treated with Ph<sub>3</sub>P (1.0 × 10<sup>-2</sup> M) in deaerated MeCN, no formation of Ph<sub>3</sub>P=O was detected. When H<sub>2</sub>O<sub>2</sub> was added to the same solution, however, Ph<sub>3</sub>P was converted to Ph<sub>3</sub>P=O quantitatively in 40 min. Thus, it is concluded that no H<sub>2</sub>O<sub>2</sub> is formed in the reaction of Q with OH<sup>-</sup> in MeCN. The products in the gas phase were also analyzed by GLC using a molecular sieve 13 X column with He gas carrier. No appreciable formation of O<sub>2</sub> has been detected, either.

(1) Sawyer, D. T.; Roberts, J. L., Jr. *Acc. Chem. Res.* 1988, 21, 469.

(2) (a) Shaik, S. S. *J. Org. Chem.* 1987, 52, 1563. (b) Bunce, E.; Shaik, S. S.; Um, I.-H.; Wolfe, S. *J. Am. Chem. Soc.* 1988, 110, 1275. (c) Shaik, S. S. *Acta Chem. Scand.* 1990, 44, 205.

(3) (a) Roberts, J. L., Jr.; Sugimoto, H.; Barreter, W. C., Jr.; Sawyer, D. T. *J. Am. Chem. Soc.* 1985, 107, 4556. (b) Srivatsa, G. S.; Sawyer, D. T. *Inorg. Chem.* 1985, 24, 1732. (c) Hojo, M.; Sawyer, D. T. *Inorg. Chem.* 1989, 28, 1201.