

Figure 1. Part of the mechanism for the β -tyrosine phenol-lyase-catalyzed elimination reaction of tyrosine. P is a phosphate group and E the enzyme.

reaction mixture. The vials were then centrifuged at 5000 rpm for a few minutes; 5–10 μ L of the quenched reaction mixtures were injected on an HPLC column,⁹ and the separated labeled substrate and product were collected in liquid scintillation bottles. The radioactivity of the fractions was measured immediately, and after total decay of the ^{11}C , the ^{14}C -radioactivity was measured with a liquid scintillation counter. The ratio $k_{11}/k_{14} = \ln(1 - f_{11})/\ln(1 - f_{14})$, where f is the fraction of reaction, was calculated for each reaction point.

The results from three KIE experiments, performed in phosphate buffer pH 6.8 at 18 $^\circ\text{C}$,¹⁰ were 1.068 ± 0.017 ($n = 10$), 1.083 ± 0.013 ($n = 11$), and 1.051 ± 0.012 ($n = 14$), where n is the number of reaction points and the standard deviation is reported. The mean $^{11}\text{C}/^{14}\text{C}$ KIE value is 1.067 ± 0.009 .

Several contributions to the elucidation of the mechanism of tyrosine phenol-lyase action have recently been reported.¹¹ For reviews, see, e.g., Snell and DiMari¹² and Miles.¹³ The chemical mechanism involves the formation of an aldimine between L-tyrosine and pyridoxal 5-phosphate (PLP). The substrate α -proton is abstracted by an enzyme-bound base (see Figure 1; B_1 , $\text{p}K_a = 7.6$ ^{11c}) with the formation of a quinonoid structure (I). Another base (B_2 , $\text{p}K_a = 8.0$ ^{12c}) then abstracts the hydroxyl proton, and the first base (B_1) returns a proton to the aromatic C-4 position with the formation of a cyclohexadienone moiety (II). The activated carbon–carbon bond now breaks with simultaneous electron-push from the PLP, and electron-pull when the hydroxyl proton is returned by the base B_2 . Phenol is released, and after transamination and hydrolysis, pyruvic acid and ammonia are released from the enzyme. In a study of the pH dependence of kinetic parameters and the primary deuterium KIE of tyrosine phenol-lyase from *C. freundii*, it was concluded that the α -proton abstraction was a partially rate-limiting step.^{11c}

For the nonenzymatic malonic acid decarboxylation, a reaction in which carbon–carbon bond breaking (as in the present case) is accompanied by formation of a double bond to the isotopic carbon atom,¹⁴ k_{12}/k_{14} for acid labeled in the 2-position was determined to be 1.076 by Ropp and Raaen.¹⁵ Using the relation

$\ln(k_{11}/k_{14})/\ln(k_{12}/k_{14}) \approx 1.6$ ^{5b} between the different carbon isotope effects, the $^{12}\text{C}/^{14}\text{C}$ KIE corresponding to our value may be estimated to be 1.04. Our results, in combination with earlier conclusions,^{11c} therefore suggest that the C–C bond breaking is at least partially rate limiting.

Acknowledgment. We thank Dr. Petter Malmberg for his help with the radionuclide production and Prof. Y. Watanabe for supplying the tyrosine phenol-lyase. This project is supported financially by the Swedish Natural Science Research Council.

One-Pot Synthesis of Aromatic Methyl Esters by Electrochemical Oxidation of Aldehydes Mediated by Biscoenzyme Catalysis

Suk-wah Tam, Leslie Jimenez, and François Diederich*

Department of Chemistry and Biochemistry
University of California at Los Angeles
Los Angeles, California 90024-1569

Received October 24, 1991

The direct oxidation of aldehydes to esters under mild conditions is a useful transformation in organic synthesis.¹ Here we describe an efficient, one-pot synthesis of aromatic methyl esters by electrochemical oxidation of aldehydes, mediated by two coenzyme catalysts: the thiazolium ions **1a/b** and flavin MeFl. The use of the macrocyclic catalyst **1b** in electroorganic synthesis elegantly combines the principles of electrocatalysis² with molecular recognition (Chart I).³

Thiazolium ions are known to catalyze the oxidation of aldehydes to esters.⁴ The thiazolium ylide **2**⁵ reacts to give the "active aldehyde" **3** (Scheme I).⁶ This reactive intermediate can condense with another aldehyde to give an acyloin or can be oxidized to give the 2-acylthiazolium ion **4**. This ion reacts readily in alcohol⁷ to give an ester (Scheme I). Stoichiometric amounts of oxidizing agents like nitrobenzene,⁸ potassium ferricyanide,⁹ and flavins¹⁰ cause solubility problems and complicate product isolation. Also, the thiazolium catalyst is destroyed oxidatively in basic solution by ferricyanide,^{9,11} iodine,¹² and air.^{11,13} Attempts to regenerate catalytic amounts of MeFl with air resulted in the oxidation of **1a/b** by air (Table II, entry e).¹³

Our investigation by cyclic voltammetry shows that the anodic peak potential of **1a/b** in a 0.05 M solution of NEt_3Br in MeOH is ~ 0.2 V (vs Ag/AgCl at 0.02V s^{-1}), while that of MeFl is ca. -0.47 V ($E_{1/2}$ ca. -0.52 V at 0.02V s^{-1}). Under argon atmosphere,

(1) (a) Larock, R. C. *Comprehensive Organic Transformations*; VCH: New York, 1989. (b) Williams, D. R.; Klingler, F. D.; Allen, E. E.; Lichtenthaler, F. W. *Tetrahedron Lett.* **1988**, 29, 5087–5090. (c) McDonald, C.; Holcomb, H.; Kennedy, K.; Kirkpatrick, E.; Leathers, T.; Vanemon, P. *J. Org. Chem.* **1989**, 54, 1213–1215.

(2) Steckhan, E. *Top. Curr. Chem.* **1987**, 142, 1–69.

(3) For electrochemical conversions in the presence of cyclodextrins, see: (a) Smith, C. I.; Utley, J. H. P. *J. Chem. Soc., Chem. Commun.* **1981**, 792–793. (b) Matsue, T.; Evans, D. H.; Osa, T.; Kobayashi, N. *J. Am. Chem. Soc.* **1985**, 107, 3411–3417.

(4) (a) Krampitz, L. O. *Annu. Rev. Biochem.* **1969**, 38, 213–240. (b) Kluger, R. *Chem. Rev.* **1987**, 87, 863–876.

(5) Breslow, R. *J. Am. Chem. Soc.* **1957**, 79, 1762–1763.

(6) (a) Breslow, R. *J. Am. Chem. Soc.* **1958**, 80, 3719–3726. (b) Zeng, X.; Chung, A.; Jordan, F. *J. Am. Chem. Soc.* **1991**, 113, 5842–5849.

(7) (a) Breslow, R.; McNelis, E. *J. Am. Chem. Soc.* **1960**, 82, 2394–2395. (b) Nash, C. P.; Olsen, C. W.; White, F. G.; Ingraham, L. L. *J. Am. Chem. Soc.* **1961**, 83, 4106–4107.

(8) (a) Castells, J.; Litjens, H.; Moreno-Manas, M. *Tetrahedron Lett.* **1977**, 18, 205–206. (b) Inoue, H.; Tamura, S. *J. Chem. Soc., Chem. Commun.* **1985**, 141–142.

(9) (a) Hilvert, D.; Breslow, R. *Bioorg. Chem.* **1984**, 12, 206–220. (b) Jimenez, L.; Diederich, F. *Tetrahedron Lett.* **1989**, 30, 2759–2762.

(10) (a) Shinkai, S.; Yamashita, T.; Kusano, Y.; Manabe, O. *J. Org. Chem.* **1980**, 45, 4947–4952. (b) Shinkai, S.; Yamashita, T.; Kusano, Y.; Manabe, O. *Tetrahedron Lett.* **1980**, 21, 2543–2546. (c) Yano, Y.; Tsukagoshi, Y. *J. Chem. Res.* **1984**, 406–407.

(11) Maier, G. D.; Metzler, D. E. *J. Am. Chem. Soc.* **1957**, 79, 4386–4391.

(12) Clarke, H. T.; Gurin, S. *J. Am. Chem. Soc.* **1935**, 57, 1876–1881.

(13) Jimenez, L. Ph.D. Thesis, University of California, Los Angeles, CA, 1989.

(9) A RP C-18 column was used, and the mobile phase was 0.01 M acetic acid, pH 3.5; methanol (75:25, v/v) isocratic flow 2.0 mL min^{-1} .

(10) The optimum is pH 8, but because of a slow sampling technique, the rate of the reaction was decreased by lowering the pH and temperature.

(11) (a) Faleev, N. G.; Lyubarev, A. E.; Martinkova, N. S.; Belikov, V. M. *Enzyme Microb. Technol.* **1983**, 5, 219. (b) Palcic, M. M.; Shen, S.-J.; Schleicher, E.; Kumagai, S.; Sawada, S.; Yamada, H.; Floss, H. G. *Z. Naturforsch.* **1987**, 42c, 307. (c) Kiick, D. M.; Phillips, R. S. *Biochemistry* **1988**, 27, 7333. (d) Faleev, N. G.; Ruvinov, S. B.; Demidkina, T. V.; Myagkikh, I. V.; Gololobov, M. Y.; Bakmutov, V. I.; Belikov, V. M. *Eur. J. Biochem.* **1988**, 177, 395.


(12) Snell, E. E.; Di Mari, S. J. *The Enzymes*, 3rd ed.; Boyer, P. D., Ed.; Academic Press: New York and London, 1970; Vol. 2, p 335.

(13) Miles, E. W. *Coenzymes Cofactors* **1986**, 1, 253.

(14) Westheimer, F. H.; Jones, W. A. *J. Am. Chem. Soc.* **1941**, 63, 3283.

(15) Ropp, G. A.; Raaen, V. F. *J. Am. Chem. Soc.* **1952**, 74, 4992.


Scheme I

Table I. Electrochemical Oxidation of Aldehydes^a

aldehyde	ratio of aldehyde to hemiacetal ^c	thiazolium catalyst	time (h) ^d	yield (%) ^e	current efficiency (%) ^f	turnover number ^b
valeraldehyde	1:16	1a	26	17	77	1.7
		1b	26	20	83	2.0
cyclohexanecarboxaldehyde	1:6.4	1a	24	7 ^g	23	0.7
2-naphthaldehyde	1:0.074	1a	21	55	72	5.5
		1b	10	74 (76)	88	7.4
benzaldehyde	1:0.10	1a	21	54	60	5.4
4-cyanobenzaldehyde	1:2.3	1a	18	72 (69)	70	7.2
		1b	6	95 (88)	90	9.5
4-chlorobenzaldehyde	1:0.22	1a	18	78 (81)	84	7.8
methyl 4-formylbenzoate	1:0.85	1a	18	85 (83)	80	8.5

^aThe oxidation used an undivided cell equipped with a glassy carbon rod cathode (6 mm o.d.) and a platinum plate anode (12 mm × 50 mm) at -0.3 V vs a Ag/AgCl/3 M NaCl reference electrode at 308 K. The cell contained 11-mL solutions of electrolytes in MeOH with mole ratios of aldehydes (1.65 mmol, 150 mM):**1**:MeFl:NEt₃:NEt₄Br = 30:1:3:30:9. ^bCalculated as the number of moles of product per mole of MeFl, the catalyst with the highest concentration. ^cDetermined by 200-MHz NMR of 150 mM solutions of aldehydes in 150 mM NEt₃ in CD₃OD. ^dThe reactions were stopped when the current dropped to <10%. ^eDetermined by gas chromatography. Numbers in brackets are isolated yields. ^fCalculated by (2 × moles of product × 100)/Faradays passed. ^gApproximately 80% of aldehyde remained unreacted.

Chart I



MeFl can be regenerated electrochemically at -0.3 V without causing oxidative destruction of **1a/b**. Therefore, we developed an unprecedented electrochemical regeneration cycle of the two coenzymes (Scheme I).

Using inexpensive **1a** and **MeFl** as catalysts, aromatic aldehydes give a 55–85% yield of methyl esters with high current efficiency (Table I). The much lower yields of aliphatic esters (<20%) are presumably due to the competing formation of hemiacetals, which significantly lowers the equilibrium concentration of aldehydes and hence the rate of formation of the “active aldehyde” intermediate, the slow step under the chosen reaction conditions (Table

Table II. Control Experiments Performed with 4-Cyanobenzaldehyde (1.65 mmol) and Thiazolium Ion **1a** (0.055 mmol)

entry	MeFl (mmol)	atmosphere	working electrode potential (V vs Ag/AgCl)	time (h)	yield (%) ^a
a	0.165	argon	-0.3	18	72
b		argon	-0.3	20	19
c	0.165	argon		20	12 ^b
d		argon		20	4
e	0.165	air		20	35
f		air		20	15 ^c

^aDetermined by gas chromatography. ^bTheoretical yield from **MeFl** is 10%. ^c0% yield in the absence of **1a**.

I). In the case of 4-cyanobenzaldehyde, the electron-withdrawing effect of the cyano group increases the reaction rate significantly and compensates for the decrease in equilibrium concentration of the aldehyde due to hemiacetal formation.

The supramolecular catalyst **1b** significantly enhances the rate^{9b,13} and yield of aromatic ester formation (Table I). This catalyst forms tight inclusion complexes with benzene and naphthalene substrates in methanol. The increased rates are probably a result of (i) entropically favorable orientation and proximity effects and (ii) microenvironmental effects¹⁴ in the apolar cyclophane cavity. Rates of thiazolium-catalyzed reactions increase with reduced solvent polarity since the relevant reaction

(14) (a) Diederich, F.; Lutter, H.-D. *J. Am. Chem. Soc.* **1989**, *111*, 8438–8446. (b) Jencks, W. P. *Adv. Enzymol. Relat. Areas Mol. Biol.* **1975**, *43*, 219–410. (c) Jordan, F.; Kuo, D. J.; Monse, E. U. *J. Org. Chem.* **1978**, *43*, 2828–2830.

transition states are less polar than the ground states.¹⁵ The acidity of the macrocyclic thiazolium ring is enhanced by the apolar environment provided by the cavity.^{14a} In deuterated acetate buffer (pD 4.7), the rate of H/D exchange at C-2 of the thiazolium ring in **1b** is about 2.6 times faster than that measured for **1a**.

The results of control experiments (Table II) show that the high yield obtained in entry a is due to the efficient regeneration of MeFl at the anode. Direct oxidation at the anode (entry b) is not an efficient enough process to trap all of the "active aldehyde" intermediates.^{16,17}

The full scope of the supramolecular electrochemical process mediated by **1b** and the very useful oxidation of aldehydes to carboxamides are now under investigation.

Acknowledgment. This work was supported by the Office of Naval Research and BASF AG, Ludwigshafen. We thank Dr. Dieter Degner, BASF, for continuing encouragement and support.

(15) Crosby, J.; Stone, R.; Lienhard, G. E. *J. Am. Chem. Soc.* **1970**, *92*, 2891–2900.

(16) The redox potential of the "active aldehyde" intermediates cannot be determined. For redox potentials of related enamines, see: (a) Bordwell, F. G.; Satish, A. V.; Jordan, F.; Rios, C. B.; Chung, A. C. *J. Am. Chem. Soc.* **1990**, *112*, 792–797. (b) Barletta, G.; Chung, A. C.; Rios, C. B.; Jordan, F.; Schlegel, J. M. *J. Am. Chem. Soc.* **1990**, *112*, 8144–8149.

(17) For electrochemical oxidation of aldehydes catalyzed by cyanide (+1.7 V vs SCE), see: Chiba, T.; Okimoto, M.; Nagai, H.; Takata, Y. *Bull. Chem. Soc. Jpn.* **1982**, *55*, 335–336. For electrochemical oxidation of aldehyde catalyzed by iodide (+0.6–0.8 V vs SCE), see: (a) Okimoto, M.; Chiba, T. *J. Org. Chem.* **1988**, *53*, 218–219. (b) Shono, T.; Matsumura, Y.; Hayashi, J.; Inoue, K.; Iwasaki, F.; Itoh, T. *J. Org. Chem.* **1985**, *50*, 4967–4969.

Facilitation of the $\Delta^2 \rightarrow \Delta^1$ Pyrroline Tautomerization of Carbapenem Antibiotics by the Highly Conserved Arginine-244 of Class A β -Lactamases during the Course of Turnover

Glenn Zafaralla and Shahriar Mobashery*

Department of Chemistry, Wayne State University
Detroit, Michigan 48202

Received October 31, 1991

The hydrolytic action of β -lactamases is the primary mechanism of bacterial resistance to β -lactam antibiotics.¹ Within the past several years a variety of new β -lactam drugs have been developed that show resistance to the action of these enzymes.² Carbapenems constitute a group of β -lactamase-resistant molecules, and they possess potent activity against a wide spectrum of bacteria.³ Studies on the mechanism of action of class A β -lactamases with carbapenems by Knowles and colleagues have indicated a biphasic profile for hydrolysis of carbapenems, with an initial fast phase for substrate turnover leading to a slower one within minutes.^{4,5} It was demonstrated that subsequent to active

(1) Bush, K. *Rev. Infect. Dis.* **1988**, *10*, 681. Sanders, C. C.; Sanders, W. E. *J. Infect. Dis.* **1985**, *151*, 399.

(2) For reviews see: Pratt, R. F. In *Design of Enzyme Inhibitors as Drugs*; Sandler, M., Smith, H. J., Eds.; Oxford Press: Oxford, UK, 1989; pp 178–205. Fisher, J. In *Antimicrobial Drug Resistance*; Bryan, J. T., Ed.; Academic Press: New York, 1984; pp 33–79.

(3) Mendell, L. *Can. Med. Assoc. J.* **1988**, *139*, 505. Jones, R. N. *Am. J. Med.* **1985**, *78*, Suppl. 6A, 22. Aukenthaler, R.; Wilson, W.; Wright, A.; Washington, J.; Durack, D.; Geraci, J. *Antimicrob. Agents Chemother.* **1982**, *22*, 448. Cullman, W.; Opferkuch, W.; Slieglitz, M.; Werkmeister, U. *Antimicrob. Agents Chemother.* **1982**, *22*, 302. Kropp, H.; Sundelof, J.; Hadju, R.; Kahan, F. *Antimicrob. Agents Chemother.* **1982**, *22*, 62. Kropp, H.; Sundelof, J.; Kahan, F.; Birnbaum, J. *Antimicrob. Agents Chemother.* **1980**, *17*, 993.

(4) Easton, C. J.; Knowles, J. R. *Biochemistry* **1982**, *21*, 2857.

(5) Charnas, R. L.; Knowles, J. R. *Biochemistry* **1981**, *20*, 2732.




Figure 1. Hydrolysis of imipenem (20 μ M) by the wild-type TEM-1 (2 μ M) and the Arg-244-Ser mutant (2 μ M) β -lactamases, in 0.1 M potassium phosphate buffer, pH 7.0, at room temperature (—). Extrapolation of the linear second phase of hydrolysis by the wild-type enzyme to time zero (---) and inhibition of activity of the wild-type (O) and the Ser-244 mutant (●) TEM-1 β -lactamases (as monitored for the turnover of benzylpenicillin) are indicated.

Scheme I





Chart I



site acylation of a β -lactamase (at Ser-70) by these molecules, the Δ^2 -pyrroline analogue **2** may either undergo deacylation or tautomerize to the corresponding Δ^1 -derivative (**3**). The ester bond of **3** is kinetically more resistant to hydrolysis because of a less favorable substrate positioning in the active site (Scheme I). We present evidence here that the highly conserved arginine-244⁶ is the essential source of proton for the $\Delta^2 \rightarrow \Delta^1$ tautomerization of carbapenem antibiotics, as depicted in Scheme I.

High-resolution crystal structures for two class A β -lactamases from *Staphylococcus aureus* PC1⁷ and *Bacillus licheniformis* 749/C^{8,9} have been reported recently. The information from crystal structure, in conjunction with kinetic findings from our laboratory, indicated that the substrate carboxylate forms hy-

(6) Arginine-244 is conserved in the majority of class A β -lactamases: Ambler, R. P.; Coulson, A. F. W.; Frère, J. M.; Ghuyssen, J. M.; Joris, B.; Forsman, M.; Levesque, R. C.; Tiraby, G.; Waley, S. G. *Biochem. J.* **1991**, *276*, 269. In a few known class A β -lactamases that do not possess arginine at position 244, the Arg-220 side chain has been shown to occupy the same space as that of Arg-244 in the majority of class A enzymes: Jacob-Dubuisson, F.; Lamote-Brasseur, J.; Dideberg, O.; Joris, B.; Frère, J. M. *Protein Eng.* **1991**, *4*, 811.

(7) Herzberg, O.; Moul, J. *Science* **1987**, *236*, 694.

(8) Moews, P. C.; Knox, J. R.; Dideberg, O.; Charlier, P.; Frère, J. M. *Proteins* **1990**, *7*, 156.

(9) Knox, J. R.; Moews, P. C. *J. Mol. Biol.* **1991**, *220*, 435.