

of MNNG may be accessible to the thiol at the active site. Given this, the chemoprotective activity of the glutathione *S*-transferase-mediated reaction of glutathione may be due to a denitrosation reaction that is predominant because the thiolate of GSH, when it reacts at carbon, generates an intermediate T that mainly reverts to starting materials due to the low conjugate

acid pK_a of the thiolate at the active site.

Supplementary Material Available: Tables S1-S4 of data from reactions of thiols with *N*-methyl-*N*'-nitro-*N*-nitroguanidine, 40 °C, ionic strength 1 M (KCl) (14 pages). Ordering information is given on any current masthead page.

Coordination Isomerism in Pentamethylcyclopentadienyl-Substituted Iminophosphanes: From Classical Structures to a π -Complexed Iminophosphenium Ion

D. Gudat,[†] H. M. Schiffner,[†] M. Nieger,[†] D. Stalke,[‡] A. J. Blake,[‡] H. Grondey,[§] and E. Niecke^{*,†}

Contribution from the Anorganisch Chemisches Institut der Universität, Universität Bonn, Gerhard Domagk Strasse 1, D-5300 Bonn 1, Germany, Anorganisch Chemisches Institut der Universität, Universität Göttingen, Tammannstrasse 4, D-3400 Göttingen, Germany, and Max Planck Institut für Kohleforschung, Kaiser Wilhelm Platz 1, D-4330 Mülheim/Ruhr, Germany. Received October 16, 1991

Abstract: Novel iminophosphanes of the type $\text{Me}_5\text{C}_5\text{P}=\text{NR}$ ($\text{R} = \text{C}_6\text{H}_2\text{tBu}_3$ (**5a**), $\text{Si-}i\text{-Pr}_3$ (**5b**), SiMe_3 (**5c**)) are prepared via either thermolytic or base-promoted elimination reactions and characterized by analytical and spectroscopic methods. The X-ray crystal structures of **5a,b** establish the presence of coordination isomerism of the cyclopentadienyl ring, which at the same time strongly alters the characteristics of the P-N multiple bond. For **5a**, η^1 -attachment of the Me_5C_5 moiety to the phosphorus and a localized phosphorus-carbon σ -bond are found, and the P-N double bond compares to that of alkylated iminophosphanes ($r_{\text{PN}} = 155.1$ (8) pm). **5b** exhibits η^2 -coordination of the cyclopentadienyl ring leading to increased P-N triple-bond character, as evidenced by shortening of the P-N distance (153.3 (3) pm) and the remarkable opening of the nitrogen valence angle (153.3 (2)°). The structure is discussed as an intramolecular π -complex between a formal cyclopentadienyl anion and an iminophosphenium cation. The ^1H and ^{13}C NMR spectra show that rapid elementotropic rearrangements around the five-membered ring take place both in solution and in the solid state. The unique temperature dependence of $\delta^{31}\text{P}$ in solution together with the considerable differences in $\delta^{31}\text{P}$ between the solution and solid states are interpreted in a model assuming a "haptotropic" mechanism for the fluxionality involving dynamic η^1/η^2 -coordination isomerizations in solution. As expected, nucleophilic displacement of the Me_5C_5 moiety takes place in the reaction of **5b,c** with $\text{LiC}_6\text{H}_2\text{tBu}_3$, yielding the iminophosphanes $\text{tBu}_3\text{H}_2\text{C}_6\text{P}=\text{NR}$ ($\text{R} = \text{Si-}i\text{-Pr}_3$ (**8b**), SiMe_3 (**8c**)) as products.

Introduction

Pentamethylcyclopentadienyl-substituted (Me_5C_5) phosphorus- π multiple-bond systems are of both high synthetic and theoretical interest due to their unique stereoelectronic properties. While the Me_5C_5 group provides sufficient steric bulk for the stabilization of the low coordination number,¹⁻³ its liability toward nucleophilic substitution^{3,4} at the same time permits a synthetically valuable functionalization of the double bond. In addition, the capacity for multihapto π -bonding⁵ leads to a versatility in structure and bonding. Thus, the cyclopentadienyls in the neutral systems $\text{Me}_5\text{C}_5\text{P}=\text{ER}_n$ ($\text{ER}_n = \text{C}(\text{SiMe}_3)_2$,¹ PC_5Me_5 ³) are η^1 -(σ)-bonded in the solid state, while "nonclassical" $\eta^2(\pi)$ -coordination prevails in the phosphenium cation $[\text{Me}_5\text{C}_5\text{PN}(\text{H})\text{tBu}]^+$.⁶ In solution, all known systems are fluxional due to fast elementotropic rearrangements.^{6,7}

Although an Me_5C_5 -substituted iminophosphane has been prepared,² no structure determinations of such compounds, which should constitute a "missing link" between $\eta^1\text{-Me}_5\text{C}_5\text{P}=\text{PR}$ and $[\eta^2\text{-Me}_5\text{C}_5\text{PNHR}]^+$, have been performed yet. Here, we report the preparation and first structural characterization of novel Me_5C_5 -substituted iminophosphanes. It is found that a substitution-induced coordination isomerism occurs between $\eta^1(\sigma)$ - and

$\eta^2(\pi)$ -bonding of the Me_5C_5 ligand, which has most interesting implications on the structure and bonding of the phosphorus-nitrogen multiple bond. The investigation of the dynamic aspects of the molecular structure was attempted by NMR spectroscopic studies of **5a** both in the solid state and in solution, and evidence is presented that the dynamic process may follow different mechanisms in both cases.

Experimental Section

All manipulations were carried out under a dry argon atmosphere using standard Schlenk techniques. Solvents were dried by refluxing over appropriate reagents and distilled before use. $\text{Me}_5\text{C}_5\text{Li}$,⁸ (2,4,6-tri-*tert*-butylanilino)dichlorophosphane (**2a**),⁹ lithium bis(trimethylsilyl)-amide (**4**),¹⁰ [bis(trimethylsilyl)amino]dichlorophosphane (**6**),¹¹ and (2,4,6-tri-*tert*-butylphenyl)lithium¹² were prepared according to literature

- (1) Gudat, D.; Niecke, E.; Krebs, B.; Dartmann, M. *Chimia* **1985**, *39*, 277.
- (2) Gudat, D.; Niecke, E.; Krebs, B.; Dartmann, M. *Organometallics* **1986**, *5*, 2376.
- (3) Jutzi, P.; Meyer, U.; Krebs, B.; Dartmann, M. *Angew. Chem., Int. Ed. Engl.* **1986**, *25*, 919.
- (4) Gudat, D.; Niecke, E. *J. Chem. Soc., Chem. Commun.* **1987**, 10.
- (5) Jutzi, P. *Adv. Organomet. Chem.* **1986**, *26*, 217.
- (6) Gudat, D.; Nieger, M.; Niecke, E. *J. Chem. Soc., Dalton Trans.* **1989**, 693.
- (7) Jutzi, P. *Chem. Rev.* **1986**, *86*, 983.
- (8) Jutzi, P.; Saleske, H. *Chem. Ber.* **1984**, *117*, 222.
- (9) Niecke, E.; Lysek, M.; Symalla, E. *Chimia* **1986**, *40*, 202.
- (10) Wannagat, U.; Niederprüm, H. *Chem. Ber.* **1961**, *94*, 1540.
- (11) Scherer, O. J.; Kuhn, N. *J. Organomet. Chem.* **1974**, *82*, C3.

[†] Universität Bonn.

[‡] Universität Göttingen.

[§] Max Planck Institut für Kohleforschung.

methods. (Triisopropylsilyl)amine (**1b**) was prepared as described in ref 13 and was purified by distillation (bp 43–45 °C/3 Torr).

Solution $^{31}\text{P}\{^1\text{H}\}$ NMR spectra (32.2 MHz) were recorded on a Varian FT80A, and ^1H (at 79.5 and 200 MHz) and $^{13}\text{C}\{^1\text{H}\}$ NMR spectra (at 20.0 and 50.3 MHz) were recorded on both Varian FT80A and Bruker AC 200 instruments. For measurements at ambient temperature, C_6D_6 (previously dried over 4-Å molecular sieves) was used as solvent unless otherwise noted. Chemical shifts are referenced to external TMS (^1H , ^{13}C) or 85% H_3PO_4 (^{31}P); multiplicities refer to coupling with ^{31}P unless stated otherwise. Variable-temperature NMR spectra of **5a,b** were obtained in toluene- d_8 (^1H , ^{13}C) or hexane solution with approximately 0.15 mL of toluene- d_8 added in order to obtain a sufficient deuterium lock signal (^{31}P), respectively. Temperature measurements were performed by means of a ^{31}P chemical shift thermometer¹⁴ which was placed in a thin-walled 5-mm NMR tube and concentrically adjusted inside the 10-mm sample tube. Solid-state CP/MAS $^{13}\text{C}\{^1\text{H}\}$ and $^{31}\text{P}\{^1\text{H}\}$ spectra were recorded on a Bruker MSL 300 spectrometer under high-power proton decoupling. Spinning rates between 2500 and 3300 Hz, relaxation delays of 4 s (3 s in ^{31}P spectra), and contact times of 1 ms were applied. The temperature was varied between 115 and 295 K with use of a temperature controller and nitrogen as the driving and bearing gas. Chemical shifts were referenced by using the methylene carbon resonance of adamantane ($\delta(^{13}\text{C})$ 38.4) or H_3PO_4 as external references. Loading of the rotors was carried out in an inert argon atmosphere using a specially designed apparatus as described elsewhere.¹⁵

Electron ionization (70 eV) mass spectra were run on a VG 12-250 spectrometer. Elemental analyses were determined on a Heraeus CHN-O-Rapid instrument; all melting points were determined in sealed glass capillaries.

Preparation of $i\text{Pr}_3\text{SiNHPCl}_2$ (2b**).** To a stirred solution of the amine **1b** (0.115 mol, 30.0 g) in hexanes (50 mL) was added dropwise 1 equiv of a 1.6 M solution of $n\text{-BuLi}$ in hexane (0.115 mol, 72 mL). The resulting mixture was stirred for 15 min and then added dropwise to a stirred solution of PCl_3 (0.120 mol, 16.5 g) in hexane (100 mL) at -78 °C. After the reaction was warmed to ambient temperature, the precipitate was removed by filtration, and the solution was evaporated in vacuo. The residue was dissolved in pentane (50 mL) and cooled to -80 °C. The product precipitated as white crystals, which were collected by filtration at low temperature and dried in vacuo: yield 87% (27.4 g); mp 0 °C; $^{31}\text{P}\{^1\text{H}\}$ NMR δ 168.3; ^1H NMR δ 1.94 (m, 21 H, $\text{SiCH}(\text{CH}_3)_2$).

Preparation of $\text{C}_6\text{H}_5\text{tBu}_3\text{NHPClC}_3\text{Me}_5$ (3a**).** A solution of compound **2a** (55.0 mmol, 19.9 g) in ether (100 mL) was added dropwise to a stirred, ice-cooled suspension of $\text{Me}_3\text{C}_3\text{Li}$ (60.0 mmol, 8.52 g) in 120 mL of pentane. After the mixture was stirred for an additional 3 h at ambient temperature, the precipitated salts were removed by filtration, and the solution was evaporated in vacuo. The residue was dissolved in pentane and cooled to -80 °C. The product separated as white crystals, which were collected by filtration and dried in vacuo: yield 72% (18.3 g); mp 165 °C; $^{31}\text{P}\{^1\text{H}\}$ NMR δ 153.0; ^1H NMR δ 7.51 (2 H, C_6H_5), 1.92 (broad, 15 H, $\text{C}_5(\text{CH}_3)_5$), 1.53 (18 H, $o\text{-CCH}_3$), 1.31 (9 H, $p\text{-CCH}_3$).

Preparation of $i\text{Pr}_3\text{SiNHPClC}_3\text{Me}_5$ (3b**).** A solution of **2b** (55.0 mmol, 15.0 g) in pentane (100 mL) was reacted with LiC_3Me_5 in the same manner as that described for **3a**. Crystallization at -35 °C yielded 81% (16.8 g) of **3b**: mp 55 °C; $^{31}\text{P}\{^1\text{H}\}$ NMR δ 148.6; ^1H NMR δ 1.73 (3 H), 1.56 (3 H), 1.50 (3 H), 1.33 (d, 4.5 Hz, 3 H), 1.16 (d, 15 Hz, 3 H, $\text{C}_5(\text{CH}_3)_5$), 0.73 (m, 21 H, $\text{SiCH}(\text{CH}_3)_2$).

Preparation of $\text{C}_6\text{H}_5\text{tBu}_3\text{N}=\text{PC}_3\text{Me}_5$ (5a**).** A precooled (-50 °C) solution of 26.2 mmol of lithium bis(trimethylsilyl)amide (**4**) (4.40 g) in THF (30 mL) was added slowly to a stirred solution of compound **3a** (26.2 mmol, 12.2 g) in THF (40 mL) at -80 °C. When the addition was complete, the resulting yellow solution was gradually allowed to warm. Precipitation of LiCl , which was accompanied by a deepening of the color, started above -60 °C and was complete at room temperature. After evaporation of the volatiles in vacuo, the residue was extracted with pentane (100 mL) and filtered. The filtrate was concentrated to 50 mL and cooled to -30 °C. **5a** precipitated as yellow crystals, which were collected by filtration and dried in vacuo: yield 63% (6.3 g); mp 115 °C; MS m/e 425 (7), M^+ , 290 (100, $\text{M}^+ - \text{C}_3\text{Me}_5$), and further fragmentation peaks; $^{31}\text{P}\{^1\text{H}\}$ NMR (26 °C, hexane/ C_6D_6) δ 194.9; $^{13}\text{C}\{^1\text{H}\}$ NMR (26 °C) δ 143.3 (d, 8 Hz), 141.5, 136.6 (d, 7 Hz), 121.7 (C_{arom}), 125.7 (d, 4 Hz, C_5), 36.3, 31.9 (d, 2 Hz, C_{quart}), 34.7, 32.2 (CCH_3), 10.8 ($\text{C}_5(\text{CH}_3)_5$); ^1H NMR (26 °C) δ 7.60 (2 H, C_6H_5), 2.00 (15 H, C_5 -

(CH_3)₅), 1.68 (18 H, $o\text{-CCH}_3$), 1.46 (9 H, $p\text{-CCH}_3$). Anal. Calcd for $\text{C}_{28}\text{H}_{44}\text{NP}$: C, 79.01; H, 10.42; N, 3.29. Found: C, 76.83; H, 10.47; N, 3.51.

Preparation of $i\text{Pr}_3\text{SiN}=\text{PC}_3\text{Me}_5$ (5b**).** Compound **5b** was prepared in the same manner as that described for **5a** from **3b** (26.2 mmol, 9.80 g) and **4** (26.2 mmol, 4.40 g). The crude product was purified by vacuum distillation: bp 85 °C (0.01 Torr); yield 70% (6.10 g); MS m/e 337 (23, M^+), 294 (97, $\text{M}^+ - i\text{Pr}$), 202 (23, $\text{M}^+ - \text{C}_3\text{Me}_5$), 160 (100, $\text{M}^+ - i\text{Pr} - \text{C}_3\text{Me}_5$), and further fragmentation peaks; $^{31}\text{P}\{^1\text{H}\}$ NMR (26 °C, hexane/ C_6D_6) δ 159 (broad); $^{13}\text{C}\{^1\text{H}\}$ NMR (26 °C) δ 124.5 (d, 5 Hz, C_5), 18.7 (SiCCH_3), 13.3 (d, 1 Hz, SiC), 10.2 (d, 2 Hz, $\text{C}_5(\text{CH}_3)_5$); ^1H NMR (26 °C) δ 1.80 (d, 0.6 Hz, 15 H, $\text{C}_5(\text{CH}_3)_5$), 1.14 (18 H) and 1.01 (3 H), AB_2X spin system, $J_{\text{AB}} = 7.5$ Hz, $J_{\text{AX}} = 0.5$ Hz, $J_{\text{BX}} = 0$, $\text{SiCH}^*(\text{CH}_3)_2$ ($\text{X} = ^{31}\text{P}$). Anal. Calcd for $\text{C}_{15}\text{H}_{36}\text{NPSi}$: C, 67.61; H, 10.75; N, 4.15. Found: C, 65.14; H, 10.89; N, 4.59.

Preparation of $\text{Me}_3\text{SiN}=\text{PC}_3\text{Me}_5$ (5c**).** Reaction of **6** (100 mmol, 26.2 g) with LiC_3Me_5 (110 mmol) in pentane (200 mL) in the same manner as that described for **3a** afforded the chlorophosphane **7**. The crude product was transferred into a distillation apparatus that was equipped with a 10-cm Vigreux column and a distillation head and was refluxed at 120 °C in vacuo. Me_2SiCl , which formed during the elimination, was collected in a cold trap (-196 °C). After the reaction was complete (approximately 1 h), the iminophosphane that formed was distilled off at the same temperature. The product was purified by fractionated redistillation to yield pure **5c** as a deep yellow liquid of bp 52–55 °C (0.01 Torr): yield 75% (18.9 g); MS m/e 253 (34, M^+), 135 (79, C_3Me_5^+), 118 (100, $\text{M}^+ - \text{C}_3\text{Me}_5$), 73 (82, Me_2Si^+), and further fragmentation peaks; $^{31}\text{P}\{^1\text{H}\}$ NMR (26 °C, hexane/ C_6D_6) δ 138 (broad); $^{13}\text{C}\{^1\text{H}\}$ NMR (26 °C) δ 124.1 (d, 4 Hz, C_5), 10.2 (d, 3 Hz, $\text{C}_5(\text{CH}_3)_5$), 3.9 (d, 2 Hz, SiC); ^1H NMR (26 °C) δ 1.80 (15 H, $\text{C}_5(\text{CH}_3)_5$), 0.31 (9 H, SiCH_3).

Preparation of $i\text{Pr}_3\text{SiN}=\text{PC}_6\text{H}_5\text{tBu}_3$ (10b**).** A solution of compound **5b** (4.20 mmol, 1.40 g) in THF (3 mL) was rapidly added via syringe to a stirred solution (2,4,6-*tert*-butylphenyl)lithium (4.20 mmol, 1.06 g) in THF (10 mL) at -80 °C. The reaction mixture was allowed to warm to ambient temperature over 1 h. After removal of the solvent in vacuo, the residue was extracted with pentane (25 mL) and filtered. The filtrate was evaporated and the residue distilled in vacuo to afford **10b** as a deep yellow liquid of bp 135–138 °C (0.01 Torr), which solidified below room temperature: yield 53% (1.0 g); mp 10–15 °C; $^{31}\text{P}\{^1\text{H}\}$ NMR (26 °C, hexane/ C_6D_6) δ 497.5; $^{13}\text{C}\{^1\text{H}\}$ NMR δ 154.6 (d, 102 Hz), 153.3 (d, 6 Hz), 150.7, 122.2 (C_{arom}), 38.5, 35.1 (CCH_3), 34.8 (d, 7 Hz), 31.4 (C_{quart}), 18.8 (SiCCH_3), 13.3 (SiC); ^1H NMR δ 7.50 (2 H, C_6H_5), 1.60 (9 H, $p\text{-CCH}_3$), 1.30 (18 H, $o\text{-CCH}_3$), 1.2 (m, 21 H, $\text{SiCH}(\text{CH}_3)_2$). Anal. Calcd for $\text{C}_{28}\text{H}_{44}\text{NP}$: C, 72.43; H, 11.25; N, 3.13. Found: C, 71.17; H, 11.48; N, 1.63.

X-ray Crystal Structure Determination of **5a,b.** The diffraction data were collected with Nicolet R3m (**5a**) and STOE AED (**5b**) diffractometers, respectively. In the case of **5b** a semiempirical absorption correction on the basis of Ψ -scans was applied ($T_{\text{min/max}} = 0.50/0.623$). The structures were solved by direct methods. All non-hydrogen atoms were refined anisotropically. H-atoms were located by difference electron density determinations and were either refined by using a riding model with isotropic thermal parameters fixed at 1.2 times that of the bonded carbon (**5a**) or refined free with common isotropic thermal parameters (**5b**). The C–H distance was fixed at 0.96 Å in both cases. Calculations were performed with SHELXTL-Plus (Sheldrick, G. M. *SHELXTL-Plus*, Siemens Analytical X-ray Instruments Inc.: Madison, WI, 1989).

Details on the structure solution are given in Table I. The atomic coordinates and isotropic temperature factors are given in Tables II (**5a**) and III (**5b**), respectively. Further details of the crystal structure determinations are available on request from the Fachinformationszentrum Karlsruhe, Gesellschaft für wissenschaftlich-technische Information mbH, D-7514 Eggenstein-Leopoldshafen 2, Germany, on quoting the depository number CSD-55060, the names of the authors, and the journal citation.

Results and Discussion

Synthesis and Characterization. The synthesis of the iminophosphanes **5a,b** followed a known procedure for the generation of $\text{P}=\text{N}$ double bonds via metalation and subsequent salt elimination from aminohalophosphanes.¹⁶ The necessary Me_3C_5 -substituted precursors **3a,b** were prepared from PCl_3 and the primary amines **1a,b** in two steps according to standard procedures (Scheme I). Treatment of THF solutions of **3a,b** with the lithium amide **4** at -78 °C (1 equiv) and warming to ambient temperature then afforded the iminophosphanes **5a,b** (Scheme I). Pure

(12) Yoshifuji, M.; Shima, I.; Inamoto, N. *Tetrahedron Lett.* **1979**, *21*, 3963.


(13) Bowser, J. R.; Neilson, R. H.; Wells, R. L. *Inorg. Chem.* **1978**, *17*, 1882.

(14) Dickert, F. L.; Hellman, S. W. *Anal. Chem.* **1980**, *52*, 996.


(15) Benn, R.; Grondy, H.; Erker, G.; Aul, R.; Nolte, R. *Organometallics* **1990**, *9*, 2493.

(16) Niecke, E.; Gudat, D. *Angew. Chem., Int. Ed. Engl.* **1991**, *30*, 217.

Scheme I



Scheme II

Table I. Crystallographic Data and Summary of Data Collection and Refinement^{a,b}

	5a	5b
formula	C ₂₈ H ₄₄ NP	C ₁₉ H ₃₆ NPSi
fw	425.6	337.5
cryst syst	monoclinic	monoclinic
space group	P2 ₁ /n (No. 14)	P2 ₁ /n (No. 14)
cryst dimens, mm	0.2 × 0.2 × 0.4	0.55 × 0.6 × 0.7
cryst color	orange	yellow
temp, K	293	120
a, Å	11.974 (6)	8.466 (2)
b, Å	13.436 (4)	11.397 (2)
c, Å	17.309 (5)	22.082 (5)
β, deg	95.80 (3)	98.75 (2)
V, Å ³	2770	2106
Z	4	4
ρ _{calcd} , g cm ⁻³	1.02	1.07
μ, mm ⁻¹	0.11	0.18
F(000)	936	744
max 2θ, deg	45	50
no. of variables	271	216
no. of reflens coll	3950	3901
no. of unique data	3611	3695
R (merge)	0.023	0.119
no. of data used in refinement	1320	3398
F > 4σ (F)		
R	0.079	0.064
R _w	0.066	0.075
s	0.0003	0.0001
largest feature in final diff map, e/Å ³	0.32	0.87 (near P)

^aAll data were taken with graphite-monochromated Mo Kα radiation with λ = 0.71073 Å by using the ω-scan (5a) and profile fitting technique^b (5b), respectively. $R = \sum ||F_o| - |F_c|| / \sum |F_o|$ and $R_w = \sum (|F_o| - |F_c|)w^{1/2} / \sum |F_o|w^{1/2}$ with weighting scheme $w^{-1} = \sigma^2(F) + gF^2$.
^bClegg, W. *Acta Crystallogr.* **1981**, *A37*, 22–28.

products were isolated after filtration, removal of volatiles in vacuo, and recrystallization from pentane (5a) or distillation (5b) of the residue. 5a may be alternatively obtained in similar yields via treatment of the iminophosphane, ClP=N(C₆H₂tBu₃), with Me₅C₅Li.¹⁷

According to a different route, the trimethylsilylated derivative 5c was accessible via thermolytic (120 °C) elimination of Me₃SiCl¹⁶ from the chlorophosphane 7, which was prepared in situ by treatment of (Me₃Si)₂NPCl₂ (6) with 1 equiv of Me₅C₅Li (Scheme II). The product was collected by distillation and was redistilled for purification.

Identification and characterization of 5a–c were established from their mass spectra and analytical and multinuclear (¹H, ¹³C, ³¹P) NMR data. 5a,b were further characterized by X-ray crystallography.

X-ray Structural Investigations. Suitable single crystals of 5a,b were obtained by recrystallization from pentane at low temperature. Although 5c was also found to solidify upon cooling below –30 °C, no sufficient single crystals could be obtained in this case. Due to the low melting point (approximately –30 °C) of 5b, the structure determination also had to be performed at low temperature.

Table II. Atomic Coordinates (× 10⁴) and Equivalent Isotropic Displacement Parameters (Å² × 10³) of 5a

	x	y	z	U(eq) ^a
P(1)	295 (2)	1209 (2)	2481 (2)	49 (1)
N(1)	–704 (6)	1947 (5)	2459 (4)	44 (3)
C(1)	–353 (9)	–111 (7)	2513 (9)	53 (5)
C(2)	410 (13)	–412 (9)	3173 (8)	68 (7)
C(3)	1379 (12)	–740 (8)	2900 (12)	80 (7)
C(4)	1241 (14)	–741 (9)	2084 (12)	81 (7)
C(5)	213 (13)	–415 (8)	1835 (8)	65 (6)
C(6)	–1614 (8)	–164 (7)	2516 (8)	104 (6)
C(7)	147 (14)	–365 (10)	4001 (8)	171 (11)
C(8)	2424 (10)	–1043 (9)	3399 (10)	194 (11)
C(9)	2113 (11)	–1071 (10)	1562 (10)	206 (11)
C(10)	–370 (13)	–361 (9)	1010 (7)	143 (9)
C(11)	–627 (7)	3011 (7)	2449 (7)	41 (4)
C(12)	–527 (8)	3514 (7)	3171 (7)	45 (5)
C(13)	–302 (7)	4550 (8)	3152 (7)	49 (5)
C(14)	–236 (8)	5075 (7)	2475 (8)	40 (4)
C(15)	–487 (7)	4559 (7)	1796 (6)	40 (5)
C(16)	–712 (8)	3535 (7)	1739 (6)	43 (5)
C(17)	–628 (11)	3039 (8)	3966 (6)	62 (5)
C(18)	–831 (11)	3822 (8)	4591 (6)	130 (7)
C(19)	434 (9)	2468 (9)	4253 (6)	94 (6)
C(20)	–1652 (8)	2354 (8)	3942 (6)	83 (5)
C(21)	68 (9)	6178 (8)	2498 (9)	67 (6)
C(22)	–697 (13)	6752 (9)	2931 (10)	187 (11)
C(23)	–46 (17)	6644 (9)	1711 (9)	206 (13)
C(24)	1216 (11)	6321 (8)	2817 (12)	239 (15)
C(25)	–1026 (9)	3058 (8)	949 (6)	53 (5)
C(26)	–1395 (9)	3847 (8)	318 (6)	96 (6)
C(27)	–2019 (8)	2344 (8)	943 (6)	82 (5)
C(28)	3 (9)	2512 (8)	669 (6)	89 (6)

^aEquivalent isotropic U defined as one-third of the trace of the orthogonalized U_{ij} tensor.

Table III. Atomic Coordinates (× 10⁴) and Equivalent Isotropic Displacement Parameters (Å² × 10³) of 5b

	x	y	z	U(eq) ^a
P	286 (1)	3226 (1)	2015 (1)	21 (1)
Si	1926 (1)	2732 (1)	850 (1)	20 (1)
N	1404 (4)	3182 (3)	1531 (1)	26 (1)
C(1)	136 (4)	5027 (3)	2319 (2)	23 (1)
C(2)	1292 (4)	4365 (3)	2732 (2)	25 (1)
C(3)	450 (5)	3825 (3)	3171 (2)	28 (1)
C(4)	–1173 (4)	4102 (3)	3024 (2)	25 (1)
C(5)	–1384 (4)	4829 (3)	2503 (2)	23 (1)
C(1M)	548 (5)	5894 (3)	1860 (2)	34 (1)
C(2M)	3062 (4)	4501 (4)	2762 (2)	36 (1)
C(3M)	1172 (6)	3051 (4)	3694 (2)	46 (2)
C(4M)	–2508 (5)	3640 (4)	3336 (2)	42 (2)
C(5M)	–2944 (5)	5325 (4)	2189 (2)	40 (1)
C(11)	3716 (4)	1732 (3)	1038 (2)	28 (1)
C(12)	4301 (6)	1209 (4)	477 (2)	51 (2)
C(13)	5124 (5)	2269 (4)	1469 (2)	37 (1)
C(21)	2270 (4)	4094 (3)	390 (2)	25 (1)
C(22)	2336 (6)	3853 (4)	–286 (2)	41 (1)
C(23)	3714 (5)	4827 (3)	676 (2)	30 (1)
C(31)	196 (5)	1890 (3)	403 (2)	29 (1)
C(32)	–217 (5)	774 (3)	738 (2)	38 (1)
C(33)	–1275 (5)	2661 (4)	253 (2)	38 (1)

^aEquivalent isotropic U defined as one-third of the trace of the orthogonalized U_{ij} tensor.

(17) Reichert, F. Ph.D. Thesis, University of Bonn, Germany, 1989.




Figure 1. Molecular structure of **5a**. Selected bond distances (pm) and angles (deg): P(1)–N(1) 155.1 (8), P(1)–C(1) 194.0 (10), C(1)–C(2) 144.7 (19), C(2)–C(3) 136.9 (23), C(3)–C(4) 140.5 (29), C(4)–C(5) 133.7 (22), C(1)–C(5) 147.0 (21); C(1)–P(1)–N(1) 106.0 (4), P(1)–N(1)–C(11) 125.9 (6).

In crystalline **5a** the η^1 -coordination of the Me_5C_5 ligand is clearly evident (Figure 1). The *E*(*trans*) P–N double-bond distance (155 pm) and the C1–P–N angle (106°) lie in the expected ranges for *trans*-iminophosphanes,¹⁶ and the bond data of the $\text{P}=\text{NC}_6\text{H}_2\text{tBu}_3$ moiety are very similar to those of comparable compounds.¹⁶ While the planarity of C2–C5 (within experimental error) and the alternate C–C bond distances in the five-membered ring agree with a description as a cyclic diene bearing the phosphorus substituent in the allylic position,¹⁸ the allylic carbon exhibits severe distortions from ideal tetrahedral geometry. The P–C1–C2 (93°) and P–C1–C5 (91°) bond angles are distinctly contracted and the C2–C1–C6 (methyl) and C5–C1–C6 angles widened (122°) as compared to the expected angle of 109.4° . At the same time the C1–P distance (194 pm) is considerably longer than a normal P–C single bond,¹⁹ corresponding to a very low Pauling bond order (PBO)²⁰ of 0.76. The observed geometry indicates a state between ideal sp^3 and sp^2 hybridization at C1, which implies a low degree of s-electron density in the C1–P bond. Characteristic bond deformations of the same type, although somewhat less pronounced as for **5a**, have previously been reported for other η^1 - Me_5C_5 -substituted phosphorus $\text{p}(\pi)$ -systems,^{1,3} demonstrating that bonding is only marginally different in all of these cases.

In contrast to the situation described above, **5b** (Figure 2) exhibits a slightly asymmetric η^2 -attachment of the planar five-membered ring (the standard deviation for the best plane fitted through C1–C5 is $\sigma = 0.87$ pm) and represents the first example of Me_5C_5 multihapto bonding in a neutral phosphorus $\text{p}(\pi)$ -system. Four out of five bond distances in the five-membered ring lie in a narrow range between 139 and 143 pm with PBOs between 1.30 (C1–C5) and 1.52 (C3–C4). The P-bridged C1–C2 bond is slightly longer (144.4 pm, PBO 1.21), so that a marked equalization of bond lengths results as compared to **5a**. Of the ring carbons, C3–C5 are planar within experimental error, and the sum of C–C–C angles at C1 ($359 \pm 1^\circ$) and C2 ($358 \pm 1^\circ$) indicates that deviations of the adjacent methyl groups out of the ring plane are negligible. Hence, sp^2 hybridization may be assumed for all five ring atoms. In the PNSiR_3 fragment, rather short distances are found for both the Si–N bond (171 pm) and the P–N double bond (153.3 pm), which is shorter than the P–N bond in **5a** by 1.8 pm. Together with the large P–N–Si angle (153°), this suggests a state between pure sp and sp^2 hybridization




Figure 2. Molecular structure of **5b**. Selected bond distances (pm) and angles (deg): P–N 153.3 (3), P–C(1) 216.8 (4), P–C(2) 212.2 (4), C(1)–C(2) 144.5 (5), C(2)–C(3) 142.8 (5), C(3)–C(4) 139.9 (5), C(4)–C(5) 140.8 (5), C(1)–C(5) 142.5 (5), Si–N 170.9 (3); P–N–Si 153.3 (2).

for the two-coordinate nitrogen, with the P–N bond adopting partial triple-bond character. A Me_5C_5 –P interaction that is much weaker than in $[\eta^2\text{-Me}_5\text{C}_5\text{PNHtBu}]^+$ (**8**)⁶ is evidenced by the long P–C1 (212 pm) and P–C2 distances (217 pm), which correspond to extremely low PBOs of 0.32 and 0.38, respectively, that are presumably the longest P–C bonds known. Even if the angle between the cyclopentadienyl ring and the plane through P, C1, and C2 (86.6°) is smaller than in **8** (90.7°), the P–C3/C4 distances (263/264 pm) indicate that bonding between these atoms may be neglected.

The η^2 - Me_5C_5 -phosphorus bonding situation described here is a common feature of cyclopentadienyl π -complexes of main group elements.⁵ Very similar structural parameters have been found for η^2 - Me_5C_5 -substituted gerylenes²¹ and the phosphonium cation **8**.⁶ An analogous geometry was also calculated for an η^2 -cyclopentadienyl in $(\text{C}_5\text{H}_5)_2\text{P}^+$.¹⁸ The longer P–C1/C2 distances and the more pronounced leveling of the ring bonds in the neutral **5b** indicate, however, that bonding between the Me_5C_5 and PN fragments is considerably weaker than in the cationic **8**. In principle, the electronic structure of **5b** can be explained as an intramolecular donor/acceptor π -complex between a cyclopentadienyl anion and an iminophosphonium cation²² with a formal P–N triple bond. Stabilization of an iminophosphonium ion in an intramolecular σ -complex has recently been proposed for the compound $[\text{tBu}_2\text{PSe}_2][\text{PNC}_6\text{H}_2\text{tBu}_3]$ (**9**).²³ The η^2 -bonding situation in **5b** is in accord with both the low electrophilic character of the nitrogen in PN triple-bond systems²² as well as the tendency of phosphorus to adopt a low formal coordination number in its π -complexes.^{6,7,18}

NMR Spectroscopic Investigations. The solution ^1H and ^{13}C NMR spectra of **5a,b** exhibit sharp average signals for all ring carbons and methyl groups of the Me_5C_5 moiety down to the lowest accessible temperatures of -90°C . Such behavior had previously been reported as a common feature of compounds $\text{Me}_5\text{C}_5\text{P}=\text{ER}$,^{1–3} and was attributed to an elementotropic migration of the phosphorus fragment around the ring occurring with a very low activation energy.⁷ Since the activation barriers for exchange are frequently higher in the solid state than in solution,²⁴ the ^{13}C -CP/MAS spectra of solid **5a** were studied in the temperature range between 20 and -158°C . However, as in solution, the spectra lack any dynamically induced broadening or splitting for the resonances of the Me_5C_5 moiety (Figure 3), thus demonstrating that the fluxional behavior still persists in the solid

(18) Lee, T. J.; Schaefer, H. F., III; Magnusson, E. A. *J. Am. Chem. Soc.* **1985**, *107*, 7239.

(19) Corbridge, D. E. C. *The Structural Chemistry of Phosphorus*; Elsevier: Amsterdam, 1974.

(20) Pauling, L. *The Nature of the Chemical Bond*, 3rd. ed.; Cornell University Press: Ithaca, NY, 1960; pp 239 ff.

(21) Jutzi, P.; Hampel, B.; Hursthouse, M. B.; Howes, A. J. *Organometallics* **1986**, *5*, 1944.

(22) Niecke, E.; Nieger, M.; Reichert, F. *Angew. Chem., Int. Ed. Engl.* **1988**, *27*, 1715.

(23) Niecke, E.; Nieger, M.; Reichert, F.; Schoeller, W. W. *Angew. Chem., Int. Ed. Engl.* **1988**, *27*, 1713.

(24) Examples for this are cited in ref 15.




Figure 3. Olefinic region (center band) of the CP/MAS- $^{13}\text{C}\{^1\text{H}\}$ NMR spectrum at 20 °C (top) and -158 °C (bottom). The intense signal marked Cp^* is assigned to the five carbons of the cyclopentadienyl ring. The resonances of the aromatic carbons are marked as C(11)–C(16) in accord with the numbering scheme in Figure 1. Due to the low symmetry of the crystal, the equivalence of the ortho and meta positions of the aryl ring is removed, and separate resonances are observed for C(12),C(16) and C(13),C(15), respectively.

state.²⁵ Even if no mechanistic information is extractable from the NMR spectra, the X-ray structures of **5a,b** give strong evidence that the rotational motion proceeds by a sequence of 1,2 shifts, which is in accord with the general interpretation of the cyclopentadienyl dynamics as [1,5]-sigmatropic rearrangements.⁷

In regard to the similarity of the solution and solid-state ^{13}C spectra of **5a**, the significant differences in the ^{31}P spectra were quite unexpected (see Figure 4). In the solid, $\delta^{31}\text{P}(\text{iso})$ 269.9 ppm at 25 °C was shifted slightly downfield ($\Delta\delta$ 8 ppm) upon cooling to -158 °C, which is sufficiently explained as the consequence of thermal expansion effects. In solution, a much stronger deshielding of the signal occurred between 25 and -90 °C ($\Delta\delta$ 42 ppm); moreover, at 25 °C, $\delta^{31}\text{P}$ differed by 75 ppm between the solid state and hexane solution ($\delta^{31}\text{P}$ 195 ppm). Temperature- or medium-induced chemical shift variations of this magnitude are unprecedented for low-coordinate phosphorus compounds.²⁶ An explanation of the effects becomes feasible, postulating a nonsigmatropic mechanism for the ring migration for the solution case, which involves dynamic exchange between discernible η^1 - and η^2 -coordination isomers and is in analogy to the η^1 -/ η^2 - Me_5C_5 coordination isomerism in solid **5a,b**. Due to the fast exchange rate, the observed chemical shift should represent the population weighted average whose variation reflects the temperature-induced changes of individual populations. Support for this model is gained from the fact that the observed temperature dependence of $\delta^{31}\text{P}$ can be excellently fitted to an appropriate theoretical expression,²⁷ with the calculated shift of the η^1 isomer in very good agreement with $\delta^{31}\text{P}(\text{iso})$ in the solid state (see Figure 4). A "haptotropic" two-step mechanism for the cyclopentadienyl migration had been suggested earlier on kinetic grounds for the complexes of Me_5C_5 -substituted acyls with AlCl_3 ,²⁸




Figure 4. Temperature dependence of $\delta^{31}\text{P}$ of **5a** in solution (Δ) and the solid state (O). The straight line shows the results (with standard deviations) of a fit of the observed ^{31}P chemical shifts (in ppm) to a functional expression according to a two-site-exchange model²⁷ (T in K), $\delta^{\text{obsd}} = [\delta(\eta^1 \text{ isomer}) + \delta(\eta^2 \text{ isomer})e^{(\Delta S/R)}e^{(\Delta H/RT)}] / [1 + e^{(\Delta S/R)}e^{(\Delta H/RT)}]$, with $\delta(\eta^1 \text{ isomer}) = 268$ (30) ppm, $\delta(\eta^2 \text{ isomer}) = 149$ (20) ppm, $\Delta S = S(\eta^1 \text{ isomer}) - S(\eta^2 \text{ isomer}) = 24$ (8) $\text{J K}^{-1} \text{ mol}^{-1}$, $\Delta H = \Delta H(\eta^1 \text{ isomer}) - \Delta H(\eta^2 \text{ isomer}) = -6.1$ (1.3) kJ mol^{-1} .

Since in that case the high electrophilicity of the migrating double-bond system was considered vital for both the stabilization of the η^2 isomer and the observed depression of the activation barrier,²⁸ the electrophilic nature of PN multiple bonds¹⁶ presents an argument in favor of the proposed model. It must be noted, however, that the available experimental data in this case provide a rather indirect clue and that alternative mechanisms of the dynamics (e.g., cis/trans isomerization of the double bond) cannot be ruled out.

In contrast to **5a**, the solution spectra of **5b** exhibit only weak deshielding of $\delta^{31}\text{P}$ with increasing temperature; however, a marked change of the line shape becomes evident: the singlet observable at low temperatures eventually broadens and above ambient temperature splits into a triplet. The 1:1:1 intensity distribution and different line widths for the inner and outer lines are characteristic features of coupling with the ^{14}N nucleus²⁹


(25) A significant broadening which is observed at low temperature for the signals of the primary carbons of the $\text{C}_6\text{H}_2(\text{tBu})_3$ moiety may be assigned to slowing of the rotation of the tBu groups in the solid state.

(26) Measurements of $\delta^{31}\text{P}$ for a number of P alkyl-, amino-, and halogeno-substituted derivatives showed that in all cases $\delta^{31}\text{P}$ remained constant within 4 ppm between ambient temperature and -80 °C: Gudat, D. Ph.D. Thesis, University of Bielefeld, Germany, 1987.

(27) Dalling, D. K.; Zilm, K. W.; Grant, D. M.; Heeschen, W. A.; Horton, W. I. *J. Am. Chem. Soc.* **1981**, *103*, 4817.

(28) Childs, R. F.; Zeya, M. *J. Am. Chem. Soc.* **1974**, *96*, 6418. Childs, R. F. *Tetrahedron* **1982**, *38*, 567.

Scheme III



($^1J^{14}\text{N},^{31}\text{P} = 62 \text{ Hz}$). The appearance of this coupling, which is to our knowledge unprecedented in the chemistry of P–N compounds, indicates a near-linear geometry or very rapidly inverting structure for the two-coordinate nitrogen which may be correlated with the partial P–N triple-bond character in the solid state.

Nucleophilic Substitution Reactions. $\eta^1\text{-Me}_5\text{C}_5$ -substituted two-coordinate phosphorus compounds are known to undergo displacement of the Me_5C_5 group very easily by organic or organometallic nucleophiles.^{1,3,4,16} In regard to the extremely weak $\text{Me}_5\text{C}_5\text{-P}$ bonding in **5b**, similar behavior is expected for the corresponding $\eta^2\text{-Me}_5\text{C}_5$ isomers.

As a test of this hypothesis, the reactions of the N-silylated compounds **5b,c** with the organic nucleophile, $\text{LiC}_6\text{H}_2\text{tBu}_3$,¹² were investigated. In both cases, treatment of the iminophosphane with an equimolar amount of the organolithium reagent afforded the corresponding substitution products **10b,c** in almost quantitative yield (as monitored by ^{31}P NMR). Whereas **10b** was identified by comparison of its spectroscopic data with an authentic sample,³⁰ **10c** was isolated in good yield after distillative workup and characterized by analytical and spectroscopic methods. Further investigations of the substitution chemistry of **5a–c** including other organic and organometallic nucleophiles are currently under way (Scheme III).

Conclusions

The synthesis and structural characterization of the iminophosphanes **5a** and **5b** as stable P–N double-bond systems con-

taining discernible η^1 - or $\eta^2\text{-Me}_5\text{C}_5$ ligands verify the coordinational isomerism which had been postulated for these compounds. The results of the X-ray structures suggest localized $\text{Me}_5\text{C}_5\text{-P}$ σ -bonding in **5a**, whereas **5b** can be described as a cyclopentadienyl π -complex of a formally triple-bonded aminophosphenium ion. The existence of coordination isomerism further demonstrates that both σ - and π -bonded species are very close in energy, as had been predicted on the basis of calculations for cyclopentadienyl-aminophosphenium ions.³¹ The preference for either configuration can be controlled by differential effects as the variation of substituents, the promotion of the η^2 -coordination in **5b** being consistent with an increased electrophilicity of the P–N multiple bond as a consequence of the silyl substitution.³²

The ^{13}C -CP/MAS spectra of **5a** prove that the fluxionality of the Me_5C_5 ligand persists at low temperature in the solid state. Further, comparison of the solution and solid-state ^{31}P spectra gives evidence that the molecular structure in solution may differ from that in the crystal, which is tentatively explained by a model assuming a "haptotropic" mechanism involving dynamic η^1 -/ η^2 -coordination isomerizations for the cyclopentadienyl migration in solution.

In accord with the low $\eta^2\text{-Me}_5\text{C}_5\text{-P}$ bond order is the chemical reactivity of **5b,c**, which allows nucleophilic substitution of the cyclopentadienyl with preservation of the double-bond system. Hence, Me_5C_5 -substituted iminophosphanes can be regarded as promising synthons for novel P-functionalized iminophosphanes or iminophosphenium ions.

Acknowledgment. We thank the Deutsche Forschungsgemeinschaft (SFB 334) and the Fonds der Chemischen Industrie for financial support.

(31) Baxter, S. G.; Cowley, A. H.; Mehrotra, S. K. *J. Am. Chem. Soc.* **1981**, *103*, 5572.

(32) (a) Gudat, D.; Niecke, E.; Schoeller, W. W.; Rademacher, P. *J. Chem. Soc., Chem. Commun.* **1985**, 1050. (b) Gudat, D.; Niecke, E.; Sachs, W.; Rademacher, P. *Z. Anorg. Allg. Chem.* **1987**, *545*, 7.

(29) Pople, J. A. *Mol. Phys.* **1958**, *1*, 168.

(30) Romanenko, V. D.; Ruban, A. V.; Markovskii, L. N. *J. Chem. Soc., Chem. Commun.* **1983**, 187.