

the same square patterns, so that the multiplet structure is preserved, but all peaks are positive. This multiplet is less prone to distortions and signal cancellation, and the comparison of in-phase and antiphase multiplets allows one to extract coupling constants with high accuracy.^{16,17} Because the undesirable terms have vanished as if by a pixie's magic wand, we like to refer to this experiment as *pure in-phase correlation spectroscopy* (PICSY).

Acknowledgment. This research was supported by the Swiss Fonds National de la Recherche Scientifique (FNRS) and by the Commission pour l'Encouragement de la Recherche Scientifique (CERS). We gratefully acknowledge a sample of cyclo-(L-Pro-L-Pro-D-Pro) from Dr. W. Bermel, Karlsruhe, and Prof. H. Kessler, Garching.

(16) Titman, J. J.; Keeler, J. J. *Magn. Reson.* 1990, 89, 640.


(17) Huber, P.; Bodenhausen, G. Manuscript in preparation.

Dimerization of Aryl Radicals as a Termination Step in $S_{RN}1$ Aromatic Nucleophilic Substitution

Rofia Ettayeb,^{1a} Jean-Michel Savéant,^{*1b} and André Thiébaud^{*1a}

Laboratoire de Chimie et Electrochimie des Matériaux Moléculaires de l'ESPCI
Unité Associée au CNRS No. 429
10 rue Vauquelin, 75005, Paris, France
Laboratoire d'Electrochimie Moléculaire de l'Université de Paris 7, Unité Associée au CNRS No. 438, 2 place Jussieu
75251 Paris Cedex 05, France
Received October 1, 1992

Aromatic nucleophilic substitution stimulated by light, electrodes, solvated electrons, and electrochemically generated electron donors is a well-documented reaction.² A large body of evidence exists to support that the reaction follows the $S_{RN}1$ reaction mechanism depicted by the following scheme:



in which the aryl radical, R^\bullet , derived from the cleavage of the anion radical of the substrate (RX) plays a central role in the formation of the substituted product (RNu). In this chain process, reduction of the aryl radical not only by the homogeneous or heterogeneous electron donors serving as initiators but also by the anion radicals included in the propagation loop ($RNu^{\bullet-}$ and to a lesser degree $RX^{\bullet-}$) has been recognized, on the basis of electrochemical evidence,^{2b,d} as a major termination step. Another possible termination step is the coupling of two aryl radicals that would lead to the corresponding dimer:



So far no such dimer has ever been found among the reaction products of $S_{RN}1$ substitution reactions. It is, however, interesting to note in this connection that a small amount of dimer has been detected in the addition of electrochemically generated cyanophenyl radicals on styrene,⁴ a reaction that bears a close resemblance




Figure 1. Cyclic voltammometry of 4-chlorobenzonitrile and reaction products in liquid NH_3 (+0.25 M KBr) at $-38^\circ C$ at a 0.5 mm diameter gold disk electrode: (a) 4-chlorobenzonitrile alone in a 4.3 mM concentration; (b) 4-chlorobenzonitrile (35 mM) in the presence of 2-pyridinethiolate ions (47 mM); (c) 2-pyridyl 4-cyanophenyl sulfide (5.4 mM); (d) 4,4'-dicyanobiphenyl (0.5 mM). Scan rate: 0.2 V/s.

to the $S_{RN}1$ reaction. Although the absence of dimerization in $S_{RN}1$ reactions may be related to the likely smallness of the concentration of the R^\bullet radical in the chain reaction, it remains surprising that it has never been detected under the largely varied experimental conditions employed heretofore.⁵

We have found that dimerization of aryl radicals may be, under appropriate experimental conditions, a quite efficient termination step yielding substantial quantities of the R-R dimer.

Figure 1 shows the cyclic voltammometry of 4-chlorobenzonitrile in liquid ammonia alone (a) and in the presence of 2-pyridinethiolate ions as the nucleophile (b). By comparison with the voltammogram of the substitution product (c), it appears that extensive substitution occurs in the diffusion layer. Some hydrolysis product (benzonitrile) also appears as demonstrated by the most negative wave in b. More important, a new reversible wave appears positive to the RX and RNu waves and located at the same potential as the first wave of 4,4'-dicyanobiphenyl (d). The second wave of the dimer is merged with the first wave of RNu . It should be noted that the concentration of RX that allows the formation of the dimer to be observed is high (35 mM), much higher than the concentrations used in past studies, and that the excess of nucleophile is small. The formation of the dimer was

(1) (a) ESPCI. (b) Université de Paris.

(2) (a) Bunnett, J. F. *Acc. Chem. Res.* 1978, 11, 413. (b) Savéant, J.-M. *Acc. Chem. Res.* 1980, 13, 323. (c) Rossi, R. A.; Rossi, R. H. *Aromatic Substitution by the $S_{RN}1$ Mechanism*; ACS Monograph Series 178; The American Chemical Society: Washington, D.C., 1983. (d) Savéant, J.-M. *Adv. Phys. Org. Chem.* 1990, 26, 1.

(3) Kim, J. K.; Bunnett, J. F. *J. Am. Chem. Soc.* 1970, 92, 7463.

(4) Chami, Z.; Gareil, M.; Pinson, J.; Savéant, J.-M.; Thiébaud, A. *J. Org. Chem.* 1991, 56, 586.

(5) (a) The absence of dimerization has even been taken recently as strong evidence against the reality of the $S_{RN}1$ mechanism which should, in the view of the authors, be replaced by an $S_{RN}2$ mechanism in which substitution would take place directly in the $RX^{\bullet-}$ anion radical with no intermediacy of the aryl radical. (d) Denney, D. B.; Denney, D. Z. *Tetrahedron* 1991, 33, 6577.

checked by preparative-scale electrolysis. Constant current (60 mA) electrolysis at a stainless steel grid electrode (9 cm² geometric surface area, 324 mesh/cm²) of a solution of 4-chlorobenzonitrile (0.06 M) and 2-pyridinethiolate ions (0.114 M) in liquid NH₃ (+ 0.25 M KBr) at -38 °C produced, after ca. 3 h, 39% of 4,4'-dicyanobiphenyl, 43% of 2-pyridyl 4-cyanophenyl sulfide, 8% of benzonitrile, and 3% of unreacted starting material. The dimer was identified after extraction from the electrolysis solution by comparison of its NMR spectrum with that of an authentic sample.

Formation of the dimer in comparable quantities was also found to occur in cyclic voltammetry with 4-iodo- and 4-bromobenzonitrile as well as with 4-chloropyridine using the same nucleophile. It was also observed with 4-iodobenzonitrile as the substrate and with thiophenoxide ions as the nucleophile.

As noted before, we have observed that large concentrations of the substrate and small excesses of the nucleophile are favorable factors for the formation of the dimer. The anion radicals of the substrates that have been selected for the present study all undergo rapid expulsion of the halide ion. It thus appears that rapid formation of R[•] from RX⁻ and rapid destruction of RNU⁻ by large concentrations of RX in the propagation loop tend to decrease the reduction of R[•] by RNU⁻ and RX⁻. R[•] then mainly undergoes dimerization in competition with the attack of the nucleophile. The latter reaction should not, however, be too slow, otherwise the chain process would not be triggered in the solution and the reduction of R[•] at the electrode surface would annihilate its chances to dimerize.

Work is now in progress to investigate in a more quantitative manner the exact effect of all the factors we have just evoked on the dimerization yield. For the moment we may conclude that it is possible to devise experimental conditions, making dimerization appear as an efficient termination step in aromatic nucleophilic substitutions. This observation provides further evidence of the intermediacy of the aryl radical in the reaction in full agreement with the S_{RN}1 mechanism.

Bis(triphenylphosphine)platinum Cycloheptadienylium Fluoborate: The Tropylium Equivalent of Benzene

Zheng Lu, Khalil A. Abboud, and W. M. Jones*

Department of Chemistry, University of Florida
Gainesville, Florida 32611


Received July 16, 1992

To our knowledge, there are no prior reports of the tropylium analogue of benzyne, either free^{1,2} or complexed to a transition metal.³ At this time we report the preparation and properties, including an X-ray diffraction crystal structure, of **3**, a Pt(0) complex of the cycloheptadienylium ion, which we shall refer to as a complex of tropyne.


Preparation of the tropyne complex is outlined in Scheme I. Reaction of the mixture of 1-, 2-, and 3-bromocycloheptatriene (**1a-c**) with LDA led to, within the limits of detection by ³¹P NMR, total regiochemical reversal from the reported reaction with *t*-BuOK; cycloheptadienyne complexes **2a** and **2b** were formed⁴ (8:1) to the exclusion of the tetraene complex.⁵ Hydride abstraction from the mixture of **2a** and **2b** with Ph₃C⁺BF₄⁻ gave the tropyne complex **3** as red, air-stable crystals.

The structure assigned to **3** is based on elemental analysis, ¹H, ¹³C, ³¹P, and ¹⁹⁵Pt NMR,⁶ and an X-ray diffraction crystal

Scheme I



structure. As expected, chemical shifts of both protons and carbons of the seven-membered ring are all shifted downfield relative to their dienyne counterparts **2a** and **2b**. However, the chemical shifts of the atoms that are not bonded directly to platinum are significantly upfield from those of the tropylium ion⁷ (¹H, δ 9.55; ¹³C, δ 160.6), indicating electron donation from platinum into the tropylium ring. This may be due to either or both a positive inductive effect and electron donation into the π-framework. Normally, symmetry limits back-bonding from a transition metal into the LUMO of the orthogonal π-bond of an alkyne to an ineffective δ-bond.⁸ However, this is not the case for the tropyne complex because the tropylium ion has a vacant low-energy orbital⁹ that can mix with a platinum "d" orbital as pictured in **4** (represented in resonance terms by **5**). This may be significant in



the tropyne complex as evidenced by the ¹³C resonance at C4/6 (lowest field resonance of remote carbons),⁶ which correlates with the smallest LUMO coefficient and, hence, the lowest predicted electron density in **4** (0.12, HMO; 0.0, EHMO),¹⁰ a Pt-C coupling constant ($J_{\text{Pt-C}} = 458.5$ Hz) that is larger than expected from strain¹¹ and is consistent with contribution from resonance forms such as **5**,^{12,13} the ¹⁹⁵Pt chemical shift (-3788 ppm) which is further

(6) Properties of **3**: ¹H NMR (300 MHz, CD₂Cl₂, δ) 7.2 (m, Ph), 7.4 (m, Ph), 7.66 (m, H3/7, ³J_{Ph-H} = 51.9 Hz), 8.34 (m, H4/6), 8.64 (t, H5, ³J_{H-H} = 10.0 Hz); ¹³C NMR (75 MHz, CD₂Cl₂, δ) 174.6 (dd, C1/2, ¹J_{Pt-C} = 458.5 Hz, ²J_{Pt-C} = 83.4 Hz, ²J_{Pt-C} = 5.0 Hz), 150.0 (d, C4/6, ³J_{Pt-C} = 53.2 Hz, ⁴J_{Pt-C} = 10.5 Hz), 141.8 (s, C5), 135.6 (m, C3/7), 134.3, 133.1, 131.4 and 129.1 for Ph; ³¹P NMR (121 MHz, CD₂Cl₂, δ, 85% H₃PO₄ as reference) 21.5 (¹J_{Pt-P} = 3121.8 Hz); ¹⁹⁵Pt NMR (64 MHz, CD₂Cl₂, δ, Na₂PtCl₆ as reference) -3788. Anal. Calcd for C₄₃H₃₃PtP₂BF₄·1/2CH₂Cl₂: C, 55.68; H, 3.84. Found: C, 55.37; H, 3.80.

(7) (a) Gansow, O. A.; Schexnayder, D. A.; Kimura, B. Y. *J. Am. Chem. Soc.* **1972**, *94*, 3406. (b) Stewart, R. P., Jr.; Isbrandt, L. R.; Benedict, J. J.; Palmer, J. G. *J. Am. Chem. Soc.* **1976**, *98*, 3215.

(8) Cf. Coleman, J. P.; Hegedus, L. S.; Norton, J. R.; Finke, R. G. *Principles and Applications of Organotransition Metal Chemistry*; University Science Books: Mill Valley, CA, 1987; p 156.

(9) Coulson, C. A.; Streitwieser, A., Jr. *Dictionary of π-Electron Calculations*; Pergamon Press Ltd: New York, 1965. EHMO shows nodes at C4 and C6.¹³

(10) W. R. Winchester, Trinity University, private communication.

(11) For example, $J_{\text{Pt-C}} = 318, 395$, and 436 Hz for complexes of cycloheptyne, cyclohexyne, and cyclopentyne, respectively.³

(12) Chisholm, M. H.; Clark, H. C.; Ward, J. E. H.; Yasufuku, K. *Inorg. Chem.* **1975**, *14*, 893.

(13) Lu, Z.; Jones, W. M. *Organometallics*, submitted for publication.

(1) Roberts, J. D.; Simmons, H. E., Jr.; Carlsmith, L. A.; Vaughan, C. W. *J. Am. Chem. Soc.* **1953**, *75*, 3290.

(2) For a review of arynes, see: Hoffmann, R. W. *Dehydrobenzene and Cycloalkynes*; Academic Press: New York, 1967.

(3) For transition metal complexes of benzyne, see: Bennett, M. A.; Schwemlein, H. P. *Angew. Chem., Int. Ed. Engl.* **1989**, *28*, 1296.

(4) Lu, Z.; Abboud, K. A.; Jones, W. M. *Organometallics*, submitted for publication.

(5) Winchester, W. R.; Jones, W. M. *Organometallics* **1985**, *4*, 2228.