

Benzoannulated [2.2]Paracyclophanes: Synthesis and Electronic Properties

Oliver Reiser,[†] Burkhard König,[†] Klaus Meerholz,[‡] Jürgen Heinze,^{*‡} Thomas Wellauer,[§] Fabian Gerson,[§] Ron Frim,^{||} Mordecai Rabinovitz,^{||} and Armin de Meijere^{*†}

Contribution from the Institut für Organische Chemie der Universität Hamburg, Martin-Luther-King-Platz 6, 2000 Hamburg, Germany, Department of Chemistry, Hebrew University Jerusalem, Jerusalem, Israel, Institut für Physikalische Chemie der Universität Freiburg, Albertstrasse 21, 7800 Freiburg, Germany, and Institut für Physikalische Chemie der Universität Basel, Klingelbergstrasse 80, 4056 Basel, Switzerland

Received November 18, 1992


Abstract: Mono- and dibenzoannulated [2.2]paracyclophanes **12** and **1** were synthesized by palladium-catalyzed two- and four-fold alkenylation of vicinal di- and tetrabromides **5** and **6**, respectively, and subsequent electrocyclicization/dehydrogenation of the resulting (*E,Z,E*)-trienes. Further extensions of the annulated ring systems, leading to the tetrahydronaphthalene derivative **15**, the bis-terphenylene derivative **16**, and the benzobis[2.2]paracyclophane **18**, were achieved through derivatization of suitable substituents introduced with the alkene coupling component. Mono- and polyanions of some derivatives were generated and studied by ESR, ENDOR, and NMR spectroscopy, as well as by cyclic voltammetry. The assembly of mutually orthogonal π -systems in arene annulated [2.2]paracyclophanes allows the reversible incorporation of up to six additional electrons per molecule, which is the upper limit for hydrocarbons reported to date.

Introduction

It has long been recognized that electron-transfer phenomena play a fundamental role in biology as well as in chemistry.¹ Consequently, considerable efforts have been made to study the mechanisms which involve electron-transfer steps. Traditionally, it has been believed that conjugated π -systems are essential for the efficient movement of electrons through molecules. However, recent studies have suggested² that electron transfer through a σ -framework might also be effective.

We have been interested in the electronic properties of compounds with mutually orthogonal π -systems such as dibenzoannulated [2.2]paracyclophanes of the general structure **1**. These molecules consist of four biphenyl units, which are held rigidly in a perpendicular conformation and are, therefore, a priori nonconjugated. By studying anions of these molecules by ESR and NMR spectroscopy, as well as by cyclic voltammetry, we hoped to address two questions:

(1) Are the π -systems in the arene subunits of **1** sufficiently isolated to be reduced independently? (2) Do they interact sufficiently to allow electron transfer between them? This way, polyanions of **1** could serve as models for an electron storage system.



Several synthetic strategies toward benzoannulated [2.2]-

[†] Universität Hamburg. New Address: Institut für Organische Chemie der Universität Göttingen, Gustav Tammannstr. 2, D-3400 Göttingen, (West) Germany.

[‡] Universität Freiburg.

[§] Universität Basel.


^{||} Hebrew University, Jerusalem.

(1) (a) Cannon, R. D. *Electron Transfer Reactions*; Butterworth: London, 1981. (b) Taube, H. *Angew. Chem.* **1984**, *96*, 315; *Angew. Chem., Int. Ed. Engl.* **1984**, *23*, 329.

(2) (a) Wegner, G. *Angew. Chem.* **1981**, *93*, 352; *Angew. Chem., Int. Ed. Engl.* **1981**, *20*, 348. (b) Hanack, M. *Nachr. Chem., Tech. Lab.* **1980**, *28*, 632.

paracyclophanes have been explored. With one exception,³ a [2.2]paracyclophane derivative was chosen as the starting material with subsequent arene annulation onto the ethano bridges, thus forming the desired benzoannulated [2.2]paracyclophane structure **1**. At first sight a [4 + 2] cycloaddition of dienes to the readily accessible⁴ [2.2]paracyclophane-1,9-diene (**2**)⁵ would seem the most straightforward approach to such skeletons. Diene **2**, however, failed to react with butadiene,⁶ substituted butadienes, and 5,5-dimethoxy-1,2,3,4-tetrachlorocyclopentadiene.⁷ The first syntheses⁸ of diene annulated [2.2]paracyclophanes were achieved via intermediates obtained by intermolecular trapping of in situ generated 1,2-benzo[2.2]paracyclophane-9-yne with dienes.^{9,10} A Diels–Alder addition of propargyl aldehyde to 1-vinyl- and 1,9(10)-divinyl[2.2]paracyclophanes afforded another approach to compounds of type **1**.⁴

Recently, we developed a general method for annulating six-membered carbocycles onto 1,2-dibromocycloalkenes¹¹ by Heck-type¹² alkenylation and subsequent cyclization/dehydrogenation of the resulting (*E,Z,E*)-1,3,5-trienes.



Preliminary studies¹³ had previously established that this protocol is applicable to vicinal dibromoalkenes derived from

(3) Jacobson, N.; Boekelheide, V. *Angew. Chem.* **1978**, *90*, 49–50; *Angew. Chem., Int. Ed. Engl.* **1978**, *17*, 46–47.

(4) Stöbbe, M.; Reiser, O.; Näder, R.; de Meijere, A. *Chem. Ber.* **1987**, *120*, 1667.

(5) Dewhirst, K. C.; Cram, D. J. *J. Am. Chem. Soc.* **1958**, *80*, 3115.

(6) (a) Hopf, H.; Psiorz, M. *Chem. Ber.* **1986**, *119*, 1836. (b) Psiorz, M. Dissertation, Universität Braunschweig, Braunschweig, Germany, 1983.

(7) Dimethyl tetrazine-3,6-dicarboxylate cleanly undergoes Diels–Alder reactions with **2** in good yields: König, B. Dissertation, Universität Hamburg, Hamburg, Germany, 1991.

(8) Chan, C. W.; Wong, H. N. C. *J. Am. Chem. Soc.* **1985**, *107*, 4790–4791.

(9) Chan, C. W.; Wong, H. N. C. *J. Am. Chem. Soc.* **1988**, *110*, 462–469.


(10) de Meijere, A.; Heinze, J.; Meerholz, K.; Reiser, O.; König, B. *Angew. Chem.* **1990**, *102*, 1443–1444; *Angew. Chem., Int. Ed. Engl.* **1990**, *29*, 1418–1419.

(11) Lansky, A.; Reiser, O.; de Meijere, A. *Synlett* **1990**, *1*, 405.

(12) Heck, R. F. *Org. React. (NY)* **1983**, *27*, 1.

(13) Reiser, O.; Reichow, S.; de Meijere, A. *Angew. Chem.* **1987**, *99*, 1285–1286; *Angew. Chem., Int. Ed. Engl.* **1987**, *26*, 1277–1278.

Scheme I



(a) Br₂, CCl₄, Δ. (b) KO^tBu, TBME, room temperature. (c) Br₂, CHCl₃, Δ. (d) KO^tBu, TBME, room temperature.

[2.2]paracyclophane. In this paper we report on the full utility of this strategy as an efficient entry to a wide variety of substituted dibenzo[2.2]paracyclophanedienes of type **1** and on our studies of their electronic properties.

Results and Discussion

Preparation of Benzoannulated [2.2]Paracyclophanes. 1,2-Dibromo[2.2]paracyclophane-1-ene (**5**) and 1,2,9,10-tetrabromo[2.2]paracyclophane-1,9-diene (**6**) were envisioned as suitable starting materials for the synthesis of benzoannulated [2.2]paracyclophanes. Hopf and Psiorz^{6a} first synthesized **5** by monobromination of [2.2]paracyclophane (**3**) followed by a sequence of dehydrobromination, bromination, and dehydrobromination steps. In our hands, however, it proved to be difficult to control the initial photobromination on a large scale. Mixtures of monobrominated and polybrominated compounds were always obtained, from which the monobromo derivative could not be separated completely.

The observation that the four-fold photobromination of **3** with bromine⁵ affords the geminal 1,1,9,9- and 1,1,10,10-tetrabromides **4b** along with considerable amounts (20–30%) of the geminal dibromide **4a** led us to the development of an alternate route to both **5** and **6**. In a sequence of dehydrobromination, exhaustive bromination, and dehydrobromination, a mixture of dibromomonoene **5** and tetrabromodiene **6** was obtained, which could easily be separated by crystallization due to their vastly different solubilities in chloroform (Scheme I).


Compounds **5** and **6**, thus accessible in multigram quantities, were subjected to palladium-catalyzed coupling reactions with various alkenes **7a–j**. According to the protocol originally suggested by Jeffery,¹⁴ which calls for addition of a phase-transfer catalyst,¹⁵ the two- and four-fold coupling products were obtained in respectable yields in most cases (Scheme II and Table I). Propene and ethylene gave poor results under the conditions employed. While no product was obtained from **5** with propene, **6** with ethene gave the four-fold adduct **9a**, which was converted to **1a** (overall yield 14%). **1a** could be obtained more conveniently and in higher overall yield (21%) by using vinyltrimethylsilane as an ethene equivalent in the coupling step. Under the basic conditions employed, the trimethylsilyl groups were cleaved off in situ to a large extent (>95% according to ¹H NMR).

Consistently better results were obtained by using alkenes with electron-withdrawing substituents (entries 1–3, 7–13 in Table I). Unfortunately, the bis-triene **9d** derived from coupling with acrolein proved to be rather unstable, and all attempts to realize further conversion to the tetraaldehyde **1d** failed. Methylacrylate (entries 1, 8) and various substituted styrenes (entries 2, 3, 9–13), however, cleanly and stereoselectively gave the corresponding (*E,Z,E*)-trienes **11** and bis-trienes **9**, which could be isolated without any apparent isomerization or decomposition. Subsequent

(14) Jeffery, T. *Tetrahedron Lett.* **1985**, 26, 2667–2670.


(15) Contrary to the procedure in ref 14, we used the cheaper tetrabutylammonium bromide instead of the chloride with good results.

Scheme II. (For Designation of Substituents R and Yields, See Table I)



(a) Pd(OAc)₂, Bu₄NBr, K₂CO₃ or NaHCO₃, DMF. (b) DDQ (or S), xylene, Δ.

Scheme III



(a) LiAlH₄, THF, Δ. (b) PBr₃, C₆H₆, room temperature. (c) Zn, THF,))) . (d) Diethyl fumarate, room temperature.

thermal 6π-electrocyclization of products **9** and **11** in the presence of sulfur or DDQ resulted in the formation of dibenzoannulated [2.2]paracyclophanes **1** and monobenzoannulated analogues **12**.

Interestingly, no monoalkenylated products were isolated. Competition experiments in a recent study¹¹ have revealed that the second coupling step in a palladium-catalyzed alkenylation of 1,2-dibromocyclohexene is about 50 times faster than the first step yielding 1-bromo-1,3-dienes. In agreement with these findings, the only intermediates we observed in the four-fold coupling of **6** were the two-fold alkenylated adducts **8**, which could be cyclized to give monobenzoannulated dibromo[2.2]paracyclophanes **10**, as established for compounds **8e** and **8f**.

This ring annelation method leads to 1,2-disubstituted arenes, which offer themselves for further elaboration, especially toward [2.2]paracyclophane derivatives with extended mutually orthogonal polycyclic π-systems.

A particularly attractive intermediate for further ring annelation appeared to be the *o*-xylylene annelated paracyclophane **14**, for which **13** ought to be an ideal precursor, since transformation of *o*-bis(bromomethyl)benzenes to *o*-xylenes is well precedented.^{16,17}

Reduction of the diester **12e** with lithium aluminum hydride to the bis-benzyl alcohol and subsequent nucleophilic substitution with phosphorus tribromide cleanly gave **13** in 70% overall yield. Attempts to isolate a tricarbonyliron(0) complex of **14** by treating **13** with enneacarbonyliron(0)¹⁸ failed. By reacting **13** with zinc under ultrasound activation,¹⁹ however, **14** could be generated in situ and trapped with diethyl fumarate to give the [4 + 2] cycloadduct **15** in 70% yield (Scheme III).

(16) Cf. (a) Johnson, B. F. G.; Lewis, J.; Thompson, D. J. *Tetrahedron Lett.* **1974**, 3789. (b) Roth, W. R.; Main, J. D. *Tetrahedron Lett.* **1967**, 2053.


(17) Cf. (a) Alder, K.; Fremery, M. *Tetrahedron* **1961**, 14, 190. (b) Han, B. H.; Boudjouk, P. *J. Org. Chem.* **1982**, 47, 751.

(18) Cf. Liu, L.; Yang, B.; Katz, T. J.; Pointdexter, M. K. *J. Org. Chem.* **1991**, 56, 3769–3775 and references cited therein.


(19) Hopf, H. *Naturwissenschaften* **1983**, 70, 349.

Table I. Palladium-Catalyzed Coupling Reactions of **5** and **6** with Various Alkenes **7** and Subsequent Cyclization/Dehydrogenation of the Resulting (*E,Z,E*)-1,3,5-Trienes **9** and **11**

| entry | starting material | 7 | alkene R | temp, °C (time) | product triene | yield, % | oxidant | product arene | yield, % |
|-------|-------------------|---|--|-----------------|----------------|----------|---------|---------------|-----------------|
| 1 | 5 | e | CO ₂ Me | 50 (12h) | 11e | 44 | DDQ | 12e | 45 |
| 2 | 5 | f | C ₆ H ₅ | 100 (3d) | 11f | 58 | DDQ | 12f | 53 |
| 3 | 5 | i | <i>p</i> -CO ₂ Me-C ₆ H ₄ | 70 (1d) | 11i | 57 | S | 12i | 65 |
| 4 | 5 | c | CH ₃ | 100 (2d) | 11c | 0 | | | |
| 5 | 6 | a | H | 100 (2d) | 9a | 24 | S | 1a | 58 |
| 6 | 6 | b | SiMe ₃ | 40 (5d) | 9b | <i>a</i> | S | 1a | 21 ^b |
| 7 | 6 | d | CHO | 40 (2d) | 9d | 38 | S | 1d | 0 |
| 8 | 6 | e | CO ₂ Me | 70 (12h) | 9e | 48 | S | 1e | 43 |
| 9 | 6 | f | C ₆ H ₅ | 100 (3d) | 9f | 55 | S | 1f | 50 |
| 10 | 6 | g | <i>p</i> -F-C ₆ H ₄ | 70 (2d) | 9g | <i>a</i> | S | 1g | 20 ^b |
| 11 | 6 | h | <i>p</i> - ^t Bu-C ₆ H ₄ | 100 (2d) | 9h | 55 | S | 1h | 45 |
| 12 | 6 | i | <i>p</i> -CO ₂ Me-C ₆ H ₄ | 70 (36h) | 9i | 58 | S | 1i | 80 |
| 13 | 6 | j | <i>p</i> -C ₆ H ₅ -C ₆ H ₄ | 100 (4d) | 9j | 15 | DDQ | 1j | 31 |

^a Not isolated. ^b Yield based on **6**.**Scheme IV**(a) LiAlH₄, THF, Δ. (b) PBr₃, C₆H₆, room temperature. (c) PhLi, Et₂O.

Since coupling of **6** with styrenes and cyclization/oxidation provides an access to *o*-terphenyl annelated [2.2]paracyclophanes **1f–j**, the photocyclization¹⁸ of these to terphenylene annelated derivatives was tested. Irradiation of **1f** at 254 nm in the presence of iodine caused the disappearance of the starting material; the isolated product, however, proved to be so sparsely soluble that it could not be sufficiently characterized. The tetra-*tert*-butyl-substituted derivative **16**, obtained from photocyclization of **1h** in 66% isolated yield, has a better solubility and could be characterized by its ¹H NMR spectrum and by cyclic voltammetry (see below).




The minimal solubility of **16** in organic solvents prevented, however, recording of a ¹³C NMR spectrum. For unknown reasons, a mass spectrum of **16** could not be obtained by electron impact, field desorption, and fast atom bombardment ionization techniques.

Finally, the first benzobis[2.2]paracyclophane **18** was synthesized starting from the di-*p*-carbomethoxyphenyl derivative **12i** (Scheme IV). Reduction of **12i** with LiAlH₄, and conversion of the resulting diol with phosphorus tribromide, gave the bis-bromomethyl derivative **17**. Upon treatment of **17** with phenyllithium in diethyl ether, **18** was isolated in 8% yield, along with dehalogenated and polymeric materials.


Despite this low yield, **18** could be completely purified and its constitution was unambiguously verified by its high-resolution mass and ¹H NMR spectrum. Prominent in the latter spectrum

is an AA'BB' system at 6.64 ppm, which is characteristic for [2.2]paracyclophanes.¹⁹ The signal for the two protons of the doubly annelated benzene appears as a singlet at 7.79 ppm, while the eight protons of the ethano bridges are represented by another singlet at 3.13 ppm.

Spectroscopic and Cyclic Voltammetric Studies of Redox Processes in Benzoannelated [2.2]Paracyclophanes. NMR studies were carried out on anions derived from the parent compound **1a**, the previous reported diphenyl derivative **19**,⁴ and the tetraphenyl derivative **1f**. These compounds were each treated with lithium, sodium, or potassium in THF-*d*₈ at -78 °C. In all cases, deeply colored solutions were obtained, which were periodically analyzed by ¹H and ¹³C NMR spectroscopy in a temperature range between 0 and -80 °C.



At no point were NMR signals observed for the reduction products of the unsubstituted dibenzo[2.2]paracyclophanediene **1a**, regardless of the alkali metal employed. This failure indicates that no diamagnetic species, in particular no stable singlet dianion, was formed from **1a**. This complements the ESR studies of **1a**,²⁰ in which only a radical monoanion, but no paramagnetic dianion, was observed.



In contrast, after 2-weeks exposure to sodium or potassium mirror at -78 °C, the solutions of **1f** and **19** showed NMR signals; these were, however, too broad to be reliably analyzed. As proved

(20) Gerson, F.; Martin, W. B., Jr.; Wong, H. N. C.; Chan, C. W. *Helv. Chim. Acta* **1987**, *70*, 79.

Table II. Reversible Reduction Potentials E° (V) of Some Arene Annulated [2.2]Paracyclophanedienes and Related Model Compounds as Measured by Cyclic Voltammetry^a

| | 1a | 1h | 1j | 16 | 19 | 20 | 21 | 22 |
|-----------------|-----------|-----------|--------------------|-----------|-----------|-----------|-----------|-----------|
| $E_{1/2}^{(1)}$ | -2.68 | -2.54 | | -2.41 | -2.54 | -2.68 | -2.62 | -2.42 |
| $E_{1/2}^{(2)}$ | -2.98 | -2.61 | | -2.50 | -2.60 | -3.18 | -2.72 | -2.97 |
| $E_{1/2}^{(3)}$ | | -2.80 | | -2.75 | -2.91 | | | |
| $E_{1/2}^{(4)}$ | | -2.89 | -2.43 ^b | -3.02 | -3.21 | | | |
| $E_{1/2}^{(5)}$ | | | -3.08 | | | | | |
| $E_{1/2}^{(6)}$ | | | -3.18 | | | | | |

^a Recorded with a Pt electrode for 10^{-4} to 10^{-5} M solutions in a dimethylamine/THF (1:1) mixture. All potentials were determined as the average of the cathodic and the anodic peak potentials and are expressed in V vs Ag/AgCl (calibration with $[\text{Cp}_2\text{Co}]^+/\text{Cp}_2\text{Co}$). The temperature was kept between 233 and 198 K. ^b This is a four-electron reduction step.

by ESR spectroscopy, both **19** and **1f** form the triplet dianions with one electron located in each of the annulated biphenyl and *o*-terphenyl systems, respectively;²¹ it might be concluded that the observed NMR signals are derived from the corresponding diamagnetic singlet dianions. An analysis of the temperature-dependent ESR signals pointed to the triplet being the ground state of these dianions with a singlet only about 2 kJ/mol higher. This finding is consistent with the observation of such singlet dianions by NMR spectroscopy, as the Boltzmann distribution would lead to a detectable fraction of them in equilibrium with the triplet species and the presence of the latter ones in solution would cause the observed line broadening.

After prolonged contact of **1f** and **19** with potassium metal, the NMR signals disappeared, indicating the formation of paramagnetic trianions which were also observed by ESR/ENDOR spectroscopy.²¹ Finally, the anion solutions were quenched with dioxigen. In all cases the starting material was recovered without any trace of decomposition, ensuring that the observed species were indeed reversibly formed anions of benzoannulated [2.2]paracyclophanes.


The cyclic voltammetric studies of some derivatives of **1** (Table II) complement the results of our ESR and NMR measurements. In all cases, the observed reduction steps are reversible, even for the highly charged species of **1j** (vide infra).

The first reduction step of **1a** occurs at $E_{1/2}^{(1)} = -2.68$ V vs SCE, which is a substantially less negative potential than that of [2.2]paracyclophane ($E_{1/2}^{(1)} = -3.00$ V vs SCE²²), as the uptake of an electron becomes easier upon extension of the π -system vs SCE, albeit with mutually orthogonal subunits. Consequently, the $E_{1/2}^{(1)}$ value of **1a** is almost equal to that of *o*-terphenyl (**21**), in which the three rings are also almost orthogonal to each other.²³ **1a** may be regarded as a sterically constrained *o*-terphenyl derivative which is bridged in the para position of its terminal phenyl units by another orthogonal benzene ring.

Although a dianion of **1a** could not be observed either by ESR or NMR spectroscopy, the cyclic voltammogram revealed a second reduction step at $E_{1/2}^{(2)} = -2.98$ V. This value is now significantly more negative than the corresponding potential of *o*-terphenyl (**21**) with $E_{1/2}^{(2)} = -2.72$ V, which demonstrates a strong Coulombic repulsion in **1a** due to the strong π - π interaction between the parallel and face to face oriented benzene rings at a distance of only 314 pm.²⁴ This π - π interaction is weaker in **21**, and consequently, its second reduction step is separated from the first by only 100 mV.

The tetrakis(*p*-*tert*-butylphenyl) derivative **1h** and the diphenyl derivative **19** of **1a**, as well as the bis-triphenylene annulated compound **16**, could be reduced to their tetraanions. The first two reduction potentials, $E_{1/2}^{(1)}$ and $E_{1/2}^{(2)}$, lie in the same range as the $E_{1/2}^{(1)}$ values of the corresponding subunits *o*-terphenyl (**21**), biphenyl (**20**), and triphenylene (**22**), respectively. This

fact supports the notion that these subunits are reduced separately and almost independently of each other, as previously found in the ESR/ENDOR studies for **1h** and **19**.²¹ Due to the rather slight interactions between the two subunits in **1h**, **16**, and **19**, the potentials, $E_{1/2}^{(1)}$ and $E_{1/2}^{(2)}$, of the three compounds are almost degenerate.



The third reduction potential of **1h**, $E_{1/2}^{(3)} = -2.80$ V, is considerably less negative than the $E_{1/2}^{(1)}$ value of -3.00 V for [2.2]paracyclophane (**3**), given the fact that both annulated *o*-terphenyl units are already charged. Nevertheless, the ESR spectrum of the radical trianion of **1h** proved unambiguously that the third electron is accommodated in the central paracyclophane subunit. In this way, the distances between the three charges are maximized. Similar to the case of **1h**, the third electron in **19** should be located in the central paracyclophane unit, all the more as the second reduction potential of **20**, $E_{1/2}^{(2)} = -3.18$ V, is markedly more negative than the corresponding value of -2.72 V for **21**. Indeed, the third reduction potential of **19**, $E_{1/2}^{(3)} = -2.91$ V, is rather close to that of **1h** (-2.80 V), and the location of the third electron in the paracyclophane subunit was confirmed by the ESR spectrum of the radical trianion of **19**.²¹ On the other hand, the fourth reduction potential of **19**, $E_{1/2}^{(4)} = -3.21$ V, is distinctly more negative than the corresponding value of -2.89 V for **1h**; this difference parallels that in the second reduction potentials, $E_{1/2}^{(2)}$, of the respective lateral subunits **20** (-3.18 V) and **21** (-2.72 V). It must, therefore, be assumed that the fourth electron in **1h** and **19** is mainly accommodated in one of these subunits, a conclusion which is difficult to confirm by ESR spectroscopy.

No ESR evidence is available for the paramagnetic anions of **16**, as the spectra observed upon alkali-metal reduction of this compound were hardly interpretable, presumably due to the occurrence of two or more different species. As mentioned above, the first two reduction steps of **16** (see Figure 1) at $E_{1/2}^{(1)} = -2.41$ and $E_{1/2}^{(2)} = -2.50$ V compare favorably with the value $E_{1/2}^{(1)} = -2.42$ V for **22**. This comparison indicates that the two steps involve a subsequent uptake of electrons into each of the triphenylene subunits of **16**. It is less certain to which subunit of **16** the admission of the third and fourth electrons should be assigned. The finding that the third reduction potential of **16**, $E_{1/2}^{(3)} = -2.75$ V, is close to the corresponding value of **1h** (-2.80 V) suggests that, here too, the central paracyclophane subunit represents the site of the third electron in the radical trianion of **16**. The fourth electron, on the other hand, may enter the lateral triphenylene subunits, as the $E_{1/2}^{(4)}$ value of -3.02 V for **16** is similar to the second reduction potential, $E_{1/2}^{(2)}$, of **22** (-2.97 V).

The tetrakisbiphenyl-substituted derivative **1j** can even be reduced to its hexaanion. The first four reduction steps occur all

(21) For a detailed discussion of this ESR study see: de Meijere, A.; Gerson, F.; König, B.; Reiser, O.; Wellauer, T. *J. Am. Chem. Soc.* **1990**, *112*, 6827.

(22) Jund, R.; Lemoine, P.; Gross, M. *Angew. Chem.* **1982**, *94*, 312; *Angew. Chem., Int. Ed. Engl.* **1982**, *21*, 305.

(23) Meerholz, K.; Heinze, J. *J. Am. Chem. Soc.* **1989**, *111*, 2325-2326.

(24) Wong, H. N. C.; Chan, C. W.; Mak, T. C. W. *Acta Crystallogr.* **1986**, *C42*, 703-705.




Figure 1. Cyclic voltammogram for the reduction of **16** in DMA/THF (1:1), 0.1 M TBABr, scan rate 100 mV s⁻¹, *T* = 213 K.

in the potential range between -2.38 and -2.48 V. As the lateral pentaphenyl units in **1j** are even more extended π -systems than *o*-terphenyl (**21**), its first two reduction potentials may be expected to be almost degenerate; therefore, a two-electron reduction step for each of these subunits is observed. Furthermore, it is assumed that the fifth excess charge is stored in the central paracyclophane subunit, although the small difference of 0.1 V between $E_{1/2}^{(5)}$ and $E_{1/2}^{(6)}$ may be taken as an indication that the fifth and the sixth electron are actually located in the lateral pentaphenyl subunits.

Conclusions

The results of the ESR, NMR, and cyclovoltammetric studies provide affirmative answers to both questions put forward in the introduction. On the one hand, the ESR/ENDOR studies of the radical anions, triplet dianions, and radical trianions of **1h**, **1j**, and **19** indicate that the π -subsystems in these bis-annelated [2.2]-paracyclophanes are sufficiently isolated to take up electrons one by one in consecutive redox steps.²¹ On the other hand, cyclic voltammograms of **1h**, **1j**, and **16** point to a substantial interaction between their "isolated" π -systems, as the reduction potentials attributed to these systems are considerably less negative than those of the corresponding parent hydrocarbons **1a**, **20**, **21**, and **22**, which are subunits in **1h**, **1j**, and **16**. Benzoannelated [2.2]-paracyclophanes are therefore highly efficient electron acceptors. The unusual geometry of their carbon framework allows reversible acceptance of four to six electrons, thus making especially **1j** an attractive model for a charge storage system.

Experimental Section

General Methods. Melting points were determined on an electrothermal melting point apparatus and are uncorrected. Infrared spectra were recorded by Perkin Elmer 297 and 399 spectrophotometers. ¹H NMR spectra were taken on Bruker AM-250 (250 MHz), WH-270 (270 MHz), and WM-400 (400 MHz) spectrometers at ambient temperature. Data are reported as follows: Chemical shifts in ppm from internal tetramethylsilane on the δ scale; multiplicity (b = broad, s = singlet, d = doublet, t = triplet, q = quartet, and m = multiplet), integration, coupling constant (Hz), and assignment. Chemical shifts are given in ppm on the δ scale, with the solvent resonance employed as the internal standard (chloroform at 7.24 ppm, benzene-*d*₅ at 7.15 ppm, and DMSO-*d*₅ at 2.49 ppm). ¹³C NMR were recorded on Bruker AM-250 (62 MHz), WM-270 (67 MHz), and WM-400 (100 MHz) spectrometers at ambient temperature; δ = 77.0 ppm for CDCl₃. Ultraviolet spectra were recorded on a Perkin-Elmer-Hitachi spectrometer. Mass spectrometry analyses were performed on a Varian MAT CH-7 spectrometer. Combustion analyses were carried out by Microanalytical Laboratory, University of Hamburg, and Beller Microanalytical Laboratory, Göttingen. Analytical thin-layer chromatography (TLC) was performed using Merck silica gel 60-F₂₅₄ on aluminum foil. Liquid chromatography was performed using

a forced flow (flash chromatography) of the indicated solvent system on Merck silica gel 60 (70–230 mesh). When necessary, solvents and reagents were purified prior to use by standard laboratory procedures.²⁵ All voltammetric measurements were performed with dimethylamine/THF solutions (1:1 mixture) and carried out with a PAR potentiostat Model 173 and a PAR universal programmer Model 175. Cyclic voltammograms were recorded with a Model HP 7004B X-Y instrument. A three-electrode configuration was employed throughout. Details are described elsewhere.²⁶ Experimental conditions for the ESR/ENDOR studies were indicated previously.²¹

1,2,9,10-Tetrabromo[2.2]paracyclophane-1,9-diene (6). A solution of 14.3 g (89.5 mmol) of bromine in 100 mL of CHCl₃ was added dropwise to a solution of 16.2 g (44 mmol) of 1,9(10)-dibromo[2.2]paracyclophane-1,9-diene⁵ in 150 mL of CHCl₃ while refluxing for 3 h. The precipitate was collected by filtration, the filtrate was concentrated to a volume of 30 mL, 100 mL of hexane was added, and the precipitated solid was again removed by filtration. The combined residues were dried in vacuo yielding 21.7 g (71%) of 1,1,2,9,9(10),10-hexabromo[2.2]paracyclophane. A small sample was recrystallized from CCl₄ for analytical characterization. ¹H NMR (270 MHz, CDCl₃): δ 5.97 [s, H-2,9,(10)], 7.03 (m, 8 H, H-aromatic). MS (70 eV) *m/z* (%): 680/682/684 (2/3/2, M⁺), 597/599/601/603/605/607 (2/7/14/13/7/1, M⁺ - Br), 518/520/522/524/526 (7/28/40/26/7, M⁺ - 2 Br), 439/441/443/445 (8/24/22/9, M⁺ - 3 Br), 360/362/364 (10/20/10, M⁺ - 4 Br), 281/283 (16/15, M⁺ - 5 Br), 202 (100, M⁺ - 6 Br). Anal. Calcd for C₁₆H₁₀Br₆: C, 28.19; H, 1.48; Br, 70.33. Found: C, 27.97; H, 1.43; Br, 70.10.

A suspension of 21.7 g (31.8 mmol) of 1,1,2,9,9(10),10-hexabromo[2.2]paracyclophane and 14.0 g (125 mmol) of potassium *tert*-butoxide (KO^tBu) in 200 mL of methyl *tert*-butyl ether (TBME) was stirred at room temperature for 12 h. The reaction mixture was evaporated to dryness and the solid residue continuously extracted with chloroform for 7 days. The precipitate in the extraction solution was filtered off, washed with 50 mL of cold chloroform, and dried in vacuo yielding 11.6 g (70%) of **6**, as tiny yellow crystals. ¹H NMR (270 MHz, CDCl₃): δ 6.74 (s, H-aromatic). ¹H NMR (270 MHz, C₆D₆): δ 6.09 (s). MS (70 eV) *m/z* (%): 516/518/520/522/524 (3/6/13/15/9, M⁺). Anal. Calcd for C₁₆H₈Br₄: C, 36.97; H, 1.55; Br, 61.48. Found: C, 37.22; H, 1.54; Br, 61.35.

1,2-Dibromo[2.2]paracyclophane-1-ene (5). Bromine (8.0 g, 2.6 mL, 50 mmol) was added dropwise to a refluxing solution of 10 g of fractions II (*R_f* = 0.22) and III (*R_f* = 0.22 and 0.09) of the column chromatographic isolation of 1,9(10)-dibromo[2.2]paracyclophane-1,9-diene⁵ consisting of a mixture of 1,9(10)-dibromo[2.2]paracyclophane-1,9-diene, 1-bromo[2.2]paracyclophane-1-ene, and 1,9(10)-dibromo[2.2]paracyclophane-1-ene in 150 mL of CHCl₃. The mixture was refluxed until the red color disappeared and then evaporated to dryness. The solid residue was washed with three portions (100 mL) of petroleum ether (30/40) [PE (30/40)], dried in vacuo, and suspended in 400 mL of dry TBME. KO^tBu (11.2 g, 100 mmol) was added, and the brown mixture was stirred for 6 h at room temperature and evaporated to dryness in vacuo. The residue was continuously extracted with chloroform for 5 days. From the extraction solution was collected by filtration 2.51 g (~35%) of precipitated **6**, and the solution was evaporated to dryness. The residue was recrystallized from heptane, yielding 3.5 g (~55%) of pure **5**, mp 196 °C. The spectroscopic data are identical with those reported.⁶

General Procedure (GP 1) for the Coupling of 5 with Alkenes. A mixture of 1.0 mmol of **5**, the specified amount of the alkene, 5.0 mmol of potassium carbonate (K₂CO₃), 2.0 mmol of tetrabutylammonium bromide (Bu₄NBr), and 25 mg (0.11 mmol) of palladium acetate Pd(OAc)₂ under nitrogen in 20 mL of dry dimethylformamide (DMF) was heated with vigorous stirring in a capped Pyrex bottle. At the end of the reaction, 100 mL of dichloromethane was added, the organic phase was washed with five portions of 50 mL of water, and the combined aqueous phases were extracted with 50 mL of dichloromethane. The combined organic phases were dried over magnesium sulfate (MgSO₄) and evaporated in vacuo. The crude products were chromatographed on silica gel. The obtained trienes were heated for 1 h in 50 mL of xylene under reflux; subsequently an equimolar amount of dichlorodicyano-*p*-benzoquinone (DDQ) or sulfur was added, and the mixture was further refluxed for 5 h. The organic phase was diluted with 80 mL of diethyl ether,

(25) Perrin, D. D.; Armarego, W. L. F.; Perrin, D. R. *Purification of Laboratory Chemicals*, 2nd ed.; Pergamon Press: New York, 1980.

(26) Hinkelmann, K.; Heinze, J.; Field, H.-T.; Vahrenkamp, H. *J. Am. Chem. Soc.* **1989**, *111*, 5078–5090.

washed twice with 100 mL of dilute NaOH, and dried over MgSO₄. The solvent was removed in vacuo and the residue chromatographed on silica gel.

4',5'-Dicarbomethoxy-1,2-benzo[2.2]paracyclophan-1-ene (12e). **5** (800 mg, 2.2 mmol) and 1.17 g (1.25 mL, 13.8 mmol) of methyl acrylate were reacted following GP 1 for 12 h at 50 °C. Column chromatography on 50 g of silica gel with ethyl acetate/PE (60/70) (1:10) as eluent (*R_f* = 0.85; ethyl acetate/dichloromethane 2:8) yielded 360 mg (44%) of dimethyl [2.2]paracyclophan-1-ene-(*E*)-1,2-diacrylate (**11e**), mp 166 °C. IR (KBr): ν 1710 (C=O) cm⁻¹. ¹H NMR (270 MHz, CDCl₃): δ 3.08 (s, 4 H, H-9,10), 3.80 (s, 6 H, CO₂CH₃), 6.20 (d, ³*J* = 15.6 Hz, H-2',2''), 6.47 (AB system, δ_A = 6.55, δ_B = 6.40, ³*J*_{AB} = 7.5 Hz, 8 H, H-arene of paracyclophane), 8.33 (d, ³*J* = 15.6 Hz, H-1',1''). ¹³C NMR (100.6 MHz, CDCl₃): δ 34.70 (-, C-9,10), 51.77 (+, CO₂CH₃), 125.46 (+, C-2',2''), 131.94 (+, C-arene of paracyclophane), 132.77 (+, C-arene of paracyclophane), 137.23 (C_{quat}), 138.38 (+, C-1',1''), 139.70 (C_{quat}), 148.00 (C_{quat}), 167.24 (C_{quat}, C=O). MS (70 eV) *m/z* (%): 290 (100, M⁺).

A solution of 747 mg (2.0 mmol) of **11e** in 40 mL of xylene was heated under reflux for 1 h; 500 mg (2.2 mmol) of DDQ was added, and the mixture was refluxed for 5 h. Chromatography on 50 g of silica gel with ethyl acetate/PE (60/70) 1:9 (*R_f* = 0.06) and ethyl acetate/dichloromethane 1:10 (*R_f* = 0.53) as eluents and recrystallization from heptane yielded 336 mg (45%) of **12e**, mp 208 °C. IR (KBr): ν 1720 (C=O) cm⁻¹. ¹H NMR (270 MHz, CDCl₃): δ 3.13 (s, 4 H, H-9,10), 3.98 (s, 6 H, CO₂CH₃), 6.55 (AB system, δ_A = 6.59, δ_B = 6.51, ³*J* = 7.2 Hz, 8 H, H-arene of paracyclophane), 7.94 (s, 2 H, H-arene) (100 MHz, CDCl₃): δ 34.60 (-, C-9,10), 52.75 (+, CO₂CH₃), 126.23 (+, C-3',6'), 130.55 (C_{quat}), 132.07 (+), 132.46 (+), 138.24 (C_{quat}), 139.97 (C_{quat}), 148.82 (C_{quat}), 167.99 (C_{quat}, C=O). MS (70 eV) *m/z* (%): 372 (100, M⁺). Anal. Calcd for C₂₄H₂₀O₄: C, 77.48; H, 5.38. Found: C, 77.41; H, 5.43.

4',5'-Diphenyl-1,2-benzo[2.2]paracyclophan-1-ene (12f). **5** (1.00 g, 2.75 mmol) and 2.5 mL (22 mmol) of styrene were reacted as described in GP 1 for 3 days at 100 °C. The crude product was chromatographed on 50 g of silica gel (dichloromethane/PE (60/70) 1:1, *R_f* = 0.7), yielding 654 mg (58%) of 1,2-distyryl[2.2]paracyclophan-1-ene (**11f**), mp 260 °C. ¹H NMR (270 MHz, CDCl₃): δ 3.15 (s, 4 H, H-9,10), 6.65 (AB system, δ_A = 6.60, δ_B = 6.70, ³*J*_{AB} = 8 Hz, 8 H, H-arene of paracyclophane), 6.88 (d, ³*J* = 15.6 Hz, 2 H), 7.32 (m, 10 H, H-phenyl), 7.90 (d, ³*J* = 15.6 Hz, 2 H). Anal. Calcd for C₃₂H₂₆: C, 93.66; H, 6.34. Found: C, 93.85; H, 6.36.

11f (650 mg, 1.58 mmol) was heated in 50 mL of xylene for 1 h at 140 °C. DDQ (430 mg, 1.9 mmol) was added, and the mixture was refluxed for 5 h. Chromatography on 50 g of silica gel (dichloromethane/hexane 1:4, *R_f* = 0.42) and recrystallization from heptane yielded 341 mg (53%) of **12f**, mp >280 °C. ¹H NMR (250 MHz, CDCl₃): δ 3.15 (s, 4 H, H-9,10), 6.65 (AB system, δ_A = 6.60, δ_B = 6.70, ³*J*_{AB} = 8 Hz, 8 H, H-arene of paracyclophane), 7.30 (m, 10 H, H-phenyl), 7.70 (s, 2 H, H-arene). ¹³C NMR (100 MHz, CDCl₃): δ 34.85 (-), 126.53, 127.95, 130.04, 132.30 and 132.43 (+), 138.89, 139.37, 139.87, 141.33 and 144.94 (C_{quat}). MS (70 eV) *m/z* (%): 408 (100, M⁺).

4',5'-Bis(*p*-carbomethoxyphenyl)-1,2-benzo[2.2]paracyclophan-1-ene (12i). **5** (2.0 g, 5.5 mmol) and 3.6 g (22 mmol) of methyl *p*-vinylbenzoate were reacted following GP 1 under sonification for 1 day at 70 °C. Chromatography on 50 g of silica gel (dichloromethane/PE (60/70) 2:1, *R_f* = 0.44) yielded 1.66 g (57%) of 1,2-bis(*p*-carbomethoxystyryl)[2.2]paracyclophan-1-ene (**11i**). ¹H NMR (270 MHz, CDCl₃): δ 3.07 (s, 4 H), 3.91 (s, 6 H, CO₂CH₃), 6.51 (AB system, δ_A = 6.55, δ_B = 6.46, ³*J*_{AB} = 8.8 Hz, 8 H, H-arene of paracyclophane), 6.85 (d, ³*J* = 15.4 Hz, 2 H), 7.78 (AB system, δ_A = 8.01, δ_B = 7.55, ³*J*_{AB} = 8.4 Hz, 8 H, H-arene), 7.93 (d, ³*J* = 15.4 Hz, 2 H). ¹³C NMR (68 MHz, CDCl₃): δ 34.90 (-), 51.89 (+, CO₂CH₃), 126.67 (+), 127.06 (+), 129.66 (C_{quat}), 130.09 (+), 132.57 (+), 132.69 (+), 134.96 (+), 139.25 (C_{quat}, relative intensity 2), 142.05 (C_{quat}), 145.42 (C_{quat}), 166.73 (C_{quat}). MS (70 eV) *m/z* (%): 526 (100, M⁺).

11i (1.61 g, 3.0 mmol) and 120 mg (3.7 mmol) of sulfur were refluxed in 50 mL of xylene for 12 h. The solvent was removed in vacuo, the solid residue was chromatographed on 50 g of silica gel (dichloromethane, *R_f* = 0.6), and the product was recrystallized from toluene/PE (60/70) 1:1, yielding 1.05 g (65%) of **12i**, mp 242 °C. IR (KBr): ν 1725 cm⁻¹ (C=O). ¹H NMR (270 MHz, CDCl₃): δ 3.11 (s, 4 H, H-9(10)), 3.89 (s, 6 H, CO₂CH₃), 6.59 (s, 8 H, H-arene of paracyclophane), 7.66 (AB system, δ_A = 7.94, δ_B = 7.39, ³*J*_{AB} = 8.4 Hz, 8 H), 7.68 (s, 2 H, H-arene). MS (70 eV) *m/z* (%): 524 (52, M⁺).

General Procedure for the Coupling of 6 with Alkenes (GP 2). **6** (1.0 mmol), 10.0 mmol of K₂CO₃, 4.0 mmol of Bu₄NBr, the specified amount of the alkene, and 25 mg (0.11 mmol) of Pd(OAc)₂ in 20 mL of anhydrous DMF under nitrogen were heated in a capped Pyrex bottle. At the end of the reaction the mixture was filtered, diluted with 100 mL of dichloromethane, and washed with five 50-mL portions of water. The aqueous phases were extracted with 50 mL of dichloromethane. The filtration residue was extracted with 200 mL of hot chloroform. All organic phases were combined, dried over MgSO₄, and evaporated to dryness in vacuo, and the crude products were chromatographed on silica gel. The obtained trienes were dissolved in 50 mL of xylene and refluxed for 1 h. After addition of equimolar amounts of DDQ or sulfur, the mixture was refluxed for 3 h, the solvent was removed in vacuo, and the crude products were purified by chromatography on silica gel.

1,2,9,10-Dibenzo[2.2]paracyclophane-1,9-diene (1a). (A) Following GP 2, 500 mg (0.96 mmol) of **6** was reacted under a pressure of 3 × 10⁵ Pa of ethylene for 2 days at 100 °C. The crude product was chromatographed on silica gel (dichloromethane/hexane 1:1, *R_f* = 0.9), yielding 70 mg (24%) of 1,2,9,10-tetravinyl[2.2]paracyclophane-1,9-diene (**9a**). ¹H NMR (270 MHz, CDCl₃): δ 5.45 (m, ³*J*_{cis} = 10 Hz, ³*J*_{trans} = 17 Hz, 8 H, H-olefin), 6.53 (s, 8 H, H-arene of paracyclophane), 7.30 (dd, ³*J*_{cis} = 10 Hz, ³*J*_{trans} = 17 Hz, 4 H, H-olefin).

According to GP 2, **9a** was cyclized and dehydrogenated with 15 mg (0.5 mmol) of sulfur, yielding after chromatography on 10 g of silica gel (hexane, *R_f* = 0.15) 54 mg of **1a**. Recrystallization from hexane afforded 40 mg (58%) of pure **1a**. All spectroscopic data are identical with those previously reported.^{8,9}

(B) **6** (1.00 g, 1.92 mmol) and 2.35 mL (14.7 mmol) of trimethylsilyl silane were reacted as described in GP 2 for 5 days at 40 °C. The crude product was chromatographed on 50 g of silica gel (dichloromethane/hexane, *R_f* = 0.9), and the so obtained material was oxidized with 55 mg (1.7 mmol) of sulfur following GP 2. Chromatography on 50 g of silica gel (hexane, *R_f* = 0.15) and recrystallization from hexane yielded 122 mg (21%) of **1a**. According to the ¹H NMR spectra of **1a**, more than 95% of the trimethylsilyl groups were cleaved. All other signals are identical with the reported ones.^{8,9}

[2.2]Paracyclophane-1,9-diene-1,2,9,10-tetraacrylaldehyde (9d). **6** (500 mg, 0.96 mmol) and 1.0 mL (15 mmol) of acrolein were reacted following GP 2 for 12 h under sonification at room temperature. Instead of K₂CO₃, 800 mg (9.6 mmol) of NaHCO₃ was used as base. The crude product was chromatographed on 50 g of silica gel (dichloromethane/ethyl acetate 3:1, *R_f* = 0.5), yielding 156 mg (38%) of **9d** as a yellow solid. IR (KBr): ν 1660 (C=O) cm⁻¹. ¹H NMR (270 MHz, CDCl₃): δ 6.45 (dd, ³*J* = 15.2 Hz, ³*J* = 7.8 Hz, 4 H, 2'-H), 6.62 (s, 8 H, H-arene of paracyclophane), 8.15 (d, ³*J* = 15.2 Hz, 4 H, 3'-H), 9.90 (d, ³*J* = 7.8 Hz, 4 H, CHO).

No products could be isolated after the dehydrogenation of **9d** with sulfur in xylene.

4',4'',5',5''-Tetracarbomethoxy-1,2,9,10-dibenzo[2.2]paracyclophane-1,9-diene (1e). **6** (1.00 g, 1.92 mmol) and 1.75 g (1.7 mL, 19.2 mmol) of methyl acrylate were reacted as described in GP 2 for 12 h at 60 °C under sonification. Chromatography on 50 g of silica gel (dichloromethane, *R_f* = 0.3) and recrystallization from dichloromethane/hexane (1:1) afforded 502 mg (48%) of tetramethyl [2.2]paracyclophane-1,9-diene-1,2,9,10-tetraacrylate (**9e**) as yellow crystals. IR (KBr): ν 1730 (C=O) cm⁻¹. ¹H NMR (270 MHz, CDCl₃): δ 3.79 (s, 12 H, CO₂CH₃), 6.16 (d, ³*J* = 15.2 Hz, 4 H, H-olefin), 6.54 (s, 8 H, H-arene of paracyclophane), 8.29 (d, ³*J* = 15.2 Hz, 4 H, H-olefin). ¹³C NMR (68 MHz, CDCl₃): δ 51.60 (+, CO₂CH₃), 126.32 (+, C-olefin), 131.58 (+, C-arene of paracyclophane), 137.62 (+, C-olefin), 138.58 (C_{quat}), 147.02 (C_{quat}, C-1(2)), 166.92 (C_{quat}, C=O). MS (70 eV) *m/z* (%): 540 (66, M⁺), 481 (100, M⁺ - CO₂CH₃), 422 (16, M⁺ - 2 CO₂CH₃), 363 (15, M⁺ - 3 CO₂CH₃), 304 (27, M⁺ - 4 CO₂CH₃). Anal. Calcd for C₃₂H₂₈O₈: 540.1784. Found 540.1807 (MS).

9e (500 mg, 0.92 mmol) and 80 mg (2.5 mmol) of sulfur were reacted following GP 2. The crude product was chromatographed on 50 g of silica gel (dichloromethane, *R_f* = 0.3); recrystallization from toluene/heptane (1:1) afforded 212 mg (43%) of **1e**. IR (KBr): ν 1720 (C=O) cm⁻¹. ¹H NMR (270 MHz, CDCl₃): δ 3.97 (s, 12 H, CO₂CH₃), 6.67 (s, 8 H, H-arene of paracyclophane), 8.00 (s, 4 H). ¹³C NMR (68 MHz, CDCl₃): δ 52.73 (+, CO₂CH₃), 126.22 (+), 131.13 (C_{quat}), 131.55 (+), 139.18 (C_{quat}), 147.59 (C_{quat}), 167.71 (C_{quat}, C=O). MS (70 eV) *m/z* (%): 536 (100, M⁺), 300 (13, M⁺ - 4 CO₂CH₃). Anal. Calcd for C₃₂H₂₄O₈: C, 71.60; H, 4.50. Found: C, 71.70; H, 4.40.

9,10-Dibromo-4',5'-dicarbomethoxy-1,2-benzo[2.2]paracyclophane-1,9-diene (10e). **6** (1.0 g, 1.92 mmol) and 1.75 g (1.7 mL, 19.2 mmol) of

methyl acrylate were reacted according to GP 2 at 60 °C under sonification for 2 h. Chromatography of the crude product mixture on 50 g of silica gel (dichloromethane) yielded 80 mg (8%) of **7d** (fraction I, $R_f = 0.7$). IR (KBr): ν 1720 (C=O) cm^{-1} . $^1\text{H NMR}$ (270 MHz, CDCl_3): δ 3.81 (s, 6 H, CO_2CH_3), 6.13 (d, $^3J = 15.4$ Hz, 2 H, H-olefin), 6.69 (AB system, $\delta_A = 6.78$, $\delta_B = 6.59$, $^3J = 8.4$ Hz, 8 H, H-arene of paracyclophane), 8.30 (d, $^3J = 15.4$ Hz, 2 H, H-olefin). $^{13}\text{C NMR}$ (68 MHz, CDCl_3): δ 51.81 (+, CO_2CH_3), 123.82 (C_{quat}), 126.54 (+), 131.05 (+), 131.34 (+), 137.65 (+), 139.55 (C_{quat}), 139.97 (C_{quat}), 146.45 (C_{quat}), 166.78 (C_{quat} , C=O). MS (70 eV) m/z (%): 526/528/530 (55/100/50, M^+), 449/447 (36/38, $\text{M}^+ - \text{Br}$), 368 (13, $\text{M}^+ - 2 \text{ Br}$). II ($R_f = 0.3$): 413 mg (40%) of **9e**.

4',4'',5',5''-Tetraphenyl-1,2,9,10-dibenzo[2.2]paracyclophane-1,9-diene (1f). **6** (1.50 g, 2.9 mmol) was reacted with 5.0 mL (43.6 mmol) of styrene for 3 days at 100 °C following GP 2. The crude product was chromatographed on 50 g of silica gel (dichloromethane/hexane 1:1, $R_f = 0.76$), yielding 971 mg (55%) of 1,2,9,10-tetrastryl[2.2]paracyclophane-1,9-diene (**9f**) as a yellow solid. $^1\text{H NMR}$ (270 MHz, CDCl_3): δ 6.68 (s, 8 H, H-arene of paracyclophane), 6.87 (d, $^3J = 15.6$ Hz, 4 H), 7.31–7.58 (m, 20 H, H-phenyl), 7.91 (d, $^3J = 15.6$ Hz, 4 H). MS (70 eV) m/z (%): 612 (100, M^+).

9f was dehydrogenated with 105 mg (3.3 mmol) of sulfur, as described in GP 2. Chromatography on 50 g of silica gel (dichloromethane/hexane 1:4) and recrystallization from toluene/hexane (2:1) afforded 482 mg (28%) of **1f**, mp >280 °C. $^1\text{H NMR}$ (270 MHz, CDCl_3): δ 6.79 (s, 8 H, H-arene of paracyclophane), 7.24–7.28 (m, 20 H, H-phenyl), 7.74 (s, 4 H). $^{13}\text{C NMR}$ (68 MHz, CDCl_3): δ 122.66, 127.91, 128.00, 130.12 and 131.76 (+), 139.42, 140.29, 141.51 and 144.22 (C_{quat}). MS (70 eV) m/z (%): 608 (100, M^+). Anal. Calcd for $\text{C}_{48}\text{H}_{32}$: 608.2504. Found: 608.2504 (MS).

9,10-Dibromo-4',5'-diphenyl-1,2-benzo[2.2]paracyclophane-1,9-diene (10f). **6** (1.0 g, 1.9 mmol) and 3.5 mL (30.5 mmol) of styrene were reacted according to GP 2 at 60 °C for 1 day. The crude product was chromatographed on 50 g of silica gel (dichloromethane/hexane 1:1, $R_f = 0.76$), yielding 610 mg of **7e**. Dehydrogenation with 65 mg (2.0 mmol) of sulfur, chromatography on 25 g of silica gel (dichloromethane/hexane 1:4, $R_f = 0.37$), and recrystallization from toluene/hexane (2:1) afforded 392 mg (36%) of 9,10-dibromo-4',5'-diphenyl-1,2-benzo[2.2]paracyclophane-1,9-diene. $^1\text{H NMR}$ (270 MHz, CDCl_3): δ 6.76 (b s, 8 H, H-arene of paracyclophane), 7.26 (b s, 10 H, H-phenyl), 7.70 (s, 2 H). $^{13}\text{C NMR}$ (68 MHz, CDCl_3): δ 124.05 (C_{quat}), 126.81 (+), 127.69 (+), 128.05 (+), 130.04 (+), 132.06 (+), 139.36 (C_{quat}), 139.79 (C_{quat}), 141.22 (C_{quat}), 141.66 (C_{quat}), 142.83 (C_{quat}). MS (70 eV) m/z (%): 562/564/566 (51/100/53, M^+). Anal. Calcd for $\text{C}_{32}\text{H}_{20}\text{Br}_2$: C, 68.11; H, 3.57; Br, 28.32. Found: C, 68.14; H, 3.60; Br, 28.29.

4',4'',5',5''-Tetrakis(*p*-fluorophenyl)-1,2,9,10-dibenzo[2.2]paracyclophane-1,9-diene (1g). **6** (300 mg, 0.58 mmol) and 550 mg (4.5 mmol) of *p*-fluorostyrene were reacted for 12 h at 100 °C according to GP 2. Chromatography on 50 g of silica gel (dichloromethane/hexane 1:2, $R_f = 0.4$) yielded 176 mg (45%) of **9g**. Without further purification **9g** was dehydrogenated with 80 mg (2.5 mmol) of sulfur following GP 2. The crude product was chromatographed on 50 g of silica gel (dichloromethane/hexane 1:2, $R_f = 0.4$), yielding 80 mg (20%) of **1g**, mp >290 °C. IR (KBr): ν 2950, 1600, 1500, 820 cm^{-1} . $^1\text{H NMR}$ (270 MHz, CDCl_3): δ 6.77 (b s, 8 H, H-arene of paracyclophane), 6.97 (m, 16 H, H-phenyl), 7.72 (b s, 4 H, H-arene). MS (70 eV) m/z (%): 680 (100, M^+). Anal. Calcd for $\text{C}_{48}\text{H}_{28}\text{F}_4$: 680.2127. Found 680.2097 (MS).

4',4'',5',5''-Tetrakis(*p*-tert-butylphenyl)-1,2,9,10-dibenzo[2.2]paracyclophane-1,9-diene (1h). **6** (1.00 g, 1.92 mmol) and 2.4 g (2.7 mL, 15 mmol) of *p*-tert-butylstyrene were reacted at 100 °C for 20 h according to GP 2. Chromatography on 50 g of silica gel (dichloromethane/hexane 1:4, $R_f = 0.29$) yielded 880 mg (55%) of 1,2,9,10-tetrakis(*E*)-*p*-tert-butylstyryl[2.2]paracyclophane-1,9-diene (**9h**) as a yellow solid. IR (KBr): ν 3030, 2960, 2900, 1450 cm^{-1} . $^1\text{H NMR}$ (270 MHz, CDCl_3): δ 1.33 [s, 36 H, $\text{C}(\text{CH}_3)_3$], 6.66 (s, 8 H, H-arene of paracyclophane), 6.84 (d, $^3J = 15.6$ Hz, 4 H, H-olefin), 7.41 (AB system, $\delta_A = 7.46$, $\delta_B = 7.35$, $^3J = 8.4$ Hz, 16 H, H-phenyl), 7.87 (d, $^3J = 15.6$ Hz, 4 H, H-olefin).

9h (870 mg, 1.04 mmol) was dehydrogenated with 80 mg (2.5 mmol) of sulfur according to GP 2. Chromatography of the crude product on 50 g of silica gel (dichloromethane/hexane 1:4, $R_f = 0.32$) and recrystallization from toluene/ethanol (1:1) afforded 390 mg (25%) of **1h**, mp >290 °C. IR (KBr): ν 3050, 2950, 2900, 1450, 1110 cm^{-1} . $^1\text{H NMR}$ (270 MHz, CDCl_3): δ 1.30 [s, 36 H, $\text{C}(\text{CH}_3)_3$], 6.77 (s, 8 H, H-arene of paracyclophane), 7.24 (AB system, $\delta_A = 7.27$, $\delta_B = 7.21$, $^3J = 8.8$ Hz, 16 H, H-phenyl), 7.72 (s, 4 H, H-arene). $^{13}\text{C NMR}$ (68 MHz, CDCl_3): δ 31.46 (+, $\text{C}(\text{CH}_3)_3$), 34.54 (C_{quat} , $\text{C}(\text{CH}_3)_3$), 124.78 (+,

127.83 (+), 129.76 (+), 131.75 (+), 138.61 (C_{quat}), 139.35 (C_{quat}), 140.42 (C_{quat}), 143.94 (C_{quat}), 149.62 (C_{quat}). MS (70 eV) m/z (%): 832 (100, M^+). Anal. Calcd for $\text{C}_{64}\text{H}_{64}$: C, 92.26; H, 7.74. Found: C, 92.12; H, 7.70.

4',4'',5',5''-Tetrakis(*p*-carbomethoxyphenyl)-1,2,9,10-dibenzo[2.2]paracyclophane-1,9-diene (1i). A mixture of 1.50 g (2.88 mmol) of **6**, 3.73 g (23.0 mmol) of methyl *p*-vinylbenzoate, 3.80 g (11.7 mmol) of Bu_4NBr , 4.10 g (29.6 mmol) of K_2CO_3 , and 25 mg (11.05 mmol) of $\text{Pd}(\text{OAc})_2$ in 40 mL of dry DMF under nitrogen was heated at 70 °C for 36 h under sonification. $\text{Pd}(\text{OAc})_2$ (25 mg) was added to the reaction mixture every 12 h. After the end of the reaction, the mixture was diluted with 500 mL of dichloromethane and washed with four portions of 150 mL of water. The organic phase was concentrated to a volume of 50 mL, and 50 mL of PE (60/70) was added. The yellow precipitate (1.3 g of **9i**) was removed by filtration, and the filtrate was evaporated to dryness. The residue was chromatographed on 50 g of silica gel (dichloromethane, $R_f = 0.5$) to yield 210 mg of **9i**. Total yield: 1.51 g (58%) of **9i**. IR (KBr): ν 1720 (C=O) cm^{-1} . $^1\text{H NMR}$ (270 MHz, CDCl_3): δ 3.92 (s, 12 H, CO_2CH_3), 6.69 (s, 8 H, H-arene of paracyclophane), 6.91 (d, $^3J = 15.7$ Hz, 4 H, H-olefin), 7.82 (AB system, $\delta_A = 8.04$, $\delta_B = 7.59$, $^3J = 8.4$ Hz, 16 H, H-arene), 8.00 (d, $^3J = 15.7$ Hz, 4 H, H-olefin). $^{13}\text{C NMR}$ (68 MHz, CDCl_3): δ 51.95 (+, CO_2CH_3), 126.74 (+, relative intensity = 2), 129.69 (C_{quat}), 130.17 (+), 132.01 (+), 135.14 (+), 140.09 (C_{quat}), 141.92 (C_{quat}), 144.66 (C_{quat}), 166.72 (C_{quat} , C=O). Anal. Calcd for $\text{C}_{56}\text{H}_{44}\text{O}_8$: C, 79.60; H, 5.25. Found: C, 79.80; H, 5.34.

A solution of 1.45 g (1.72 mmol) of **9i** and 200 mg (6.25 mmol) of sulfur in 150 mL of xylene was refluxed for 48 h. After 12 h at 4 °C the white precipitate (743 mg of **1i**) was removed by filtration, the filtrate was evaporated to dryness, and the residue was chromatographed on 50 g of silica gel with dichloromethane ($R_f = 0.9$) and dichloromethane/ethyl acetate (4:1) as eluents. Recrystallization of the crude product from toluene yielded 420 mg of **1i**. Total yield: 1.16 g (80%) of **1i**, mp >290 °C. IR (KBr): ν 1720 (C=O) cm^{-1} . $^1\text{H NMR}$ (270 MHz, CDCl_3): δ 3.90 (s, 12 H, CO_2CH_3), 6.79 (s, 8 H, H-arene of paracyclophane), 7.67 (AB system, $\delta_A = 8.00$, $\delta_B = 7.34$, $^3J = 8.0$ Hz, 16 H, H-arene), 7.76 (s, 4 H). $^{13}\text{C NMR}$ (68 MHz, CDCl_3): δ 51.94 (+, CO_2CH_3), 127.83 (+), 129.02 (C_{quat}), 129.54 (+), 130.05 (+), 131.79 (+), 138.58 (C_{quat}), 139.96 (C_{quat}), 144.93 (C_{quat}), 145.68 (C_{quat}), 166.62 (C_{quat} , C=O). MS (70 eV) m/z (%): 840 (32, M^+), 91 (100, C_7H_7^+). Anal. Calcd for $\text{C}_{56}\text{H}_{40}\text{O}_8$: C, 79.98; H, 4.79. Found: C, 80.05; H, 4.93.

4',4'',5',5''-Tetra(biphenyl)-1,2,9,10-dibenzo[2.2]paracyclophane-1,9-diene (1j). **6** (960 mg, 1.7 mmol), 2.40 g (13.3 mmol) of 4-vinylbiphenyl, 2.36 g (17.1 mmol) of K_2CO_3 , 2.2 g (6.8 mmol) of Bu_4NBr , and 43 mg (0.19 mmol) of $\text{Pd}(\text{OAc})_2$ in 35 mL of anhydrous DMF was heated under nitrogen for 4 days at 100 °C. The reaction mixture was diluted with 200 mL of dichloromethane, filtered, washed with five portions of 100 mL of water, dried over MgSO_4 , and evaporated in vacuo. Chromatography on 80 g of silica gel (dichloromethane/PE (60/70) 1:1, $R_f = 0.2$) yielded 233 mg (15%) of 1,2,9,10-tetrakis(*p*-phenylstyryl)[2.2]paracyclophane-1,9-diene (**9j**) as a yellow solid. IR (KBr): ν 3026, 1600, 1486, 761 cm^{-1} . $^1\text{H NMR}$ (250 MHz, CDCl_3): δ 6.75 (s, 8 H, H-arene of paracyclophane), 6.95 (d, $^3J = 15.4$ Hz, 4 H), 7.25–7.70 (m, 36 H), 8.05 (d, $^3J = 15.4$ Hz, 4 H).

9j (220 mg, 0.24 mmol) in 100 mL of xylene was refluxed for a total of 7 h; after 1 h 210 mg (0.93 mmol) of DDQ was added. Chromatography on 80 g of silica gel (dichloromethane, $R_f = 0.5$) yielded 68 mg (31%) of **1j**, mp >280 °C. IR (KBr): ν 3027, 1486, 765, 697 cm^{-1} . $^1\text{H NMR}$ (250 MHz, CDCl_3): δ 6.84 (s, 8 H), 7.35, 7.45 and 7.60 (m, 36 H), 7.84 (s, 4 H). MS (70 eV) m/z (%): 912 (8, M^+), 359 (82), 180 (100).

4',5'-Bis(bromomethyl)-1,2-benzo[2.2]paracyclophane-1-ene (13). A solution of 320 mg (0.86 mmol) of **12e** in 20 mL of dry THF was added dropwise to a suspension of 100 mg (2.6 mmol) of LAH in 40 mL of THF at 0 °C. The mixture was allowed to warm up to room temperature, stirred for 1 h, and was hydrolyzed with 10 g of ice. The solvent was removed in vacuo and the white residue purified by flash chromatography on 30 g of silica gel (ethyl acetate/dichloromethane 1:10, $R_f = 0.01$), yielding 230 mg (85%) of 4',5'-bis(hydroxymethyl)-1,2-benzo[2.2]paracyclophane-1-ene, mp 221 °C. IR (KBr): ν 3290 (OH) cm^{-1} . $^1\text{H NMR}$ (270 MHz, CDCl_3): δ 2.89 (s, 2 H, OH), 3.11 (s, 4 H, H-9,10), 4.91 (s, 4 H, H-benzyl), 6.53 (AB system, 8 H, H-arene of paracyclophane), 7.62 (s, 2 H). $^{13}\text{C NMR}$ (68 MHz, CDCl_3): δ 35.68 (-), 62.96 (-), 126.79 (+), 133.38 (+), 133.49 (+), 138.85 (C_{quat}), 140.63 (C_{quat}), 141.52 (C_{quat}). MS (70 eV) m/z (%): 316 (79, M^+).

To a suspension of 210 mg (0.66 mmol) of 4',5'-bis(hydroxymethyl)-1,2-benzo[2.2]paracyclophane-1-ene in 20 mL of dry benzene was added

at room temperature 450 mg (1.7 mmol, 1% solution in benzene) of phosphorus tribromide, and the mixture was stirred for 12 h. The reaction mixture was poured onto 200 g of ice, diluted with 150 mL of diethyl ether, and extracted with four portions of 50 mL of water. The organic phase was dried over MgSO_4 and the solvent evaporated in vacuo. Chromatography on 30 g of silica gel (ethyl acetate/dichloromethane 1:10, $R_f = 0.69$) yielded 130 mg (44%) of **13**, mp 252 °C. IR (KBr): ν 3070, 1600, 1500 cm^{-1} . $^1\text{H NMR}$ (270 MHz, CDCl_3): δ 3.12 (s, 4 H, H-9,10), 4.80 (s, 4 H, H-benzyl), 6.55 (AB system, $\delta_A = 6.58$, $\delta_B = 6.51$, $^3J = 8$ Hz, 8 H, H-arene of paracyclophane), 7.60 (s, 2 H). $^{13}\text{C NMR}$ (100 MHz, CDCl_3): δ 30.10 (-), 34.63 (-), 128.44 (+), 132.22 (+), 132.36 (+), 135.03 (C_{quat}), 138.69 (C_{quat}), 139.50 (C_{quat}), 146.96 (C_{quat}). MS (70 eV) m/z (%): 444/442/440 (26/51/27, M^+), 283 (100).

5',6'-Dicarboxy-4',5',6',7'-tetrahydro-1,2-naphtho[2.2]paracyclophane-1-ene (15). A mixture of 80 mg (0.18 mmol) of **13**, 249 mg (0.23 mL, 1.45 mmol) of diethyl maleate, and 980 mg (15 mmol) of activated zinc²⁷ in 8 mL of dry dioxane was sonicated at room temperature for 16 h. The reaction mixture was filtered, diluted with 250 mL of dichloromethane, washed with 100 mL of saturated NH_4Cl and 100 mL of water, dried over MgSO_4 , and evaporated in vacuo. The residue was recrystallized from toluene/heptane (1:3), yielding 37 mg (45%) of **15**, mp 175 °C. IR (KBr): ν 1730 ($\text{C}=\text{O}$) cm^{-1} . $^1\text{H NMR}$ (400 MHz, C_6D_6): δ 1.02 (t, 6 H, CH_3), 2.79 (s, 4 H, H-9,10), 2.90 (m, 2 H), 3.08 (m, 2 H), 3.21 (m, 2 H), 4.02 (q, 4 H, OCH_2), 6.45 (AB system, $\delta_A = 6.55$, $\delta_B = 6.35$, $^3J_{\text{AB}} = 7.5$ Hz, 8 H, H-arene of paracyclophane), 7.23 (s, 2 H). $^{13}\text{C NMR}$ (100 MHz, CDCl_3): δ 14.21 (+, CH_3), 31.66 (-, OCH_2), 34.62 (-, C-9(10)), 42.31 (+), 60.63 (-), 125.61 (+), 132.19 (+), 132.33 (+), 132.37 (C_{quat}), 139.25 (C_{quat}), 139.68 (C_{quat}), 143.64 (C_{quat}), 174.40 (C_{quat} , $\text{C}=\text{O}$). MS (70 eV) m/z (%): 454 (100, M^+). Anal. Calcd for $\text{C}_{30}\text{H}_{30}\text{O}_4$: C, 79.31; H, 6.60. Found: C, 78.71; H, 6.64.

7',7'',10',10''-Tetra-tert-butyl-1,2:2',3';9,10:2'',3''-bis(triphenyleno)-[2.2]paracyclophane-1,9-diene (16). A solution of 30 mg (0.04 mmol) of **1h** and 20 mg (0.08 mmol) of iodine in 20 mL of benzene was irradiated in a quartz tube under nitrogen with a 250-W Hg medium-pressure lamp for 24 h. The white precipitate was removed by filtration, washed with 10 mL of hexane, and dried in vacuo, to yield 20 mg (66%) of **16**, mp >290 °C. $^1\text{H NMR}$ (250 MHz, CDCl_3): δ 1.39 [s, 36 H, $\text{C}(\text{CH}_3)_3$], 6.67 (s, 8 H, H-arene of paracyclophane), 8.24 (ABX system, $\delta_A = 8.71$, $\delta_B = 8.68$, $\delta_X = 7.77$, $^3J_{\text{BX}} = 8.0$ Hz, 12 H, H-arene), 8.91 (s, 4 H).

4',5'-Bis(*p*-bromomethyl)phenyl)-1,2-benzo[2.2]paracyclophane-1-ene (17). A solution of 143 mg (0.27 mmol) of **12i** in 100 mL of dry THF was added dropwise to a suspension of 40 mg (1.1 mmol) of LAH in 80 mL of THF at 0 °C. The reaction mixture was allowed to warm to room temperature and stirred for 1 h. After hydrolysis with 20 g of ice, the mixture was extracted with 100 mL of chloroform, the organic phase was dried over MgSO_4 and evaporated to dryness, and the solid residue was subjected to chromatography on 50 g of silica gel (dichloromethane/ethyl acetate 10:1, $R_f = 0.1$), yielding 75 mg (60%) of 4',5'-bis(*p*-

(hydroxymethyl)phenyl)-1,2-benzo[2.2]paracyclophane-1-ene, mp >290 °C. $^1\text{H NMR}$ (270 MHz, CDCl_3): δ 1.60 (b s, 2 H, OH), 3.10 (s, 4 H, H-9,10), 4.62 (b s, 4 H, CH_2O), 6.58 (AB system, $\delta_A = 6.62$, $\delta_B = 6.56$, $^3J_{\text{AB}} = 8.0$ Hz, H-arene of paracyclophane), 7.35 (AB system, $\delta_A = 7.38$, $\delta_B = 7.30$, $^3J_{\text{AB}} = 8.0$ Hz, H-phenyl), 7.60 (s, 2 H, H-3',6').

To a suspension of 70 mg (0.15 mmol) of 4',5'-bis(*p*-(hydroxymethyl)phenyl)-1,2-benzo[2.2]paracyclophane-1-ene in 10 mL of benzene was added at 0 °C 2 mL (0.2 mmol, 0.1 M solution in benzene) of phosphorus tribromide. The mixture was allowed to warm up to room temperature and stirred for 12 h. Water (0.5 mL) was added, the solution was dried over MgSO_4 , and the solvent was evaporated to dryness. The solid residue was chromatographed on 25 g of silica gel (dichloromethane/hexane 1:1, $R_f = 0.39$), yielding 73 mg (82%) of **17**, mp >290 °C. $^1\text{H NMR}$ (270 MHz, CDCl_3): δ 3.12 (s, 4 H, H-9,10), 4.49 (s, 4 H, CH_2Br), 6.60 (AB system, $\delta_A = 6.62$, $\delta_B = 6.57$, $^3J_{\text{AB}} = 8.4$ Hz, 8 H, H-arene of paracyclophane), 7.27 (AB system, $\delta_A = 7.30$, $\delta_B = 7.23$, $^3J_{\text{AB}} = 8.4$ Hz, 8 H, H-phenyl), 7.65 (s, 2 H, H-3',6'). $^{13}\text{C NMR}$ (68 MHz, CDCl_3): δ 33.26 (-), 34.96 (-), 127.96 (+), 128.79 (+), 130.39 (+), 132.36 (+), 132.45 (+), 136.24 (C_{quat}), 138.27 (C_{quat}), 139.45 (C_{quat}), 139.71 (C_{quat}), 141.47 (C_{quat}), 145.39 (C_{quat}). MS (70 eV) m/z (%): 596/594/592 (44/100/49, M^+). Anal. Calcd for $\text{C}_{34}\text{H}_{26}\text{Br}_2$: C, 68.70; H, 4.41; Br, 26.89. Found: C, 68.33; H, 4.38; Br, 26.48.

1,2:4,5-Bis(2.2)paracyclophane-1-eno)benzene (18). A suspension of 100 mg (0.17 mmol) of **17** in 100 mL of diethyl ether was added dropwise over 1 h at room temperature to a solution of 1.2 mmol of phenyllithium in 20 mL of diethyl ether. Water (1 mL) was added, the solvent was evaporated in vacuo, and the solid residue was chromatographed on 25 g of silica gel (dichloromethane/hexane 1:1). Fraction I ($R_f = 0.95$): biphenyl, not isolated. II ($R_f = 0.80$ – 0.85): 22 mg of an unidentified product mixture. III ($R_f = 0.58$ – 0.68): 24 mg of three compounds. Separation by preparative thin-layer chromatography (dichloromethane/hexane 1:1) afforded 8 mg (11%) ($R_f = 0.68$) of **18**. $^1\text{H NMR}$ (270 MHz, CDCl_3): δ 3.13 (s, 8 H), 6.64 (AB system, $\delta_A = 6.69$, $\delta_B = 6.59$, $^3J_{\text{AB}} = 8.0$ Hz, 16 H, H-arene of paracyclophane), 7.79 (s, 2 H, H-arene). MS (70 eV) m/z (%): 434 (100, M^+). Anal. Calcd for $\text{C}_{34}\text{H}_{26}$: 434.2035. Found: 434.2032 (MS).

Acknowledgment. This work was supported by the Volkswagen-Stiftung, the Deutsche Forschungsgemeinschaft, the Schweizerischer Nationalfonds zur Förderung der wissenschaftlichen Forschung, the Fonds der Chemischen Industrie, BASF AG, Hoechst AG, Bayer AG, and Degussa AG. O.R. and B.K. thank the Studienstiftung des Deutschen Volkes for a doctoral fellowship, O.R. thanks the Minerva foundation for a young scientists' travel award. We thank Dr. B. Knieriem for his careful revision of the manuscript.