

A Ready, Convergent Synthesis of the Heptasaccharide GPI Membrane Anchor of Rat Brain Thy-1 Glycoprotein¹

Uko E. Udodong,^{2a} Robert Madsen,^{2b}
Carmichael Roberts,^{2c} and Bert Fraser-Reid*

Paul M. Gross Chemical Laboratory
Department of Chemistry, Duke University
Durham, North Carolina 27708


Received April 12, 1993

That certain proteins are covalently attached to cell surfaces via glycosylphosphatidylinositol (GPI) anchors was first established eight years ago by Ferguson and co-workers for the variant surface glycoprotein (VSG) of the parasitic protozoan *Trypanosoma brucei*.³ Since then over 100 GPI-anchored proteins have been identified in eukaryotes⁴ with diverse functions, including adhesion molecules, hydrolases, receptors, and transmembrane signal inducers.^{4,5} Common structural features in the complex oligosaccharide anchoring domain, indicative of a high degree of evolutionary conservation, would seem to suggest functional interrelationships.⁶ However, Ferguson has cautioned that "x does y rules' may be hard to realize".⁴ Furthermore, a structure/activity debate may be premature since complete structural details⁷ are available for only two GPI anchors, VSG³ and the rat brain Thy-1 glycoprotein,⁸ the latter being of special interest since it was the first to be isolated from a mammalian source. In this communication we describe the first synthesis of the complete glycan moiety of Thy-1, by a route which is convergent, provides the material in multimilligram amounts, and makes provisions for attaching the protein components.


Although the structures of the Thy-1 (1) and VSG anchors (Scheme I) may seem similar biologically, their laboratory syntheses^{9,10} require vastly different strategies. We envisaged a triply convergent approach based on subunits 2 and 3 in the form of *n*-pentenyl glycosides (NPGs)¹¹ and the pseudodisaccharide 4.

The retrosynthetic plan in Scheme I gives subunits 2, 3, and 4. The trisaccharide 2 presented an opportunity to explore the advantages of the pentenyl activating group for reducing the numbers of "starting materials" required for oligosaccharide syntheses. As described in a recent report from this laboratory, the ability to readily protect and deprotect pentenyl double bonds enabled the preparation of a nonamannan from only two starting materials.¹² In this communication we focus attention on the divergent pathways followed by pentenyl ortho esters upon

Scheme I



In VSG, M4 is missing and M1 has a tetraacetate at O3 instead of O4 GalNAc residue.



treatment with different electrophiles (Scheme II). For example, reaction of 5a¹¹ with a protic acid induces transfer of the pentenyloxy group to the anomeric center of 10a.¹³ However, when titrated with bromine, pentenyl ortho esters yield glycosyl bromides such as 9.¹¹ These reaction pathways can be rationalized by invoking the protonated intermediate 7 and the furanylium ion 6, both of which proceed to the dioxolenium ion 8, which then reacts with the best nucleophile available to give 9 or 10. Coupling of the former is best carried out *in situ* after discharging excess bromine with a stream of argon. In this way, 9b and 10b furnished disaccharide 11a in 89% yield and, in turn, acceptor 11b reacted with 9a to give trisaccharide 12a. The protecting groups of the latter were then changed to those in 12b to provide for future phosphorylations.

With respect to the disaccharide segment 3 (Scheme I), the galactosyl donor is disarmed by virtue of the phthalimide residue, and hence the armed/disarmed strategy employing two NPGs¹¹ would be problematic. Hence a different glycosyl donor, the trichloroacetimidate,¹⁴ was chosen since it had been tested in a comparable coupling.¹⁵ Accordingly, compound 13¹⁶ was coupled with 14, obtained from *n*-pentenyl mannopyranoside¹¹ by standard transformations,¹⁷ to give disaccharide 15a (Scheme III). Cleavage of the phthalimide¹⁸ and selective acylation then led to 15b.

Synthetic routes to variously protected forms of the NI pseudodisaccharide 4 (Scheme I) have emerged from this⁹ and other^{10,19} laboratories, and so there were several options that could be retooled to fit our present needs. A critical factor in our choice was the desire to protect the future site of inositol phosphorylation with a functionality that would be refractory to the conditions of NPG activation¹¹ (*vide infra*). An allyl group fit this need, and hence the known²⁰ derivative 17 was chosen for coupling

(1) This work was supported by grants from the NIH (GM 41071) and NSF (CHE 8920003).

(2) (a) Present address for U.E.U.: Eli Lilly Corporation, Indianapolis, IN; (b) R.M. thanks the Danish Technical Research Council for a P.D. Fellowship; (c) C.R. is the recipient of an NIH Award for Underrepresented Minorities.

(3) Ferguson, M. A. J.; Low, M. G.; Cross, G. A. M. *J. Biol. Chem.* **1985**, *260*, 14547. Ferguson, M. A. J.; Homans, S. W.; Dwek, R. A.; Rademacher, T. W. *Science* **1988**, *239*, 753.

(4) Ferguson, M. A. J. *Curr. Opin. Struct. Biol.* **1991**, *1*, 522.

(5) Robinson, P. J. *Immunol. Today* **1991**, *12*, 35.

(6) Ferguson, M. A. J.; Williams, A. F. *Annu. Rev. Biochem.* **1988**, *57*, 2985. Low, M. G. *Biochim. Biophys. Acta* **1989**, *988*, 427.

(7) For reviews, see: Cross, G. A. M. *Annu. Rev. Cell. Biol.* **1990**, *6*, 1. Thomas, J. R.; Dwek, R. A.; Rademacher, T. W. *Biochemistry* **1990**, *29*, 5413.

(8) (a) Homans, S. W.; Ferguson, M. A. J.; Dwek, R. A.; Rademacher, T. W.; Anand, R.; Williams, A. F. *Nature* **1988**, *333*, 269. (b) For a picturesque model of the Thy-1 glycoprotein, see *Chem. Eng. News* **1993**, *61* (Jan 11), 26.

(9) Mootoo, D. R.; Konradsson, P.; Fraser-Reid, B. *J. Am. Chem. Soc.* **1989**, *111*, 8540.

(10) Murakata, C.; Ogawa, T. *Carbohydr. Res.* **1992**, *235*, 95.

(11) Fraser-Reid, B.; Udodong, U. E.; Wu, Z.; Ottosson, H.; Merritt, J. R.; Rao, C. S.; Roberts, C.; Madsen, R. *Synlett* **1992**, 927.

(12) Merritt, J. R.; Fraser-Reid, B. *J. Am. Chem. Soc.* **1992**, *114*, 8334.

(13) For proton-induced rearrangements β -D-mannopyranose 1,2-(alkyl) ortho esters, see: (a) Franks, N. E.; Montgomery, R. *Carbohydr. Res.* **1968**, *6*, 286. (b) Ogawa, T.; Nukada, T. *Carbohydr. Res.* **1988**, *136*, 135.

(14) Schmidt, R. R. *Angew. Chem., Int. Ed. Engl.* **1986**, *25*, 212. Schmidt, R. R. *Pure Appl. Chem.* **1989**, *61*, 1257.

(15) Ito, Y.; Nunomura, S.; Shibayama, S.; Ogawa, T. *J. Org. Chem.* **1992**, *57*, 1821.



(16) For the tetraacetate related to 13, see: Paulsen, H.; Bünsch, A. *Carbohydr. Res.* **1982**, *100*, 143. Five standard steps involving glycosidation with *p*-methoxyphenol, deacetylation, benzylation, oxidation with CAN, and reaction with Cl₃CCN afforded 13.

(17) Patroni, J. J.; Stick, R. V.; Skelton, B. W.; White, A. H. *Aust. J. Chem.* **1988**, *41*, 91. Nashed, M. A. *Carbohydr. Res.* **1978**, *60*, 200.

(18) Wolfe, S.; Hasan, S. K. *Can. J. Chem.* **1970**, *48*, 3572.

(19) See for examples: Zapata, A.; Martin-Lomas, M. *Carbohydr. Res.* **1992**, *234*, 93. Plourde, R.; d'Alarco, M.; Saltiel, A. R. *J. Org. Chem.* **1992**, *57*, 2606. Verduyn, R.; Elie, C. J. J.; Dreef, C. E.; van der Marel, G. A.; van Boom, J. H. *Recl. Trav. Chim. Pays-Bas* **1990**, *109*, 591.

(20) Elie, C. J. J.; Verduyn, R.; Dreef, C. E.; Broutns, D. M.; van der Marel, G. A.; van Boom, J. H. *Tetrahedron* **1990**, *46*, 8243.

Scheme II^aScheme III^a

with the glycosyl bromide **16**,²¹ the product **18a** being obtained in 63% yield.

Assembly of the heptasaccharide was now undertaken. Coupling of acceptor **18b** and donor **15b** in 1.2:1 ratio²² was complete within 15 min to give tetrasaccharide **19a** in 55% yield based on **15b**, substantial amounts of acceptor **18b** also being recovered.²³ Dechloroacetylation then yielded **19b** for coupling to trisaccharide **12b**. For this task, compounds **12b** (250 mg) and

19b (200 mg) were used in 1.5:1 ratio, affording 115 mg of heptasaccharide **20** (39% yield based on recovered acceptor **19b**).

The sites for future couplings in **20** have been provided with temporary protecting groups that can be removed selectively. These tasks are now being carried out as a prelude to biological testing.

Acknowledgment. We are grateful to Dr. M. A. J. Ferguson of the University of Dundee, Scotland, for his advice and interest.

Supplementary Material Available: Listings of experimental procedures for the preparation of all key compounds and their NMR data (16 pages). Ordering information is given on any current masthead page.

(21) Paulsen, H.; Richter, A.; Sinnwell, V.; Stenzel, W. *Carbohydr. Res.* **1978**, *64*, 339. Hori, H.; Nishida, Y.; Ohri, H.; Meguro, H. *J. Org. Chem.* **1989**, *54*, 1346.

(22) The unusual use of the acceptor **18b** in excess was due to the observation that the material was highly prone to silylation by the NIS/Et₃SiOTf promoter.

(23) The recovered silylated glycosyl acceptor was treated with tetra-*n*-butylammonium fluoride in THF to regenerate the corresponding alcohol.