

Nonplanarity of Tetrafluorocyclobutadiene

E. James Petersson,^{†,‡} Jason C. Fanuele,[†] Mark R. Nimlos,[§]
David M. Lemal,[†] G. Barney Ellison,[‡] and
J. George Radziszewski^{*,§}


National Renewable Energy Laboratory (NREL)
1617 Cole Blvd., Golden, Colorado 80401
Department of Chemistry, Dartmouth College
Hanover, New Hampshire 03755
Department of Chemistry & Biochemistry
University of Colorado, Boulder, Colorado 80309

Received June 10, 1997

The examination of tetrafluorocyclobutadiene, C₄F₄ (**1**), illuminates striking differences between fluorocarbons and their hydrocarbon analogues.^{1,2} While cyclobutadiene, C₄H₄, has been studied extensively,^{3–6} due to interest in its antiaromaticity, so far only indirect evidence for the formation of **1** has been provided.⁷

Analysis of the IR spectrum of C₄F₄ leads us to believe that this unstable molecule is nonplanar. This result, though not anticipated by any treatment of C₄F₄ as a mere analog of cyclobutadiene, is reminiscent of the nonplanarity of the calculated structure for the perfluoroallyl radical.⁸

We observe the IR spectrum of **1** (Figure 1, center) when we irradiate tetrafluorocyclobutene-3,4-dicarboxylic anhydride (**2**)^{7a} isolated in an Ar matrix at 12 K with a 248 nm light from KrF excimer laser. In addition to C₄F₄, CO₂ and CO are formed. The C₄F₄ was also generated, with lower yield, by irradiation of *cis*- or *trans*-tetrafluoro-3,4-diiodocyclobutene.⁹



As for an IR spectrum of C₄F₄, we identify a set of peaks which (a) grows in at the same rate on irradiation of **2**, (b) is produced at the same rate that the precursor is destroyed, and (c) disappears at the same rate when **1** is destroyed by prolonged irradiation. The photochemical transformation of **2** to **1** is very clean, with only a trace amount of the perfluorocyclopenta-




Figure 1. Calculated (B3LYP/cc-pVDZ), unscaled spectrum of C₄F₄ (top). Calculated transition moment directions are referenced to the carbon skeleton plane and are determined in a counterclockwise sense about the C₂ axis. The experimental infrared absorption spectrum of tetrafluorocyclobutadiene (C₄F₄) isolated in Ar matrix at 12 K (center). For abbreviations see, Table 1. Linear dichroism spectrum obtained after photoorientation with linearly polarized light at 248 nm (bottom). C₄F₄ bands are marked with appropriate polarizations. Experimental transition moment directions were determined according to ref 13.

dienone^{7c} detected. The equimolar concentrations of CO₂, CO, and C₄F₄ in the matrix allow us to estimate the absolute intensities of the C₄F₄ absorptions.¹¹

Partial bleaching of the C₄F₄ by linearly polarized 248 nm light leads to partial photoorientation of **1**. Polarization measurements indicate that electronic absorption¹² at 248 nm proceeds along the C₂ axis, and the three a_u vibrations show negative dichroism in the IR; for each of them absorption along the photoselection axis Z (E_Z) is smaller than along direction Y (E_Y) perpendicular to it (E_Z < E_Y). At the same time, the two observed b_u modes show positive dichroism (Figure 1, bottom).

The fundamental vibrations for tetrafluorocyclobutadiene (Table 1) were identified by comparison of observed IR bands

[†]Dartmouth College.

[‡]University of Colorado.

[§]NREL.

(1) *Chemistry of Organic Fluorine Compounds II: a Critical Review*; Hudlicky, M.; Pavlath, A. E., Eds.; American Chemical Society: Washington, DC, 1995.

(2) *Organofluorine Chemistry, Principles and Commercial Applications*; Banks, R. E., Smart, B. E., Tatlow, J. C., Eds.; Plenum: New York, 1994.

(3) Bally, T.; Masamune, S. *Tetrahedron* **1980**, *36*, 343.

(4) Hess, B. A.; Carsky, P.; Schaad, L. *J. Am. Chem. Soc.* **1983**, *105*, 695.

(5) Maier, G. *Angew. Chem., Int. Ed. Engl.* **1988**, *27*, 309.

(6) Arnold, B. R.; Michl, J. in *Kinetics and Spectroscopy of Carbenes and Biradicals*; Platz, M. S., Ed.; Plenum Publishing Corporation: New York, 1990; pp 1–35. Bonačić-Koutecký, V.; Schöffel, K.; Michl, J. *J. Am. Chem. Soc.* **1989**, *111*, 6140.

(7) (a) Gerace, M. J.; Lemal, D. M.; Ertl, H. *J. Am. Chem. Soc.* **1975**, *97*, 5584. (b) Fanuele, J. C., Senior Honors Thesis, Dartmouth College, 1996. Earlier workers had transformed 3,4-diiodotetrafluorocyclobutene into a polymer of tetrafluorocyclobutadiene (Anderson, R. W.; Frick, H. R., US Pat. 3 682 876, 1972), a trimer of cyclobutadiene (Hertler, W. R. *J. Fluor. Chem.* **1975**, *6*, 171), and octafluorocyclooctatetraene (Barlow, M. G.; Crawley, M. W.; Haszeldine, R. N. *J. Chem. Soc., Perkin Trans. 1* **1980**, 122), but there was no clear evidence for the intermediacy of the cyclobutadiene in these reports. (c) Grayston, M. W.; Saunders, W. D.; Lemal, D. M. *J. Am. Chem. Soc.* **1980**, *102*, 413.

(8) Hammons, J. H.; Coolidge, M. N.; Borden, W. T. *J. Phys. Chem.* **1990**, *94*, 5468.

(9) 3,4-Dichlorotetrafluorocyclobutene (Haszeldine, R. N.; Osborne, J. E. *J. Chem. Soc.* **1955**, 3880) was transformed into the 3,4-diiodides as described by Dreyfuss, M. P., Ph.D. Dissertation, Cornell University, NY, 1957.

(10) Radziszewski, J. G.; Hess, B. A.; Zahradník, R. *J. Am. Chem. Soc.* **1992**, *114*, 52.

(11) Radziszewski, J. G.; Nimlos, M. R.; Winter, P. R.; Ellison, G. B. *J. Am. Chem. Soc.* **1996**, *118*, 7400.

(12) The UV absorption spectrum was obtained in the instrument where electronic and IR spectral observations are performed simultaneously. This allows for precise correlation between IR and UV absorption growth or decay rates during the course of photolysis at various wavelengths, and thus the extraction of the weak UV absorption bands.

(13) The experimental transition moment directions were determined according to the following formula:¹⁴ $\tan^2 \varphi = (K_x - K_y)/(K_x + K_y)$, where K_x and K_y are the principal orientation factors, and K_i values are the orientation factors of individual observed vibrations ν_i . Since the orientation is uniaxial, as dictated by the photoselected direction Z, we can adapt for $K_z = 0.318$ (an average of K_z values for ν_{10} , ν_{11} , and ν_{12}). From the condition that $\sum_i K_i = 1$ we determine $K_x = K_y = 0.341$. Although the error bars for our experimental transition moment direction angles are quite large (at least $\pm 12^\circ$), this result supports our claim for a nonplanar structure for **1**. The calculated transition moment directions are notoriously unreliable, and advanced *ab initio* treatment is required to achieve some acceptable precision.¹⁵


(14) Thulstrup, E. W.; Michl, J. *Elementary Polarization Spectroscopy*; VCH: New York, 1989. Michl, J.; Thulstrup, E. W. *Spectroscopy with Polarized Light. Solute Alignment by Photoselection, in Liquid Crystals, Polymers, and Membranes*; VCH: New York, 1986.

(15) Radziszewski, J. G.; Downing, J. W.; Gudipati, M. S.; Balaji, V.; Thulstrup, E. W.; Michl, J. *J. Am. Chem. Soc.* **1996**, *118*, 10275.

Table 1. Infrared Absorption Spectrum of Perfluorocyclobutadiene (C₄F₄)^a

| $\Gamma_{\text{vib}}(C_{2h}) = 5a_g \oplus 4b_g \oplus 5a_u \oplus 4b_u$ | | | | | | | | |
|--|----------------|---------------|-----|---------------|--------------------|-------|-------------|-------------------------|
| ν | exptl | | | | calcd ^b | | | |
| | sym | $\tilde{\nu}$ | I | K_i^c (pol) | $\tilde{\nu}$ | I | φ^d | assignment ^e |
| 1 | | — | — | — | 1867 | 0 | | C=C sym s |
| 2 | a _g | — | — | — | 1214 | 0 | | C-C sym s |
| 3 | | — | — | — | 673 | 0 | | rb |
| 4 | | — | — | — | 261 | 0 | | F-C-C-F sym b |
| 5 | | — | — | — | 154 | 0 | | F-C=C-F sym b |
| 6 | b _g | — | — | — | 1354 | 0 | | C-F sym s |
| 7 | | — | — | — | 755 | 0 | | ip ring t |
| 8 | | — | — | — | 489 | 0 | | ip ring d |
| 9 | | — | — | — | 415 | 0 | | ip ring d |
| 10 | a _u | 1291 | 535 | 0.319 (z) | 1306 | 509.7 | z | C-F sym s |
| 11 | | 914 | 32 | 0.322 (z) | 912 | 43.8 | z | ip ring d |
| 12 | | 596 | 37 | 0.313 (z) | 610 | 40.3 | z | rp |
| 13 | | 221 | 0.2 | — | 224 | 0.1 | z | F-C=C-F asym b |
| 14 | | — | — | — | 159 | 0.1 | z | C-F sym op b |
| 15 | b _u | 1683 | 143 | 0.337 (25°) | 1849 | 158.6 | 23.7° | C=C asym s ^f |
| 16 | | 968 | 101 | 0.340 (16°) | 976 | 193.3 | 7.4° | C-F asym s |
| 17 | | 300 | 4 | — | 303 | 5.2 | 20.1° | F-C=C-F asym sc |
| 18 | | 188 | 1 | — | 194 | 1.8 | -10.5° | C-F sym op d |

^a Argon matrix, 12 K; $\tilde{\nu}$, in cm⁻¹; I , absolute absorption intensity, in km/mol. ^b Calculated frequencies (B3LYP/cc-pVDZ) are unscaled. ^c K_i is the orientation factor for transition i ; $K_i = \langle \cos^2 i \rangle$, $i = x, y, \text{ or } z$, and i is the angle between the molecular axis i and the photoselection direction Z , averaged over all possible orientations, as indicated by the pointed brackets. ^d Calculated transition moment directions, the angles φ are determined with respect to the carbon skeleton plane. ^e Approximate mode description: sym = symmetric, asym = asymmetric, ip = in-plane, op = out-of-plane, s = stretch, b = bend, d = deformation, sc = scissors, t = twist, rb = ring breathing, rp = ring puckering. ^f The C=C stretching vibration, at 1683 cm⁻¹ proceeds in the direction perpendicular to the C=C bond, and its transition moment is inclined by nearly 24° from the plane created by the carbons.

**Figure 2.** Calculated structure of C₄F₄ (B3LYP/cc-pVDZ). Bond lengths are given in angstroms and angles are in degrees.

with the frequencies and intensities obtained from *ab initio* (B3LYP/cc-pVDZ) calculations. The optimized nonplanar structure of **1** is shown in Figure 2.

Our experimental data point to a nonplanar structure for C₄F₄. We observe a peak at 595 cm⁻¹ which we assign as the ring-puckering mode on the basis of its frequency and polarization and comparison with C₄H₄.¹⁶ This band shows negative dichroism, as expected for all modes active along the molecular z axis. Were C₄F₄ planar, symmetry would prevent a change in dipole moment for this mode and make it IR inactive. Were C₄F₄ to tautomerize, as does C₄H₄, we would record the same dichroism for all observed in-plane transitions. Instead, we measure distinctly different dichroic ratios for vibrations proceeding in the directions along and perpendicular to the C₂ axis (see Table 1).


The nonplanarity of tetrafluorocyclobutadiene is most likely the result of a conjunction of two effects. Electron withdrawal by the highly electronegative fluorines favors pyramidalization of the carbons, increasing the p-character in the C-F bonds, and this tendency is reinforced by the resulting attenuation of the cyclic conjugation in the π system. A natural atomic orbital (NAO) and natural bond orbital (NBO) population analysis of C₄F₄ (B3LYP/cc-pVDZ) supports this interpretation. The NBO calculation shows that the bonding orbitals of C₄F₄ are all nearly equivalent, having s and p contributions which resemble those of a tetrahedral sp³ hybrid for carbon atoms, in contrast to cyclobutadiene (C₄H₄) which exhibits the classical bonding

Table 2. B3LYP/cc-pVDZ NBO and NAO Population Analysis of Atomic Orbital Character in Bonding Orbitals

| C ₄ H ₄ (D _{2h}) | | | C ₄ F ₄ (C _{2h}) | | |
|--|-----|-----|--|-----|-----|
| orbital | % S | % P | orbital | % S | % P |
| C=C σ bond | 38 | 62 | C=C σ bond | 28 | 72 |
| C=C π bond | 0 | 100 | C=C π bond | 15 | 85 |
| C-C bond | 27 | 73 | C-C bond | 30 | 70 |
| C-H bond | 35 | 65 | C-F bond | 27 | 73 |
| ideal: sp ² | 33 | 67 | ideal: sp ³ | 25 | 75 |

characteristics of a carbon atoms with three equivalent sp² hybrid orbitals and one π -bonding orbital (Table 2).

The spectra of C₄F₄ show no indications of the valence tautomerization which is observed in C₄H₄.¹⁷ The bond-shift isomerization in C₄H₄ happens *via* heavy atom tunneling,¹⁸ and this is out-of-the-question for **1** in light of the large changes in geometry required.



Unlike the situation in C₄H₄, where interconversion of the two valence tautomers amounts to bond length alternations in the carbon skeleton accompanied by minimal movement of hydrogens, in C₄F₄ substantial additional change of the C-F bond angles (calculated to be over 23°) would be required in order for tautomerization to occur. Perhaps the matrix environment further inhibits tautomerization.¹⁷

Acknowledgment. This work is dedicated to Professor Günther Maier on the occasion of his 65th Birthday. This project was supported by the internal funds at NREL ("First Program"). E.J.P. was supported by Petroleum Research Fund, grant number 30676-AC6 and the National Science Foundation (CHE-9215164). Continuing support by Dr. Helena Lee Chum (NREL) is gratefully acknowledged.

JA971930U

(16) Orendt, A. M.; Arnold, B. R.; Radziszewski, J. G.; Facelli, J. C.; Malsch, K. D.; Strub, H.; Grant, D. M.; Michl, J. *J. Am. Chem. Soc.* **1988**, *110*, 2648.

(17) Arnold, B. R.; Radziszewski, J. G.; Campion, A.; Perry, S. S.; Michl, J. *J. Am. Chem. Soc.* **1991**, *113*, 692.

(18) Carpenter, B. K. *J. Am. Chem. Soc.* **1983**, *105*, 1700.