

## Total Synthesis of (–)-Laulimalide

Arun K. Ghosh\* and Yong Wang

Department of Chemistry  
University of Illinois at Chicago  
845 West Taylor Street  
Chicago, Illinois 60607

Received July 25, 2000

Laulimalide (**1**), also known as figianolide B, a 20-membered macrolide isolated from the Indonesian sponge *Hyattella* sp., has shown remarkable antitumor activities.<sup>1</sup> Recently, Laulimalide was also isolated from an Okinawan sponge *Fasciospongia rimosa*.<sup>2</sup> It displayed potent cytotoxicity against the KB cell line with an IC<sub>50</sub> value of 15 ng/mL.<sup>1</sup> Furthermore, it has shown cytotoxicity against P388, A549, HT29, and MEL28 cell lines in the range of 10–50 ng/mL (IC<sub>50</sub> values).<sup>2b</sup> The structure of **1** was initially established by NMR studies. Subsequently, its absolute configuration was established by X-ray analysis by Higa and co-workers.<sup>2</sup> The significant clinical potential of laulimalide has stimulated considerable interest in its synthesis and structure–function studies.<sup>3</sup> Herein, we report the first synthesis of (–)-laulimalide **1**.

As outlined in Figure 1, our synthetic strategy of laulimalide is convergent and involves the assembly of C<sub>3</sub>–C<sub>16</sub> segment **2** and C<sub>17</sub>–C<sub>28</sub> segment **3** by Julia olefination, followed by an intramolecular Horner–Emmons reaction between the C<sub>19</sub> phosphonoacetate and C<sub>3</sub> aldehyde. The sensitive epoxide at C<sub>16</sub>–C<sub>17</sub> was selectively introduced at the final stage of the synthesis by Sharpless epoxidation.<sup>4</sup> The construction of both dihydropyran rings of laulimalide was achieved by ring-closing olefin metathesis utilizing Grubbs' catalyst as the key step.<sup>5</sup>

The synthesis of C<sub>3</sub>–C<sub>16</sub> segment is accomplished by modifications of the previously published sequence.<sup>3a</sup> As outlined in Scheme 1, diastereomerically pure  $\delta$ -lactone **5** was prepared efficiently by exposure of acryloyl ester **4** to Grubbs' catalyst (10 mol %) in CH<sub>2</sub>Cl<sub>2</sub>.<sup>3a</sup> DIBAL reduction of **5** at –78 °C in CH<sub>2</sub>Cl<sub>2</sub> followed by reaction with ethanol and CSA afforded the ethyl glycoside.<sup>6</sup> Reaction of this ethyl acetal with *tert*-butyldimethylsilyl vinyl ether and Montmorillonite K-10 as the Lewis acid in CH<sub>2</sub>Cl<sub>2</sub> at 23 °C followed by NaBH<sub>4</sub> reduction of the resulting aldehyde afforded the corresponding dihydropyran as a single isomer (by <sup>1</sup>H and <sup>13</sup>C NMR). Protection of the alcohol with TBSCl and imidazole furnished the TBS ether **6**. Removal of the benzyl group by lithium in liquid ammonia provided the alcohol which was subsequently converted to iodide **7**. To install the C<sub>13</sub> methylene unit and the C<sub>15</sub> hydroxyl group, alkylation of lactone **8** was carried out by treatment with NaH in DMF at 0 °C for 15 min followed by reaction with iodide **7** at 23 °C for 15 min and then 60 °C for 12 h, which furnished lactone **9** as a mixture (4.2:1 by <sup>1</sup>H NMR) of isomers. Reduction of the mixture of lactone **9**





Figure 1.

Scheme 1<sup>a</sup>

<sup>a</sup> (a) Dibal-H, –78 °C then CSA, EtOH, 23 °C; (b) K-10, CH<sub>2</sub> = CHOTBS, 23 °C; (c) NaBH<sub>4</sub>, MeOH, 0 °C (54%); (d) TBSCl, imidazole, DMF, 23 °C (75%); (e) Li, NH<sub>3</sub> (95%); (f) I<sub>2</sub>, PPh<sub>3</sub>, Imidazole (96%); (g) **8**, NaH, DMF, 0 °C then iodide **7**, 60 °C (89%); (h) Red-Al, THF, 0 °C; (i) PhCOCl, Et<sub>3</sub>N, DMAP (cat.); (j) Na(Hg), Na<sub>2</sub>HPO<sub>4</sub>, MeOH, –20 °C to 23 °C (72%); (k) MOMCl, *i*-Pr<sub>2</sub>NEt, 23 °C; (l) DDQ, pH 7 buffer, 23 °C (81%); (m) DMSO, (COCl)<sub>2</sub>, *i*-Pr<sub>2</sub>NEt, –60 °C (85%).

by Red-Al provided the diol **10**. Benzylation of **10** afforded the corresponding dibenzoate which was exposed to Na–Hg in MeOH at –20 °C to provide the olefin **11**.<sup>7</sup> Protection of the C<sub>15</sub> hydroxyl group as a MOM ether, removal of the PMB group and Swern oxidation of the resulting alcohol furnished the C<sub>3</sub>–C<sub>16</sub> segment, aldehyde **2**.

The synthesis of the C<sub>17</sub>–C<sub>28</sub> segment **3** has been achieved by modification of reaction sequences published previously.<sup>3b</sup> Optically active dihydropyran derivative **12** was prepared in multigram quantity by utilizing ring-closing olefin methathesis<sup>5</sup> and Corey–Fuchs' homologation<sup>8</sup> reactions as the key steps. The alkynyl anion of **12** was readily obtained by treatment with *n*-BuLi at –78 °C for 1 h followed by warming to 23 °C for 1 h (Scheme 2). Reaction of the resulting alkynyl anion with optically active

(1) (a) Quinoa, E.; Kakou, Y.; Crews, P. *J. Org. Chem.* **1988**, *53*, 3642. (b) Corley, D. G.; Herb, R.; Moore, R. E.; Scheuer, P. J.; Paul, V. J. *J. Org. Chem.* **1988**, *53*, 3644.

(2) (a) Jefford, C. W.; Bernardinelli, G.; Tanaka, J.-I.; Higa, T. *Tetrahedron Lett.* **1996**, *37*, 159. (b) Tanaka, J.-I.; Higa, T.; Bernardinelli, G.; Jefford, C. W. *Chem. Lett.* **1996**, 255.

(3) (a) Ghosh, A. K.; Wang, Y. *Tetrahedron Lett.* **2000**, *41*, 2319. (b) Ghosh, A. K.; Wang, Y. *Tetrahedron Lett.* **2000**, *41*, 4705. (c) Mulzer, J.; Hanbauer, M. *Tetrahedron Lett.* **2000**, *41*, 33. (d) Shimizu, A.; Nishiyama, S. *Synlett* **1998**, 1209. (e) Ghosh, A. K.; Mathivanan, P.; Cappiello, J. *Tetrahedron Lett.* **1997**, *38*, 2427.


(4) Johnson, R. A.; Sharpless, K. B. In *Catalytic Asymmetric Synthesis*; Ojima, I., Ed.; VCH Publishers: New York, 1993, p 103–158.

(5) Grubbs, R. H.; Chang, S. *Tetrahedron* **1998**, *54*, 4413 and references therein.

(6) For a related method, see: Dahanukar, V. H.; Rychnovsky, S. D. *J. Org. Chem.* **1996**, *61*, 8317.

(7) Lee, G. H.; Lee, H. K.; Choi, E. B.; Kim, B. T.; Pak, C. S. *Tetrahedron Lett.* **1995**, *36*, 5607.

(8) Corey, E. J.; Fuchs, P. L. *Tetrahedron Lett.* **1972**, 3769.


Scheme 2<sup>a</sup>

<sup>a</sup> (a) *n*-BuLi,  $-78\text{ }^{\circ}\text{C}$ , 1 h and  $23\text{ }^{\circ}\text{C}$ , 1 h then **13**,  $-78\text{ }^{\circ}\text{C}$  (64%); (b) Dess–Martin,  $\text{CH}_2\text{Cl}_2$ ,  $23\text{ }^{\circ}\text{C}$  (81%); (c) L-Selectride, THF,  $-78\text{ }^{\circ}\text{C}$  (87%); (d) Red-Al, THF,  $-20\text{ }^{\circ}\text{C}$  (81%); (e)  $\text{CF}_3\text{CO}_2\text{H}$ ,  $\text{CH}_2\text{Cl}_2$ ,  $23\text{ }^{\circ}\text{C}$ ; (f) *p*-MeO-Ph-CH(OMe)<sub>2</sub>, CSA,  $\text{CH}_2\text{Cl}_2$ ,  $23\text{ }^{\circ}\text{C}$  (71%); (g) Dibal-H,  $\text{CH}_2\text{Cl}_2$ ,  $-78\text{ }^{\circ}\text{C}$  (74%).

aldehyde **13** at  $-78\text{ }^{\circ}\text{C}$  furnished a diastereomeric mixture (1.8:1) of alcohols **14** and **15**, which upon oxidation with Dess–Martin periodinane<sup>9</sup> gave the corresponding alkynyl ketone. L-Selectride reduction of this ketone at  $-78\text{ }^{\circ}\text{C}$  furnished the desired *syn*-alkynyl alcohol **15** as a single diastereomer (by <sup>1</sup>H NMR and <sup>13</sup>C NMR). Red-Al reduction of **15** set the C<sub>21</sub>–C<sub>22</sub> *trans*-olefin geometry. Removal of the C<sub>19</sub> PMB group by exposure to TFA<sup>10</sup> and subsequent reaction of the resulting diol with *p*-methoxybenzylidene acetal and CSA provided acetal **16**. DIBAL reduction of **16** at  $-78\text{ }^{\circ}\text{C}$  in  $\text{CH}_2\text{Cl}_2$  afforded the C<sub>17</sub>–C<sub>28</sub> segment **3** as a single regio isomer (by <sup>1</sup>H and <sup>13</sup>C NMR).

Our subsequent synthetic strategy calls for the assembly of fragments **2** and **3** by Julia olefination (Scheme 3). Thus, lithiation of sulfone derivative **3** with 2.1 equiv of *n*-BuLi in THF at  $-78\text{ }^{\circ}\text{C}$  for 15 min followed by reaction of the resulting dianion with the aldehyde **2** at  $-78\text{ }^{\circ}\text{C}$  to  $-40\text{ }^{\circ}\text{C}$  for 2 h furnished the expected  $\alpha$ -hydroxy sulfone derivatives. The hydroxy sulfone was transformed into the corresponding C<sub>16</sub>–C<sub>17</sub> olefin in a two step sequence involving (1) acylation of the hydroxy sulfone derivatives with Ac<sub>2</sub>O, Et<sub>3</sub>N and DMAP (cat.) and (2) exposure of the resulting acetates to Na(Hg) in methanol at  $-20\text{ }^{\circ}\text{C}$  for 2 h followed by warming the reaction to  $23\text{ }^{\circ}\text{C}$  for 30 min. The C<sub>16</sub>–C<sub>17</sub> *trans* olefin **17** was obtained in 34% yield along with 10% *cis*-olefin, which was readily separated by silica gel chromatography.

Subsequent elaboration to the macrolactone possessing C<sub>2</sub>–C<sub>3</sub> *cis*-olefin geometry proved to be a formidable task. We finally relied upon an intramolecular Horner–Emmons reaction of C<sub>19</sub> phosphonoacetate and C<sub>3</sub> aldehyde using Still's protocol.<sup>11</sup> Thus, acylation of C<sub>19</sub> hydroxyl group with bis-(2,2,2-trifluoroethyl)-phosphonoacetic acid followed by removal of the TBS group by exposure to aqueous acetic acid in THF at  $23\text{ }^{\circ}\text{C}$  furnished the acetate derivative **18** in near quantitative yield.<sup>12</sup> Oxidation of **18** with Dess–Martin periodinane provided the C<sub>3</sub> aldehyde which upon treatment with K<sub>2</sub>CO<sub>3</sub> in the presence of 18-C-6 at  $-20\text{ }^{\circ}\text{C}$  for 30 min and then at  $0\text{ }^{\circ}\text{C}$  for 2.5 h furnished a mixture (2:1) of macrolactones **19** and **20** in 84% yield (Scheme 3). Both *cis*- and *trans*-lactones were separated by silica gel chromatography. Horner–Emmons reaction of the corresponding (diphenylphosphono)acetate derivative provided slight improvement of the *cis*-selectivity (*cis:trans* = 1:1.7, 69% isolated yield).<sup>13</sup> Further attempts to improve the ratio for **20** by changing reaction conditions or C<sub>20</sub> protecting group have been unsuccessful. The overall yield of the desired *cis*-macrolactone **20** was however

Scheme 3<sup>a</sup>

<sup>a</sup> (a) *n*-BuLi,  $-78\text{ }^{\circ}\text{C}$ , 15 min, Then **2**,  $-78$  to  $-40\text{ }^{\circ}\text{C}$ , 2 h; (b) Ac<sub>2</sub>O, Et<sub>3</sub>N, DMAP (cat.); (c) Na(Hg), Na<sub>2</sub>HPO<sub>4</sub>, MeOH,  $-20$  to  $23\text{ }^{\circ}\text{C}$  (34%); (d)  $(\text{CF}_3\text{CH}_2\text{O})_2\text{P}(\text{O})\text{CH}_2\text{CO}_2\text{H}$ , Cl<sub>3</sub>C<sub>6</sub>H<sub>2</sub>COCl, *i*-Pr<sub>2</sub>NEt, DMAP; (e) AcOH–THF–H<sub>2</sub>O (3:1:1),  $23\text{ }^{\circ}\text{C}$  (99%); (f) Dess–Martin,  $\text{CH}_2\text{Cl}_2$ ,  $23\text{ }^{\circ}\text{C}$  (79%); (g) K<sub>2</sub>CO<sub>3</sub>, 18-C-6,  $-20$  to  $0\text{ }^{\circ}\text{C}$  (84%); (h) *h* $\nu$ , Et<sub>2</sub>O, 50 min (66%); (i) PPTS, *t*-BuOH,  $84\text{ }^{\circ}\text{C}$  (45%) (j) Ti(O<sup>*i*</sup>Pr)<sub>4</sub>, (+)-DET, *t*-BuOOH,  $-20\text{ }^{\circ}\text{C}$ ; (k) DDQ, pH 7,  $23\text{ }^{\circ}\text{C}$  (48%).

improved to 47% after photoisomerization of the *trans*-macrolactone **19**.<sup>14</sup> Thus, irradiation of **19** in ether under UV in a Rayonet photochemical reactor for 50 min afforded a mixture of *trans*-lactone **19** (33%) and the *cis*-lactone **20** (33%) which were separated by chromatography. The identity of the *cis*-olefin geometry of **20** was established by its observed coupling constant ( $J = 11.6\text{ Hz}$ ). Macrolactone **20** was converted to synthetic (–)-laulimalide **1** as follows: removal of the MOM group by refluxing with PPTS in *t*-BuOH,<sup>15</sup> exposure of the resulting alcohol to Sharpless epoxidation<sup>4</sup> with (+)-DET and removal of the C<sub>20</sub> PMB ether by exposure to DDQ. Spectral data (<sup>1</sup>H and <sup>13</sup>C NMR) of synthetic **1** ( $[\alpha]_D^{23} -196\text{ c } 0.23$ , CHCl<sub>3</sub>) are identical to that from a sample of natural laulimalide (lit.<sup>2a</sup>  $[\alpha]_D^{29} -200\text{ c } 1.03$ , CHCl<sub>3</sub>) kindly provided by Professor Higa.

Thus, a stereocontrolled synthesis of (–)-laulimalide has been achieved. Considering its clinical potential as an antitumor agent, the present synthesis will enable important structure–function studies as well as synthesis of structural variants of laulimalide. Further improvement in synthesis and biological studies are currently in progress.

**Acknowledgment.** Financial support by the National Institutes of Health (GM 55600) is gratefully acknowledged. We also thank Professor Higa for providing a sample of natural laulimalide and Professor Forsyth for the experimental details of bis-(2,2,2-trifluoroethyl)phosphonoacetic acid.

**Supporting Information Available:** Experimental procedures and spectral data for compounds **1–3**, **6–11**, **15–20**; <sup>1</sup>H NMR spectra for compounds **1**, **3**, **11**, **17**, and **19** (PDF). This material is available free of charge via the Internet at <http://pubs.acs.org>.

JA0027416

(9) Meger, S. D.; Schreiber, S. L. *J. Org. Chem.* **1994**, *59*, 7549.(10) Yan, L.; Kahne, D. *Synlett* **1995**, 523.(11) Still, W. C.; Gennari, C. *Tetrahedron Lett.* **1983**, *24*, 4405.(12) This protocol was recently employed in the synthesis of phorbokazole A. See: Forsyth, C. J.; Ajmed, F.; Cink, R. D.; Lee, C. S. *J. Am. Chem. Soc.* **1998**, *120*, 5597.(13) Ando, K. *J. Org. Chem.* **1999**, *64*, 8406 and references therein.(14) Smith, A. B.; Lupio, A. T.; Ohba, M.; Chen, K. *J. Am. Chem. Soc.* **1989**, *111*, 6648 and references therein.(15) Monti, H.; Leandri, G.; Klos-Ringquet, M.; Corriol, C. *Synth. Commun.* **1983**, *13*, 1021.