

Chromophore Formation in Resorcinarene Solutions and the Visual Detection of Mono- and Oligosaccharides

Ming He, Rolanda J. Johnson, Jorge O. Escobedo, Patricia A. Beck, Kyu Kwang Kim, Nadia N. St. Luce, Claude J. Davis,[†] Patrick T. Lewis,[‡] Frank R. Fronczek, Bruce J. Melancon, Anthony A. Mrse,[§] W. Dale Treleaven, and Robert M. Strongin*

Contribution from the Department of Chemistry, Louisiana State University, Baton Rouge, Louisiana 70803

Received December 8, 2001

Abstract: The colorimetric properties of resorcinarene solutions had not been investigated since Baeyer's initial synthesis. We recently reported that solutions containing resorcinarene macrocycles develop color upon heating or standing. In the presence of saccharides, these solutions exhibit significant color changes which are easily seen. We herein present strong evidence that the solution color is due to macrocycle ring opening and oxidation. The optical responses to saccharides are due to complexation of the sugar with the acyclic chromophores. We apply these mechanistic insights toward the challenging problem of the visual detection of neutral oligosaccharides by simple chromogens. In addition, we also report the first single-crystal X-ray crystal structure determination of a rarely observed "diamond" resorcinarene stereoisomer.

Introduction

In 1872, a year after his landmark synthesis of the xanthen dye fluoresceine,¹ Baeyer studied the condensation reaction of benzaldehyde and resorcinol in the presence of acid.² The red solution containing his product mixture changed to violet upon the addition of base. It is now well known that Baeyer's reaction afforded the first resorcinarenes, cyclic tetramers of resorcinol.³

The broad impact of the resorcinarenes in the fields of molecular recognition, materials science, and supramolecular chemistry has been the subject of extensive study as noted in several recent reviews.³ For instance, they were the first compounds shown to bind sugars in apolar media.⁴ Because boronic acids are the basis of carbohydrate affinity chromatography, we reasoned that resorcinarenes functionalized with boronic acids would constitute powerful sugar receptors. We thus synthesized **1** and **2a** and investigated their properties in the presence of sugars.⁵

We observed 11 different solution colors of 11 different heated sugar solutions containing **1**.⁶ The analytes included structurally related neutral carbohydrates, glucose phosphates, carboxylic acid, and amino sugars. The solution color changes were rapid, quantifiable, and reproducible. We also showed that solutions containing acetaldehyde-derived **2b** exhibited relatively paler color changes in the presence of sugars as compared to those containing boronic acids **1** and **2a** or acyclic congener **3a** (Figure 1).⁷ The colorimetric properties of resorcinarene solutions had not received prior attention since Baeyer's initial report.²

Sugars are a relatively challenging class of compounds to analyze. They exhibit great structural similarity as well as transparency in the visible region. The sensing of specific saccharides could aid the monitoring of disease or industrial fermentation products. The visual determination of saccharides has been of interest for well over a century.

Resorcinol and its derivatives have long been used in simple color tests for sugars. In 1887, Seliwanoff reported a resorcinol color test which was followed by other resorcinol-derived methods.⁸ These latter and numerous other related reducing sugar assays, while based on simple reagents, typically require toxic materials, harsh and often tedious procedures.⁹ In the 1990s, significant progress was made toward the improved

* To whom correspondence should be addressed. Phone: 225-578-3238. Fax: 225-578-3458. E-mail: rstrong@lsu.edu.

[†] Present address: Kraft Foods, Inc., Research and Development, 801 Waukegan Road, Glenview, IL 60025.

[‡] Present address: United States Patent and Trademark Office, Crystal Mall 1, 8-D-12, Washington, D.C. 20231.

[§] Present address: Department of Chemistry and Biochemistry, University of California, San Diego, 9500 Gilman Drive, La Jolla, CA 92093-0332.

(1) Baeyer, A. *Chem. Ber.* **1871**, *5*, 255.

(2) (a) Baeyer, A. *Ber. Dtsch. Chem. Ges.* **1872**, *5*, 25. (b) Baeyer, A. *Ber. Dtsch. Chem. Ges.* **1872**, *5*, 280.

(3) (a) Schneider, H.-J.; Schneider, U. *J. Inclusion Phenom.* **1994**, *19*, 67. (b) Cram, D. J.; Cram, J. M. *Container Molecules and Their Guests*; The Royal Society of Chemistry: Cambridge, U.K., 1994. (c) Sherman, J. C. *Tetrahedron* **1995**, *51*, 3395. (d) Timmerman, P.; Verboom, W.; Reinhoudt, D. N. *Tetrahedron* **1996**, *52*, 2663. (e) Jasat, A.; Sherman, J. C. *Chem. Rev.* **1999**, *99*, 931. (f) Rudkevich, D. M.; Rebek, J. *Eur. J. Org. Chem.* **1999**, *9*, 1991.

(4) Aoyama, Y.; Tanaka, Y.; Sugihara, S. *J. Am. Chem. Soc.* **1989**, *111*, 5397.

(5) Lewis, P. T.; Davis, C. J.; Saraiva, M.; Treleaven, W. D.; McCarley, T. D.; Strongin, R. M. *J. Org. Chem.* **1997**, *62*, 6110.

(6) Davis, C. J.; Lewis, P. T.; McCarroll, M. E.; Read, M. W.; Cueto, R.; Strongin, R. M. *Org. Lett.* **1999**, *1*, 331.

(7) Lewis, P. T.; Davis, C. J.; Cabell, L. A.; He, M.; Read, M. W.; McCarroll, M. E.; Strongin, R. M. *Org. Lett.* **2000**, *2*, 589.

(8) For example: (a) Seliwanoff, T. *Chem. Ber.* **1887**, *20*, 181. (b) Bial, M. *Dtsch. Med. Wochenschr.* **1902**, 253. (c) Kulka, R. G. *Biochem. J.* **1956**, *63*, 542.




Figure 1. We previously reported that solutions containing resorcinarenes **1**, **2a**, **2b**, and related condensation product **3a** exhibit significant color changes in the presence of sugars.

selective and mild detection of monosaccharides via relatively strong solution color changes observable by visual inspection. The recent advances were due mainly to the pioneering efforts of Shinkai and co-workers. Their studies were based primarily on aniline-functionalized azo dyes containing appended arylboronic acids.¹⁰

The field of glycobiology has recently undergone great resurgence due to the exciting therapeutic potential of oligosaccharides.¹¹ A vast number of oligosaccharides are found in glycoproteins and on cell surfaces. Most of the recent progress toward the color detection of sugars has, however, involved monosaccharide analysis.^{10,12} The great variety of linear and branched oligosaccharides magnifies the problems associated with monosaccharide analysis.

New methods allowing for simple and rapid oligosaccharide detection are in demand. The colorimetric detection of oligosaccharides eluting from chromatographic columns is one of the biggest challenges in sugar analysis.¹³ The older color tests used for automated HPLC postcolumn detection of monosaccharides

fail to effectively directly detect oligosaccharides containing more than three residues. For example, in a representative assay, maltohexaose's response is only 18% of that observed for the same weight of glucose.¹³ Colorimetric methods thus typically require prior complete hydrolysis to monosaccharides or covalent attachment of the oligosaccharides to a chromophore, which can lead to diminished separation.^{13,14}

Herein we present the first evidence that xanthenes form and serve as the active chromophores in resorcinarene solutions. We also describe our investigations of the saccharide-induced optical signal transduction mechanism. We extend the scope and utility of our sensing methodology to include the color detection of a homologous series of prototypical neutral oligosaccharides, the linear maltodextrins, containing up to seven glucose residues. We thus address the fundamental problem of diminished optical detection of larger oligosaccharides using simple colorimetric agents.

Results

Formation and Structure of the Chromophore in Resorcinarene Solutions. We reported the synthesis of compounds **1** and **2a** in one step in a combined 90% yield via the HCl-promoted condensation of commercially available 4-formylphenyl boronic acid and resorcinol.⁵ Separation of the two stereoisomers by fractional crystallization afforded white crystalline solids. We obtained X-ray quality crystals of the half-methyl tetraboronate ester of **1** upon slow crystallization from a 9:1 MeOH:EtOH solution. The solid-state architecture was characterized as an infinite, antiparallel two-dimensional network of macrocycles, each of which exhibited 12 intermolecular hydrogen bonds.¹⁵

We found that colorless DMSO solutions of freshly crystallized **1** or **2a** (5.2 mM), upon standing in solution for several hours or upon heating at 90 °C for 1 min, developed a pinkish-purple color. The color formation was monitored via UV-vis spectroscopy. The appearance of a new λ_{max} at 535 nm was accompanied by a less intense absorbance at 500 nm.⁶

Initial attempts at understanding the origin of the solution color involved heating solutions of **1** in the dark or in O₂ degassed conditions. In both cases, we found that the color intensities diminished, as evidence by both visual inspection and UV-vis spectroscopy.⁶ For instance, heating a solution of **1** (5.2 mM in DMSO) under O₂ degassed conditions led to a 61% decrease in absorbance at 536 nm. Light and O₂ apparently promote color formation. In addition, when we acylated the phenolic hydroxyls of **1** and heated a DMSO solution of the resultant octaacetate to reflux, the solution remained colorless.⁷ The phenolic hydroxyls thus also play a key role in chromophore formation. We reasoned that the chromophore arises via oxidation of a resorcinol moiety to a quinone.^{6,7}

We also heated solutions of resorcinol or benzeneboronic acid separately or as an equimolar mixture using the aforementioned conditions and concentrations, with and without added monosaccharides. We observed only very faint solution colors by visual inspection.⁶ This result showed that a methine-bridged resor-

(9) Chaplin, M. F. *Monosaccharides*. In *Carbohydrate Analysis. A Practical Approach*; Chaplin, M. F., Kennedy, J. F., Eds.; Oxford University Press: Oxford, 1994; pp 1–40.
 (10) Review: (a) James, T. D.; Samankumara Sandanayake, K. R. A. S.; Shinkai, S. *Angew. Chem., Int. Ed. Engl.* **1996**, *35*, 1910. More recent examples of boronic acid-based dyes and optical sensing of neutral sugars: (b) Koumoto, K.; Takeuchi, M.; Shinkai, S. *Supramol. Chem.* **1998**, *9*, 203. (c) Koumoto, K.; Shinkai, S. *Chem. Lett.* **2000**, 856. (d) Ward, C. J.; Patel, P.; Ashton, P. R.; James, T. D. *Chem. Commun.* **2000**, 229. (d) DiCesare, N.; Lakowicz, J. R. *Org. Lett.* **2001**, *3*, 3891.
 (11) Recent examples: (a) Bertozzi, C. R.; Kiessling, L. L. *Science* **2001**, *291*, 2357. (b) Axford, J. *Trends Immunol.* **2001**, *22*, 237 and references therein.
 (12) Solution color changes in response to monosaccharides, maltose, maltotriose, and maltotetraose using a resorcinarene-methyl red dye competition process: Rusin, O.; Kral, V. *Tetrahedron Lett.* **2001**, *42*, 4235.

(13) Kennedy, J. F.; Pagliuca, G. *Oligosaccharides*. In *Carbohydrate Analysis. A Practical Approach*; Chaplin, M. F., Kennedy, J. F., Eds.; Oxford University Press: Oxford, 1994; pp 46–62.
 (14) Review: LoGuidice, J. M.; Lhermitte, M. *Biomed. Chromatogr.* **1996**, *10*, 290.
 (15) Davis, C. J.; Lewis, P. T.; Billodeaux, D. R.; Fronczek, F. R.; Escobedo, J. O.; Strongin, R. M. *Org. Lett.* **2001**, *3*, 2443.






Figure 2. **2a** (1.0 mg) and **3a** (1.0 mg) each in 0.9 mL of DMSO were heated to gentle reflux over 2 min and cooled to room temperature before 0.1 mL of H₂O was added to each solution. The final concentrations of **2a** and **3a** in 9:1 DMSO:H₂O are 1.03×10^{-3} and 1.96×10^{-3} M, respectively. A solution of **6b** (5.0×10^{-6} M) was prepared at room temperature in 9:1 DMSO:H₂O.

cinol/aldehyde condensation framework was needed for effective chromophore formation and optical sugar detection. Interestingly, methine-bridged condensation product resorcinarene substructures, of which **3a–c** are examples, were noted as

Scheme 1. Dehydration and Oxidation of Methine-Bridged Resorcinol Oligomers Leading to a Xanthene



reaction intermediates in standard xanthene dye syntheses (e.g., the transformation of **4** to **5**, $n = m = 0$, Scheme 1).¹⁶



Xanthenes are some of the oldest known synthetic dyes such as fluorescein, rhodamine, **6a** and **6b** and ethyl eosin (**6c**), as well as many others. Importantly, the colorimetric properties of xanthenes are a function of the ionization state of the C-6 moiety.¹⁷ They typically exhibit two absorbance maxima in the visible region.¹⁷ The absorption spectrum of **6b** (5.0×10^{-6} M) in 9:1 DMSO:H₂O is shown in Figure 2. It exhibits a λ_{max}

at 530 nm and a less intense λ_{max} at 500 nm. Interestingly, the λ_{max} absorbance values and spectral features are strikingly similar to those observed for colored DMSO solutions of **1** as well as **2a**, **2b**, and **3a** which we previously reported.⁷

Incorporation of a planar xanthene within a resorcinarene macrocycle framework via the transformation shown in Scheme 1 would lead to a considerable increase in strain energy. Simulations (Sybil 6.6) show that an increase in strain energy of 34.2 kcal/mol would occur upon formation of a xanthene substructure within **2b**. Prior studies of the related calixarenes (macrocycles formally derived from phenol/formaldehyde condensations) showed that xanthenes did not form in cyclic tetrameric structures.¹⁸

Ring opening to acyclic oligomers could thus be a prerequisite for xanthene formation from resorcinarenes. It is known that the condensation reactions producing resorcinarenes are reversible under acidic conditions.³ The detailed mechanism of resorcinarene macrocycle genesis has been studied thoroughly by Weinelt and Schneider.¹⁹ They found that **2b** and its macrocyclic stereoisomers interconverted via the intermediacy of acyclic oligomers. Their studies included the rapid quenching of condensation reactions between resorcinol and either acetaldehyde or paraldehyde in MeOH in the presence of anhydrous HCl. Because the opening of a resorcinarene ring has only been previously shown to occur upon the addition of strong acid, our hypothesis of acyclic oligomer formation in aqueous or neat DMSO solutions without added acid warrants further analysis.

We had observed that ¹H and ¹³C NMR spectra of DMSO-*d*₆ solutions of **1** (5.2 mM), heated at 90 °C for 3 min (our initial sugar colorimetric detection conditions), exhibited no readily observable change in chemical shifts or peak area integrals as compared to fresh, colorless samples.⁶ Xanthenes are strongly absorbing materials which need be only produced in trace (ca. 0.5% conversion; see, for example, the concentrations and absorbances shown in Figure 2) amounts to afford solution colors under our conditions.

Evidence for Acid Formation in DMSO Solutions. We find that more vigorous thermolytic conditions are necessary to afford conversion to significant amounts of products. Heating a DMSO (10 mL) solution of freshly recrystallized **2b** (100 mg, 18.4 mM) for 8 h at 120 °C followed by analysis via reversed phase HPLC reveals the formation of numerous new products representing a 74% conversion of **2b** to products based on relative peak areas.

Strong literature precedent allows us to propose that acyclic oligomers arise from **2b** via the in situ formation of strong acids. It is known that acid production from DMSO is promoted by the presence of O₂ and peroxides.²⁰ In addition, in situ acid formation has been attributed as the cause of certain oxidations in DMSO.^{20b} We have noted the effect of O₂ on resorcinarene solution color intensity (vide supra).

Acid formation observed during DMSO decomposition has been inhibited by free radical scavengers.^{20c} Under the same thermolysis conditions as noted above, but in the presence of free radical scavengers (either BHT or phenothiazene, 10 mol %), we observe less than 28% conversion to products by HPLC analysis.

(18) Agbaria, K.; Biali, S. E. *J. Org. Chem.* **2001**, *66*, 5842.

(19) Weinelt, F.; Schneider, H.-J. *J. Org. Chem.* **1991**, *56*, 5527.

(20) (a) Gillis, B. T.; Beck, P. E. *J. Org. Chem.* **1963**, *28*, 1388. (b) Santusosso, T. M.; Swern, D. *Tetrahedron Lett.* **1974**, 4255. (c) Emert, J.; Goldenberg, M.; Chiu, G. L. *J. Org. Chem.* **1977**, *42*, 2012.

(16) Sen, R. N.; Sinha, N. N. *J. Am. Chem. Soc.* **1923**, *45*, 2984.

(17) Gupta, S. N.; Linden, S. M.; Wrzyszczyński, A.; Neckers, D. C. *Macromolecules* **1988**, *21*, 51.




Figure 3. Compound **7** and ORTEP.

Important further evidence that strong acids form under our conditions was presented in our recent report describing the first X-ray crystal structure of $(\text{CH}_3)_3\text{S}^+\text{CH}_3\text{SO}_3^-$.²¹ The latter compound was obtained from a thermolysis reaction of **2b** in DMSO. It is known that $(\text{CH}_3)_3\text{S}^+\text{CH}_3\text{SO}_3^-$ forms, along with $\text{CH}_3\text{SO}_3\text{H}$, $\text{CH}_3\text{SO}_2\text{H}$, and CH_3SOH (and several other products), via the radical- and acid-promoted decomposition of DMSO.²²

Macrocycle Bond Breaking and Oxidation of the Acyclic Products. We next turned our attention to identifying the products of macrocycle ring opening. We isolate **7** (Figure 3), a rarely observed “diamond” resorcinarene stereoisomer,²³ in 2.3% yield from the thermolysis of **2b** in DMSO, via flash column chromatography. The structure of **7** was previously assigned (as the octabutyrate derivative) via NMR evidence during the acid-catalyzed condensation/isomerization studies of Schneider and Weinelt.¹⁹

Single crystals suitable for X-ray analysis, grown via slow evaporation of a 9:1 CH_2Cl_2 :MeOH solution of **7**, allow us to report the first known crystal structure of a resorcinarene diamond isomer. The macrocyclic molecule lies on a mirror plane in the crystal, with three methyl groups (C10, C8, and its mirror-related equivalent) relatively syn. The fourth methyl group, C18, lies on the mirror, and is anti to the other three. The resulting conformation is such that OH groups O1 and O4 both form intramolecular hydrogen-bonding contacts with their mirror-related equivalents on adjacent aromatic rings. The H atoms in these contacts are disordered, with the two O atoms alternately donor and acceptor. The $\text{O}\cdots\text{O}$ distances are 2.712(2) Å for O1 and 2.671(2) Å for O4. There are two independent




Figure 4. Top: Expansion of a ^1H NMR spectrum of semipurified thermolysis reaction products of **2b** showing the formation of **3b**. Bottom: Expansion of a ^1H NMR spectrum of pure **3b**.

methanol molecules, both lying on mirrors and forming hydrogen bonds with external macrocyclic OH groups O2 and O3. Importantly, stereoisomer **7** can only form from **2b** via bond rupture and reformation.¹⁹ If **7** were a conformer of **2b**, the

Table 1. MALDI MS Evidence for the Formation of Acyclic Oxidized and Unoxidized Products from the Thermolysis of **2b**

entry	structure	TLC R_f	(m/z) calcd	(m/z) obsd
1	4 , R = Me, $m = 1$, $n = 0$	0.29	382.41	381.89
2	4 , R = Me, $m = 3$, $n = 2$	0.44	926.36	926.28
3	6a	0.44	226.23	225.61
4	5 , R = Me, $m + n = 4$	0.26	906.01	906.33
5	5 , R = Me, $m + n = 3$	0.84	770.79	770.82
6	5 , R = Me, $n = 1$, $m = 0$	0.79	362.51	361.38

methyl group (C18) would reside outside, rather than above, the plane of the macrocycle cavity.

We also observe acyclic products during the thermolysis of **2b**. A key product, **3b**, is found in a broad HPLC fraction eluting from 18 to 20 min. The ^1H NMR spectrum of the isolate exhibits several peaks including each of the resonances associated with **3b**¹⁹ (CH_3OD δ 1.46, d, $^3J = 7.3$ Hz, 4.53, q, $^3J = 7.3$ Hz, 6.18–6.22, m, 6.89, d, $^3J = 8.0$ Hz). Overlay of the ^1H NMR spectrum of the HPLC isolate with a sample of independently synthesized and isolated **3b** confirms the assignment (Figure 4). In addition, the MALDI MS of the HPLC fraction contains a peak at 245.59 amu (246.26 amu calcd). The production of compounds **3b** and **7** under our conditions constitutes an important initial link between our investigations and the prior acid-catalyzed macrocycle genesis mechanism studies.¹⁹

We also find evidence for higher order oligomer production under our conditions (vide supra) involving thermolysis of **2b** in DMSO. At least five sets of doublets appear between 0.72 and 1.53 ppm in the ^1H NMR of each of two flash column fractions (TLC $R_f = 0.54$ and 0.63, 9:1 CH_2Cl_2 : CH_3OH , δ 1.53, 1.08, 1.01, 0.97, 0.83, 0.72 ppm, and δ 1.29, 1.15, 1.00, 0.89, 0.84 ppm, CH_3OD , respectively). In addition, the MALDI mass spectra (anthracene matrix) of other fractions ($R_f = 0.29$ and 0.44) exhibits peaks for higher homologues of **3b** (entries 1 and 2, Table 1). MALDI MS evidence also suggests the formation of xanthene materials not previously reported in previous fragmentation and equilibration studies of **2b** (entries 3–6, Table 1).^{24,25}


To study the formation of the oxidation products, we heat **3b**, the parent acyclic unoxidized homologue attained via ring opening of **2b**. Heating an air-saturated solution of **3b** (0.880

(21) Fronczek, F. R.; Johnson, R. J.; Strongin, R. M. *Acta Crystallogr.* **2001**, *E5*, o447–o448.

(22) (a) Chen, C.-T.; Yan, S.-J. *Tetrahedron Lett.* **1969**, 3855. (b) Head, D. L.; McCarty, C. G. *Tetrahedron Lett.* **1973**, 1405.

(23) For the structures of the different possible resorcinarene stereoisomers, see ref 3d.

g, 3.576 mmol) dissolved in DMSO (78 mL) at 100 °C for 28 h leads to the formation of several products. The complex ^1H NMR of the crude mixture reveals the presence of resorcinol as the predominant (90%) product as well as minor conversion to 2,4-dihydroxyacetophenone **8** (ratio of integrals of resorcinol triplet 6.94 ppm to **8** doublet at 7.76 ppm is 153:1, CH_3OD) and very small traces of xanthene **6a** (d, 7.65 ppm).



The production of resorcinol and **8** is consistent with the reversible opening and fragmentation of the resorcinarenes in acidic media.¹⁹ This result also complements our recent report describing the production of 4-formylphenylboronic acid from **3c**.²⁶ Furthermore, in acidic media, the addition of water at the methine carbon of **4** ($\text{R} = \text{Ar}$, $n = 0$, $m = 0$) followed by elimination has been described as an intermediate step in the synthesis of xanthenes.²⁷

We attain better conversion to xanthene **6a** from **3b** by limiting thermolysis time to 2 h. The ^1H NMR spectrum ($\text{DMSO-}d_6$) of the crude reaction mixture clearly shows a doublet at 7.65 ppm characteristic of **6a** with improved S/N as compared to the 28 h experiment (vide supra). Resonances centered at 5.26, 6.49, and 6.60 ppm are also discernible, overlaying with the ^1H NMR of an analytical sample²⁷ of **6a**. Because oxidation to xanthenes can be promoted by peroxides and acid,^{27,28} we should attain better conversion to xanthenes upon addition of these latter reagents. Indeed, we find that heating a solution of **3b** (50 mg, 0.203 mmol), H_2SO_4 (0.15 mL), and $\text{K}_2\text{S}_2\text{O}_8$ (1.0 mg) in 1.5 mL of MeOH at reflux for 2 h produces the most significant conversion (4% yield) of **3b** to **6a** we have observed to date.

We have shown that xanthenes form in solutions containing resorcinarene macrocycles (Scheme 1). The O_2 -induced radical decomposition of DMSO leads to in situ strong acid formation. The acid catalyzes a reverse condensation reaction to afford acyclic oligomers. The acyclic oligomers undergo oxidation also via the action of acid and peroxide.

Solution Color Changes Promoted by Mono- and Oligosaccharides in the Presence of Resorcinol Condensation Products. The linear maltodextrins **9a–f** are a prototypical class of readily available neutral oligosaccharides composed of glucose residues with α -1,4-glycoside linkages. We heat solutions (9:1 DMSO:H₂O, 1 mL) containing specific maltodextrins **9a–f** (2 mg) in the presence of **2a** (5 mg, 5.2 mM) until a temperature of 120 °C is reached in 3.0 min. In these studies, the resorcinarene is initially preheated alone for 15 min in 9:1





Figure 5. Colored solutions of **2a** after heating alone or with maltodextrins **9a–f**.

DMSO:H₂O at reflux prior to sugar addition to maximize chromophore formation.




Upon cooling the respective solutions containing mixtures of **2a** and each of **9a–f**, we observe solution colors of deep orange-yellow, yellow-gold, pale orange, orange, pale pink, and peach for maltose **9a** through maltoheptaose **9f**, respectively (Figure 5). The UV-vis spectra of each of the sugar-containing solutions exhibit increased absorbances at 460 nm (and decreased absorbances at 530 nm) as compared to a solution of **2a** heated without sugar (Figure 6). Heated solutions containing **1** and **9a–f** also exhibit absorbance maxima at 460 nm that are greater than that of **1** heated alone. The boronic acid-containing resorcinol condensation products thus allow for potentially useful colorimetric detection of larger oligosaccharides at 460 nm.

In analogous experiments to those above, but employing **2b** instead of **1** or **2a**, we find the absorbance changes in the oligosaccharide-containing solutions are greatly diminished as compared to the solution containing **2b** alone. The presence of boronic acid moieties is apparently a key to attaining enhanced

- (24) In the previous work (ref 19), acyclic oligomeric products (**3b** and two stereoisomeric trimeric compounds, three resorcinol rings, **4**, $\text{R} = \text{Me}$, $m = 1$, $n = 0$, Scheme 1) were isolated and characterized. Higher order acyclic oligomers (e.g., pentamers and hexamers) were also observed as major reaction products. Methyl ^1H NMR resonances, appearing as several doublets between 0.7 and 2.0 ppm (CH_3OD) that corresponded to neither **3b**, **4** ($\text{R} = \text{Me}$, $m = 1$, $n = 0$), nor resorcinarene macrocycles, thus were assigned to acyclics with five or more resorcinol moieties.
- (25) Flash column chromatography and TLC analysis of the thermolysis products of **2b** are complicated by the multiple product formation and fraction streaking.
- (26) Fronczek, F. R.; St. Luce, N. N.; Strongin, R. M. *Acta Crystallogr.* **2001**, C57, 1423.
- (27) Sen, R. N.; Sarkar, N. N. *J. Am. Chem. Soc.* **1925**, 47, 1079.
- (28) Deno, N. C.; Booker, E. L.; Kramer, K. E.; Saines, G. *J. Am. Chem. Soc.* **1969**, 91, 5237.

absorbance values with larger oligosaccharides. This result is in accord with our prior studies involving the comparison of monosaccharide-induced solution color changes in solutions containing **1**, **2b**, and **3a**.⁷

Sugar Solution Color Changes in the Presence of Resorcinol Condensation Products not Possessing Boronic Acids. To understand the different optical responses to sugars observed with boronic acid- and non-boronic acid-containing compounds, we first examine solutions containing **2b** and reducing and nonreducing sugars. We find that nonreducing sugars do not promote significant optical changes when heated in the presence of **2b**. For example, when we heat a solution of **2b** (5.2 mM) in 9:1 DMSO:H₂O (1 mL) in the presence of nonreducing sugars, either methyl- α -glucopyranoside (**10**, 15.6 mM) or α -methyl-mannopyranoside (**11**, 15.6 mM), for 2.5 min to gentle reflux, each solution color and absorption spectrum is unchanged from the heated solution of **2b** without sugar.



Reducing sugars undergo fragmentation to chain-shortened carbohydrates including aldonic acids, in the presence of peroxide.²⁹ The ¹H NMR spectra of respective DMSO-*d*₆ solutions containing three reducing sugars (28 mM), maltose (**9a**), D-(−)-fructose, and D-(+)-glucose, heated until gentle reflux (2.5 min), exhibit at least five new aldehyde resonances between 9 and 10 ppm as well as a broad proton resonance in the carboxylic acid region at 11.7 ppm.³⁰ None of these observed resonances is present in the NMR spectrum of heated neat DMSO-*d*₆. The region between 9 and 15 ppm in the spectra of the two nonreducing sugars **10** and **11** exhibits only very weak resonances which are also observed in the spectrum of heated neat DMSO-*d*₆. We thus hypothesize that the color changes observed in **2b** solutions are due to the new products formed upon reducing sugar thermolysis. When unheated solutions of either maltose (**9a**), D-fructose, or D-glucose are added to a colored solution of **2b** at room temperature, no color change is observed.

If reducing sugar reaction products are a cause of the solution color changes in the presence of the resorcinol condensation products, then the addition of preheated reducing sugars to preheated colored solutions of **2b** should produce a color change. We heat DMSO (0.9 mL) solutions of maltose (**9a**), D-fructose, and D-glucose each to a gentle reflux, cool to room temperature, and add 0.1 mL of H₂O. We add each of the solutions to three

preheated 9:1 DMSO:H₂O solutions of **2b** (5.2 mM). The three sugar (15.6 mM) solutions exhibit color changes observable by visual inspection and exhibit UV–vis spectra that are different from that of a heated solution containing only **2b**.³¹

We propose therefore that acyclic, xanthene-containing oligomers, which we have demonstrated to be the active chromophores in resorcinarene solutions, function as colorimetric receptors for sugar-derived acids. To prove this hypothesis, we chose to study the interaction of **6b** with *N*-acetylneuraminic acid (the most naturally abundant sialic acid, **12**), an important cell surface residue, in buffered media. Compound **6b** is a xanthene dye with hydroxyls adjacent to the C-6-OH and carbonyl moieties. The structure of **6b** is analogous to oxidized acyclic oligomers derived from resorcinarenes, in that it contains proximal potential hydrogen-bonding sites for charged carboxylates (Figure 7).

Addition of varying amounts of sialic acid **12** (1.9×10^{-4} – 1.5×10^{-3} M) to a peach-colored solution of compound **6b** (1.0×10^{-5} M) in a 9:1 DMSO:H₂O solution (HEPES buffer, 5.0×10^{-3} M final concentration, adjusted to pH = 7.5 in H₂O before mixing with DMSO) at room temperature results in a solution color change to yellow. The color changes monitored by UV–vis spectroscopy are concentration dependent (Figure 8). We calculate an apparent equilibrium constant, $K_{\text{exp}}(\text{app}) = 271.3 \text{ M}^{-1}$, with a reproducibility of $\pm 6\%$ (triplicate runs) for the binding of **6b** to **12**. We observe a color change to pink under the same conditions when NaOAc (2.2×10^{-4} M) instead of **12** is added to a solution of **6b**. This latter result suggests that the carboxylate moiety of the sugar is responsible for the change in solution color.³²

IR absorption shifts serve as evidence supporting a hydrogen-bonding interaction between the carboxylate moiety of **12** and the hydroxyls of **6b**. The FT-IR spectrum of a 1.0×10^{-2} M solution of **12** in DMSO exhibits a carbonyl absorption at 1726 cm^{−1}. Upon addition of an equimolar amount of **6b**, a new absorption appears at 1661 cm^{−1} accompanied by significant weakening of the absorption at 1726 cm^{−1}. This is in keeping with hydrogen bonding of **6b** to the sialic acid carbonyl oxygen.³³ Further support for hydrogen bonding is the fact that

(29) Arts, S. J. H. F.; Mombarg, E. J. M.; van Bekkum, H.; Sheldon, R. A. *Synthesis* **1997**, 597.

(30) Analysis of NMR integral ratios reveals that 59% of starting D-fructose is converted to new products under these conditions. The new product mixture contains 27% aldehydes and 0.9% carboxylic acids.

(31) We heat solutions of D-(−)-fructose, **9a**, or D-(+)-glucose in DMSO to a gentle reflux for 2 min and cool to room temperature. A solution of **2b** in DMSO is also heated to gentle reflux and cooled to room temperature. The three sugar solutions are added to the solution of **2b**, respectively, followed by the addition of water to afford mixtures in 9:1 DMSO:H₂O. The final concentration is 5.2×10^{-3} M for **2b** and 1.6×10^{-2} M for all sugars. Each solution is stirred for 6 min prior to UV–vis analysis. The absorbance decrease (as compared to the solution of **2b** without sugar) at 530 nm for the solution containing D-(+)-glucose is $(17.9 \pm 1.0)\%$, $(23.7 \pm 1.3)\%$ for D-(−)-fructose, and $(9.1 \pm 2.0)\%$ for **9a** (three trials).

(32) Having shown that acyclic aldonic acids, rather than uncharged sugars, produce a solution color change at room temperature in the presence of our active, non-boronic acid-containing chromophores, we next chose to generalize this finding to include sialic acid (**12**) because it is a prototypical, readily available, and bioactive cyclic sugar molecule possessing a carboxylate moiety. In addition, in our first communication in this study (ref 6), we used boron-containing **1** to produce significant solution color changes with **12** as well as other carboxylate-, phosphate-, and amine-functionalized saccharides. The results reported herein thus show that the boronic acid moieties may not be required (see also ref 7) to produce solution color changes in the presence of charged sugars. The fact that acetate also induces a color change at room temperature in buffered media demonstrates that simpler carboxylic acids could also be determined under these conditions. Research is in progress in our laboratory to develop selective sensing conditions for bioactive charged phosphates and carboxylates, including sialic acids, on the basis of these findings.

(33) The IR absorption changes we observe are in keeping with those reported by Zhang and co-workers who have recently studied the formation of hydrogen bonds between anthracene carboxylic acid and xanthene dyes using several techniques including IR and ¹H NMR: Zhang, H.; Zhang, M.; Shen, T. *Dyes Pigm.* **1999**, *43*, 15.






Figure 6. UV-vis spectra of solutions of **2a**, **1**, and **2b** heated alone or in the presence of **9a–f**. The boronic acid-containing receptors **1** and **2a** exhibit larger absorbance responses to oligosaccharides.

solution color changes upon sialic acid addition to **6b** are no longer observed when the proportion of H₂O to DMSO is $\geq 50\%$.³⁴

Aldonic acids, formed from reducing sugars in situ upon heating in DMSO, thus promote solution color changes via a charged hydrogen-bonding interaction with the xantheno chromophore. This latter interaction perturbs the ionization state of the C-6 hydroxyl of the dye. In contrast, heating reducing sugars in the presence of boronic acids should disfavor aldonic acid formation since anomeric hydroxyls are typically those involved in boronate ester formation. We therefore propose that the

(34) In addition, **6b** (10 mM) in DMSO-*d*₆ was titrated with 0.13–2.6 equiv of **12**. The ¹H NMR spectrum shows that a **6b** hydroxyl resonance exhibits a successive downfield chemical shift from 10.53 to 10.70 ppm upon increasing the concentration of **12**.




Figure 7. Charged hydrogen bonding between carbohydrates and proximal hydroxyls of **5** and **6b**.

heightened optical responses exhibited in colored solutions derived from boronic acid-containing resorcinol condensation products (Figure 6) are due, in large part, to sugar boronate formation.




Figure 8. UV-vis spectra of **6b** (1.0×10^{-5} M) in the absence and presence of **12** (1.9×10^{-4} – 1.5×10^{-3} M) in 9:1 DMSO:H₂O (pH 7.5 HEPES buffer). Inset shows the concentration of **12**.

Sugar Solution Color Changes in the Presence of Resorcinol Condensation Products Possessing Boronic Acids. It is known that boronic acid-appended dyes can produce color changes in the presence of saccharides.¹⁰ Importantly, when saccharides form cyclic boronates, the Lewis acidity of boron is enhanced.³⁵ Thus, upon saccharide binding, *sp*²-hybridized neutral boron is more readily converted to an *sp*³-hybridized anionic boronate via the addition of H₂O or HO[−] as the fourth ligand. It was shown by several researchers that the spectral changes of boronic acid-appended chromophores and fluorophores induced by saccharide binding arise due to the change from a neutral, *sp*² boronic acid to an *sp*³-hybridized anionic boronate-saccharide complex.^{10,35b} We thus set out to determine whether the formation of *sp*³-hybridized sugar boronates occurs under our experimental conditions.

In the presence of **2a** (40 mM), D-fructose-2-¹³C (1 equiv) in 9:1 DMSO-*d*₆:D₂O exhibits several new ¹³C-2 resonances which correspond to cyclic sugar boronic esters (Figure 9). The ¹³C chemical shifts (Table 2) are in agreement with the values obtained by Norrild and Eggert for the analogous *p*-tolylboronic acid sugar complexes.³⁶ Importantly, we observe resonances for

(35) (a) Lorand, J. P.; Edwards, J. D. *J. Org. Chem.* **1959**, *24*, 769. (b) Yoon, J.; Czarnik, A. W. *J. Am. Chem. Soc.* **1992**, *114*, 5874.

(36) Norrild, J. C.; Eggert, H. *J. Chem. Soc., Perkin Trans. 2* **1996**, 2583.




Figure 9. Structures of the four **2a**: β -D-fructofuranose complexes (**13**–**16**) observed in 9:1 DMSO- d_6 :D₂O. Boronates **15** and **16** are “exo” and “endo” isomers exhibiting ¹³C-2 chemical shifts of equal intensity at 114.4 and 114.3 ppm, respectively.

three anionic complexes all corresponding to known β -D-fructofuranose esters (Figure 9).

Further proof that we are forming anionic sugar boronates derives from ¹¹B NMR spectroscopy. The ¹¹B NMR chemical shifts of boronates change as a result of complexation with

Table 2.

fructose mutarotational isomer bound to boron	¹³ C-2 chemical shift (ppm)	
	literature value ^c (<i>p</i> -tolylboronic ester)	experimental value ^d (2a boronic ester)
β -D-fructofuranose	113.4 ^a	113.8 (14)
	114.7 ^a	114.4 (15)
	114.6 ^a	114.3 (16)
	115.2 ^a	115.5 (13)
β -D-fructopyranose	105.0 ^b	105.0
	107.6 ^b	107.7
α -D-fructofuranose	99.8 ^b	99.9
α -D-fructopyranose	92.6 ^b	93.2

^a In D₂O at pD = 11–12. ^b In DMSO- d_6 . ^c Reference 36. ^d Experimental values are obtained in 9:1 DMSO- d_6 :D₂O.

sugars due to differential electronic shielding of the ¹¹B atom. An upfield shift of the ¹¹B NMR signal accompanies the conversion of sp²-hybridized neutral species into sp³-hybridized boronate anions.³⁷ At pH = 6.5, compound **2a** (10 mM) in 1:1 DMSO:H₂O (pH value refers to the buffered aqueous portion before mixing) exhibits a single broad resonance at –19.1 ppm which we assign to the neutral sp²-hybridized boronic acid (**17**, Figure 10). At pH = 11.0, but in the presence of 0.5 equiv of D-fructose, a new resonance appears at –32.9 ppm which intensifies when the amount of D-fructose is increased to 5 equiv. We thus assign the resonance at –32.9 ppm to D-fructose cyclic boronate anion **20**. A solution of **2a** (20 mM) in DMSO also exhibits a resonance at –32.9 ppm upon D-fructose (3 equiv) addition. The observation of the resonance at –32.9 ppm





Figure 10. Equilibria for solutions of **2a** and D-fructose.

corresponding to boronate **20** in DMSO is consistent with our ¹³C NMR results.

The formation of the sugar boronate anion allows us to establish a mechanism for the sugar-induced color changes with our receptors. Anionic boronate formation, favored in the presence of sugars, leads to the diminished acidity of the C-6 hydroxyl. One way to envision this is via examination of xanthene resonance forms **21** and **22**. Structure **22** possesses a more stable cation than does **21**, rendering the C-6 hydroxyl of **22** relatively less ionizable.



We have reported that colored aqueous DMSO solutions of resorcinol condensation products exhibited enhanced absorbances at 532 nm upon raising solution pH.⁷ If sugar complexation results in a change of receptor pK_a, we should observe a change in receptor absorbance in pH-controlled media in the presence of sugars. Figure 11 shows the results of a pH versus absorbance titration monitored at 534 nm. As expected, the absorbance of colored 9:1 DMSO:H₂O solutions containing **2a** exhibits a sharp intensity onset beginning at ca. pH = 9. When the same pH titration experiment is run in the presence of D-fructose, the absorbance increase at and beyond pH = 9 is significantly lessened. This is evidence of the diminished ionizability of the C-6 hydroxyl induced by sugar complexation.

A similar pH titration profile is observed for colored solutions of **3a** alone and with added D-fructose (Figure 11), demonstrating that this finding appears general to these materials. Solutions containing **2a** or **3a** each exhibit color changes from dark brown to pale orange when D-fructose is added beyond pH = 9 (of the aqueous portion of the solution mixture). We are thus able to observe monosaccharide-induced solution color changes at room temperature in buffered media using these methods.³⁸

Oligosaccharides should also promote solution color changes via boronate ester formation with chromophore-appended bo-

(37) Nagai, Y.; Kobayashi, K.; Toi, H.; Aoyama, Y. *Bull Chem. Soc. Jpn.* **1993**, *66*, 2965.




Figure 11. Spectrophotometric study ($\lambda = 534$ nm) of colored solutions containing **2a** (top, 3.5×10^{-4} M) and **3a** (bottom, 3.5×10^{-4} M) alone and with D-fructose (3.5×10^{-2} M) in buffered 9:1 DMSO:H₂O (pH values refer to the H₂O portion before mixing with DMSO).

ronic acids. Prior studies (using fluorescence detection) showed that binding constants observed for the interaction of oligosaccharides and fluorophore-appended arylboronic acids increased with larger oligosaccharide size.³⁷ This has been attributed to sugar binding via boronate formation involving the hydroxyls of the oligosaccharide reducing termini. The binding is enhanced by secondary sugar-dye contacts including hydrophobic and CH- π interactions. Larger oligosaccharides have greater flexibility and can thus more readily fold back and interact with the dye molecule when a boronate ester is formed.

The linear maltodextrins **9a–f** ($0–7.5 \times 10^{-2}$ M) are dissolved in 0.5 mL of carbonate buffer (2.5×10^{-2} M, pH = 11) and mixed with 0.5 mL of DMSO colored preheated solutions of **2a** (5.0×10^{-4} M), respectively, to furnish a 1:1 DMSO:H₂O solvent system. The solution colors change from orange to yellow upon sugar addition. We thus detect neutral oligosaccharides in buffered media at room temperature by visual inspection. The apparent equilibrium constant ($K_{\text{exp}}(\text{app})$) is determined by UV-vis spectroscopy. The expected trend³⁷ of enhanced binding affinity correlating with longer oligosaccharide length is observed (Figure 12). We conclude that the

(38) Derivative analysis reveals slope maxima for the titration curves of colored solutions of **2a** and **3a** occur at pH values corresponding to 9.45 and 9.80, respectively. However, these values have no physical significance towards assigning a respective pK_a to the materials since the pH values on the x-axis of Figure 11 refer only to the aqueous sugar-containing solutions prior to mixing with solutions derived from heated **2a** and **3a** in DMSO. A known method for the determination of resorcinarene pK_a values in a mixed aqueous solvent system (Schneider, H.-J.; Güttes, D.; Schneider, U. *J. Am. Chem. Soc.* **1988**, *110*, 6449) more appropriate for potentiometric titrations was applied to colored solutions of **2a** and **3** with and without added fructose. Unfortunately, solubility problems precluded us from obtaining pK_a data via this method. Although to date we have not been able to determine pK_a values, the strikingly similar effect of fructose on the UV-vis spectra of distinct but related materials (colored solutions of **2a** and **3a**) shows that the sugar-induced perturbation of the chromophore ionization state is apparently significant and reproducible within this class of materials.




Figure 12. Plot of binding constant (three trials, monitored at 460 nm) vs molecular weight. Increasing molecular weight beginning with α -D-glucose through the series of linear maltodextrins (**9a–f**) results in an increase of binding constants (65.0, 108.3, 145.0, 189.7, 246.0, and 275.0 M^{-1}) in colored solutions containing **2a**. The binding constants are reproducible within $\pm 10\%$.

significant solution color changes observed for the oligosaccharides (**9a–f**) are due, in large part, to the formation of cyclic sugar boronate esters.

The acyclic oligomeric xanthenes existing in colored resorcinarene solutions function as receptors. They produce solution color changes in the presence of saccharides due to factors including cyclic boronate formation and hydrogen bonding. Each of these latter binding motifs can perturb the ionization state of the C-6 hydroxyl, the key moiety responsible for modulating the color of xanthene dyes.¹⁷ Acid-containing analytes, whether added directly or derived via in situ formation from the oxidation of reducing sugars, produce color changes via hydrogen bonding. Boronate ester formation, however, is the predominant factor leading to solution color changes with neutral sugars. The important problem of diminished optical response correlating with increasing oligosaccharide chain length^{13,14} may be overcome via the use of boronic acid-containing chromophores. This is apparently due to the enhanced binding affinities of larger oligosaccharides for boronic acids.³⁷

Conclusion

We have presented strong evidence that the color observed in Baeyer's initial resorcinarene condensation reaction² was due to the presence of xanthenes. In other words, he had created fluorescein¹ analogues at apparently low levels. Interestingly, Fischer, who won the Nobel Prize for sugar research, presented his doctoral thesis to Baeyer in 1874 on fluorescein and orcinol dyes.³⁹

We have shown that the colored products existing in solutions of resorcinarene macrocycles can serve as colorimetric indicators. The main findings of the present study include (i) the determination of the origin and structure of the active chromophores, (ii) elucidation of mechanisms associated with the solution color changes induced by sugars, and (iii) optical signaling for larger oligosaccharides.

The investigation of the colorimetric and fluorimetric properties of resorcinarenes, xanthenes, and related chromophoric materials is ongoing in our laboratory. Our efforts are now guided by the mechanistic studies described herein. Xanthene dyes containing well-positioned boronic acid or related binding

(39) Fischer, E. Ph.D. Dissertation, Strasbourg University, Strasbourg, France, 1874.

moieties should find application as powerful receptors for saccharides and other polar analytes such as carboxylates and phosphates.

The resorcinarenes, however, do offer potential advantages as compared to functionalized dye materials. A great attraction is their ease of synthesis in one step on ca. 200 g scale.^{3d} Having addressed many of the main mechanistic issues associated with the colorimetric sugar detection process, we are now also focusing on the study and optimization of important applied sensing parameters such as detection selectivity, sensitivity, and reversibility in aqueous and biological media. We are additionally developing new automated detection methods and expanding the detection capabilities of our receptors to include various analytes of biomedical significance.

Experimental Section

General. Matrix assisted laser desorption ionization mass spectra were acquired using a Bruker Proflex III MALDI mass spectrometer with either anthracene or dithranol matrixes. FT-IR spectra were recorded at room temperature on a Perkin-Elmer 1760X FT-IR spectrophotometer. UV–visible spectra were recorded at room temperature on a Spectramax Plus (Molecular Devices). Analytical thin-layer chromatography (TLC) was performed using general purpose silica gel on glass (Scientific Adsorbants). Flash chromatography columns were prepared with silica gel (Scientific Adsorbants, 32–63 μm particle size, 60 \AA). Analytic and preparative-scale HPLC were performed on a CM4000 multiple solvent delivery system (Milton Roy) and a Spectromonitor 5000 photodiode array detector (LDC Analytical) using a Dynamax 60 \AA C18 (21.4 mm ID \times 25 cm L) with a flow rate of 5 mL/min and a gradient of 50% water/MeOH to 100% MeOH in 20 min unless otherwise stated. The following compounds were prepared according to literature methods: **1**,⁵ **2a**,⁵ **2b**,⁴⁰ **3a**,⁷ **3b**,¹⁹ and **6a**.¹⁶ Isotopically labeled d-fructose-2-¹³C was purchased from Isotec. All other chemicals were purchased from Sigma or Aldrich and used without further purification. Proton NMR spectra were acquired in either CD₃OD, CH₃OD, or DMSO-*d*₆ on a Bruker DPX-250, DPX-400, or AMX-500 spectrometer. All δ values are reported with (CH₃)₄Si at 0.00 ppm or DMSO at 2.45 ppm as references.

Boron NMR Spectra. ¹¹B ($I = 3/2$) NMR data were collected at room temperature using a Chemagnetics Infinity solid-state spectrometer at a frequency of 128.35 MHz. All experiments were performed static (nonspinning), in the liquid-state, using a Chemagnetics 5 mm MAS probe and quartz NMR tubes. Data were recorded (64 k data points) following a 1.5 μs pulse (45° flip angle; aqueous boric acid) over a spectral width of 1.0 MHz. Each spectrum is the sum of 50 000 transients. Spin–lattice relaxation time (T_1) was not measured, but no saturation was observed using a 1.0 s pulse delay. All spectra were externally referenced using aqueous boric acid by setting the single ¹¹B resonance to –18.8 ppm.⁴¹ Experiments were first attempted on several commercial high-resolution, solution-state spectrometers from both major vendors with no success. We attribute this lack of success to a combination of factors including (a) high ¹¹B background signals from the probe, even with quartz tubes, (b) long receiver dead times as compared to the perceived short transverse relaxation times (T_2) of the sample, and (c) insufficient transmitter power required to uniformly irradiate large spectral windows to detect spectral components with a line width as large as 10 kHz.

X-ray Crystallographic Data. Intensity data were collected on a Nonius Kappa CCD diffractometer equipped with Mo K α radiation

and a graphite monochromator. The sample was cooled to 120 K by an Oxford Cryosystems Cryostream chiller. Data collection parameters and crystallographic data are provided in Supporting Information. Absorption and decay effects were negligible. The structure was solved by direct methods, using SIR97,⁴² and refined using SHELXL97.⁴³ H atoms were observed in difference maps, but were constrained to be in idealized positions in the refinement. OH hydrogen atoms are all disordered into two sites, all of which were treated as half populated. O–H distances were constrained to be 0.84 \AA , but otherwise, these H positions were refined.

pH Titration Experiments. Compounds **2a** or **3a** (3.9×10^{-4} M) in DMSO (stock solutions) are heated to gentle reflux and cooled to room temperature. Additionally, 6.3 mg of D-fructose is dissolved in 0.1 mL of H₂O containing buffer (2.5×10^{-1} M) and mixed with 0.9 mL of a receptor stock solution aliquot. The final concentrations: **2a** or **3a** = 3.5×10^{-4} M, D-fructose = 3.5×10^{-2} M, buffer = 2.5×10^{-2} M. For the pH titration experiments, buffers of pH from 4.5 to 5.0 were prepared in H₂O containing AcONa and titrated with HCl. Buffers of pH from 5.5 to 8.5 were prepared in H₂O containing KH₂PO₄ and titrated with NaOH. Buffers with higher pH were prepared in H₂O containing NaHCO₃ titrated with NaOH. For the study of sialic acid binding to **6b** in H₂O at pH 7.5, HEPES dissolved in H₂O was titrated with NaOH.

Binding Constant Determinations. The colored solutions of **6b** or **2a** in 9:1 DMSO:H₂O each exhibit a linear dependence of absorbance on concentration in the ranges measured: 3.9×10^{-6} – 3.1×10^{-5} M (**6b**) and 5.0×10^{-4} – 1.0×10^{-2} M (**2a**). Apparent equilibrium constant ($K_{\text{exp}}(\text{app})$)³⁷ data for the interaction of sugars and receptors derive from UV–vis spectroscopy. The equilibrium constants are evaluated via Benesi–Hildebrand treatment⁴⁴ (eqs 1 and 2) using UV–vis absorbance changes observed at 460 nm for the linear maltodextrin series in colored solutions of **2a** (complexation coverage = 50–60%). A 1:1 stoichiometry is observed for the interaction of **12** and **6b** by the continuous variation method (Job plot)⁴⁴ at $[\mathbf{6b}] + [\mathbf{12}] = 1.0 \times 10^{-5}$ M (constant). At 530 nm, **12** and **6b** solutions exhibit a complexation coverage = 25%. In each case, the double reciprocal plots according to eq 1 afforded very good linearity with a correlation coefficient $r \geq 0.99$.

$$\frac{1}{[A_0] - [A]} = \frac{1}{[A_0] - [A_\infty]} + \frac{1}{[A_0] - [A_\infty]} \cdot \frac{1}{K} \cdot \frac{1}{[\text{sugar}]} \quad (1)$$

$$K = \frac{\text{y-intercept}}{\text{slope}} \quad (2)$$

A_0 and A are UV–vis absorbances at 460 nm of chemosensor in the absence and presence of a sugar, respectively, while A_∞ is that of the complex of sugar and chemosensor.

Acknowledgment. We gratefully acknowledge the National Institutes of Health (GM-61915), the Arnold and Mabel Beckman Foundation through the Beckman Young Investigator Award Program, and the donors of the Petroleum Research Fund administered by the American Chemical Society (36497-AC4) for support of this research.

Supporting Information Available: Data collection parameters and crystallographic data for compound **7** (PDF/CIF). This material is available free of charge via the Internet at <http://pubs.acs.org>.

JA017713H

- (40) (a) Niederl, J. B.; Vogel, H. *J. Am. Chem. Soc.* **1940**, *62*, 2512. (b) Erdtmann, H.; Hoegberg, S.; Abramsson, S.; Nilsson, B. *Tetrahedron Lett.* **1968**, 1679. (c) Nilsson, B. *Acta Chem. Scand.* **1968**, 22732. (d) Cram, D. J.; Karbach, S.; Kim, H.-E.; Knobler, C. B.; Maverick, E. F.; Ericson, J. L.; Helgeson, R. C. *J. Am. Chem. Soc.* **1988**, *110*, 2229.
- (41) Onak, T. P.; Landesman, H.; Williams, R. E.; Shapiro, I. *J. Phys. Chem.* **1959**, *63*, 1533.

- (42) Altomare, A.; Burla, M. C.; Camalli, M.; Cascarano, G. L.; Giacovazzo, C.; Guagliardi, A.; Moliterni, A. G. G.; Polidori, G.; Spagna, R. *J. Appl. Crystallogr.* **1999**, *32*, 115.
- (43) Sheldrick, G. SHELXL97, University of Göttingen, Germany, 1997.
- (44) Benesi, H.; Hildebrand, J. H. *J. Am. Chem. Soc.* **1949**, *71*, 2703.