

Intramolecular [4 + 2] Cycloadditions of Iminoacetonitriles: A New Class of Azadienophiles for Hetero Diels–Alder Reactions

David T. Amos, Adam R. Renslo, and Rick L. Danheiser*

Department of Chemistry, Massachusetts Institute of Technology, Cambridge, Massachusetts 02139

Received February 12, 2003; E-mail: danheiser@mit.edu

Activated imines which can function as reactive 2π components in cycloadditions are valuable building blocks for the construction of nitrogen heterocycles.¹ In this communication we report on the preparation and intramolecular Diels–Alder reactions of *iminoacetonitriles*, a class of electron-deficient imines² whose cycloaddition chemistry has not previously been examined.³ Our interest in this class of imines derives from the expectation that they should function as reactive partners in a variety of cycloaddition and annulation processes, providing access to cyclic α -amino nitriles of diverse ring size (Scheme 1). α -Amino nitriles are exceptionally versatile intermediates for the synthesis of nitrogen heterocycles.⁴ Metalation provides opportunities for alkylation and other carbon–carbon bond-forming processes, while exposure to Lewis acids furnishes iminium ions which can be intercepted with Grignard reagents (Bruylants reaction) and organosilanes, or engaged in Mannich reactions and other useful “cation- π ”-type cyclization processes.

Previously, the synthesis of iminoacetonitriles has been achieved by the chlorination of α -amino nitriles followed by elimination of HCl with base.² In our hands this approach indeed proved workable, but for the preparation of our cycloaddition substrates we have developed a more expeditious route that begins with readily available alcohols⁵ and employs a Mitsunobu coupling reaction⁶ with the previously unknown, easily prepared reagent, $\text{HN}(\text{Tf})\text{CH}_2\text{CN}$.⁷ Base-promoted elimination of trifluoromethanesulfonate⁸ then furnishes the desired iminoacetonitriles as a mixture of *E* and *Z* isomers, both of which undergo cycloaddition to afford the same product(s). As summarized in Table 1, this protocol allows the efficient conversion of a variety of functionalized alcohols to the desired cycloaddition substrates in excellent overall yield.

Table 2 delineates the scope of the intramolecular iminoacetonitrile hetero Diels–Alder reaction. Most cycloadditions proceed smoothly at 85–120 °C in the presence of BHT (as radical inhibitor), and the *E*- and *Z*-imines appear to react at similar rates.^{9,10} Interestingly, in most cases the cycloadduct with *exo*-oriented cyano group is obtained as the major or exclusive product of the reaction. Our observations suggest that the initially formed epimeric cycloadducts equilibrate to afford the axial cyano isomer which is favored as a consequence of the “ α -amino nitrile anomeric effect.”¹¹

The α -amino nitrile moieties incorporated in these cycloadducts constitute *latent iminium ions*, which upon exposure to mild protic or Lewis acids are unmasked, setting the stage for further useful synthetic transformations. For example, reductive decyanation with NaBH_3CN excises the cyano group and furnishes unadorned quinolizidines (e.g., **21**) in excellent yield. Especially valuable is the application of the Bruylants reaction for further carbon–carbon bond construction, as illustrated with the conversion of cycloadduct **20** to the α -alkynylamine **22** (Scheme 2).

Of particular significance are the stereocomplementary transformations outlined in Scheme 3. Alkylation of the metalated nitrile

Scheme 1




Table 1. Synthesis of Iminoacetonitriles

entry	alcohol	triflamide yield (%) ^c	iminoacetonitrile	yield (%) ^c (<i>E/Z</i> ratio)
1		6 87–94		87–90 (80:20)
2		7 93		64–75 (71:29)
3		8 81–91		80–89 (77:23)
4		9 81–87		89–90 ^d (80:20)
5		10 81–94		86–93 (77:23)

SiR₃ = Si*t*-BuMe₂

^a 1.05 equiv $\text{HN}(\text{Tf})\text{CH}_2\text{CN}$, 1.2 equiv Ph_3P , 1.2 equiv DEAD, 1:1 THF–toluene, rt, 0.5–4 h (22 h for entry 2). ^b 3–4 equiv Cs_2CO_3 , THF, 45–55 °C, 2–4 h. ^c Isolated yield of products purified by column chromatography. ^d Enone **9** was converted to the dienol silyl ether (*t*-BuMe₂SiCl, NaI, Et₃N, CH₃CN, rt, 16 h, 96–97% yield) prior to elimination with Cs_2CO_3 .



with ethyl iodide followed by reductive decyanation furnishes exclusively the *endo*-ethyl product **23**, while Bruylants reaction with

Table 2. [4 + 2] Cycloadditions of Iminoacetonitriles^a

entry	iminoacetonitrile	cycloadduct	yield (%) ^b
1			67-70
2			61-64
3			44-45 ^c
4			71-78
5			79-87


SiR₃ = Si*t*-BuMe₂

^a Cycloadditions were run in toluene (0.05 M) with 3 equiv of BHT in a resealable threaded Pyrex tube at 120 °C for 15–36 h (CH₃CN, reflux for entry 2). ^b Isolated yield of products purified by column chromatography. ^c Obtained as a 79:21 mixture of isomers at C-1.

Scheme 2**Scheme 3**

EtMgBr leads predominantly to the diastereomeric quinolidine **24** via *exo* addition to the less sterically encumbered face of the intermediate iminium ion.

The availability of substituted quinolidines such as **23** and **24** via this strategy is noteworthy, since systems of this type are not available via cycloadditions of iminium ions according to the method of Grieco.¹² Thus, while the formiminium ion derivative of amine **25** undergoes smooth cycloaddition in water at 65 °C (eq 1), analogous reaction of **25** with propanal fails to deliver any of the corresponding α -substituted product (R¹ = Et).¹³



Further studies on the application of iminoacetonitriles in organic synthesis are underway and will be reported in due course.

Acknowledgment. We thank the National Institute of Health (GM 28273), Pharmacia, and the Merck Research Laboratories for generous financial support. D.T.A. was supported in part by NIH Training Grant CA 09112.

Supporting Information Available: Experimental procedures and characterization for key compounds (PDF). This material is available free of charge via the Internet at <http://pubs.acs.org>.

References

- (1) (a) Tietze, L. F.; Ketschau, G. *Top. Curr. Chem.* **1997**, *189*, 1. (b) Boger, D. L.; Weinreb, S. M. *Hetero Diels–Alder Methodology in Organic Synthesis*; Academic: San Diego, 1987.
- (2) (a) Boyer, J. H.; Dabek, H. *J. Chem. Soc., Chem. Commun.* **1970**, 1204. (b) Boyer, J. H.; Kooi, J. *J. Am. Chem. Soc.* **1976**, *98*, 1099. (c) De Kimpe, N.; Verhé, R.; De Buyck, J. C.; Schamp, N. *Synthesis* **1978**, 895.
- (3) For examples of oximinomalonitrile cycloadditions, see: (a) Biehler, J.-M.; Fleury, J.-P. *J. Heterocycl. Chem.* **1971**, *8*, 431. (b) Fleury, J.-P.; Desbois, M.; See, J. *Bull. Soc. Chim. Fr.* **1978**, II-147. (c) Bland, D. C.; Raudenbush, B. C.; Weinreb, S. M. *Org. Lett.* **2000**, *2*, 4007.
- (4) (a) Enders, D.; Shilvock, J. P. *Chem. Soc. Rev.* **2000**, *29*, 359. (b) Rubiralta, M.; Giralt, E.; Diez, A. In *Piperidine: Structure, Preparation, Reactivity, and Synthetic Applications of Piperidine and its Derivatives*; Elsevier: Amsterdam, 1991; pp 225–312. (c) Husson, H.-P.; Royer, J. *Chem. Soc. Rev.* **1999**, *28*, 383.
- (5) Alcohols **1–5** are available in two to four steps. For **1**, see Margot, C.; Rizzolio, M.; Schlosser, M. *Tetrahedron* **1990**, *46*, 2411. For **4**, see Ihara, M.; Taniguchi, T.; Yamada, M.; Tokunaga, Y.; Fukumoto, K. *Tetrahedron Lett.* **1995**, *36*, 8071. Preparation of alcohols **2**, **3**, and **5** will be reported in the full paper.
- (6) (a) Hughes, D. L. *Org. React.* **1992**, *42*, 335. (b) Hughes, D. L. *Org. Prep. Proced. Int.* **1996**, *28*, 127.
- (7) Prepared using the general method of Hendrickson, J. B.; Bair, K. W.; Bergeron, R.; Giga, A.; Skipper, P. L.; Sternbach, D. D.; Wareing, J. A. *Org. Prep. Proced. Int.* **1977**, *9*, 173.
- (8) For a related elimination, see: Hendrickson, J. B.; Bergeron, R.; Giga, A.; Sternbach, D. *J. Am. Chem. Soc.* **1973**, *95*, 3412.
- (9) Cycloadditions of substrates with three-atom tethers (leading to indolizidine products) also proceed in good yield, albeit more slowly; details will be discussed in the full paper.
- (10) Control experiments indicate that the *E*- and *Z*-imines are in equilibrium under the conditions of the reaction.
- (11) Bonin, M.; Romero, J. R.; Grierson, D. S.; Husson, H.-P. *J. Org. Chem.* **1984**, *49*, 2392.
- (12) (a) Larsen, S. D.; Grieco, P. A. *J. Am. Chem. Soc.* **1985**, *107*, 1768. (b) Grieco, P. A.; Parker, D. T. *J. Org. Chem.* **1988**, *53*, 3325.
- (13) Grieco similarly reports that only iminium ions derived from formaldehyde participate in efficient aza Diels–Alder reactions in aqueous solution.^{12a}

JA0346290