


## Rhodium-Catalyzed Disulfide Exchange Reaction

Mieko Arisawa, and Masahiko Yamaguchi

*J. Am. Chem. Soc.*, **2003**, 125 (22), 6624-6625 • DOI: 10.1021/ja035221u • Publication Date (Web): 10 May 2003

Downloaded from <http://pubs.acs.org> on March 29, 2009



### More About This Article

Additional resources and features associated with this article are available within the HTML version:

- Supporting Information
- Access to high resolution figures
- Links to articles and content related to this article
- Copyright permission to reproduce figures and/or text from this article

[View the Full Text HTML](#)

## Rhodium-Catalyzed Disulfide Exchange Reaction

Mieko Arisawa and Masahiko Yamaguchi\*

Department of Organic Chemistry, Graduate School of Pharmaceutical Sciences, Tohoku University, Aoba, Sendai 980-8578 Japan

Received March 19, 2003; E-mail: yama@mail.pharm.tohoku.ac.jp

A disulfide bond plays an important role in the construction of the secondary and the tertiary structures of polypeptides and proteins. Industrially important polymers such as rubber contain polysulfide bonds.<sup>1</sup> The cleavage and recombination of organodisulfides and polysulfides, resulting in the disulfide exchange is, therefore, an important process for the production, modification, and degradation of such biologically active substances and materials. The cleavage of sulfur–sulfur bonds is known to occur either homolytically or heterolytically: (a) short-lived sulfonyl radicals are generated by homolytic scission via photolysis,<sup>2</sup> oxidation,<sup>3</sup> or heating;<sup>4</sup> and (b) ionic scission generates mercaptides (XS<sup>−</sup>) under basic/nucleophilic conditions<sup>5</sup> or sulfenium cation (XS<sup>+</sup>) under acidic/electrophilic conditions.<sup>6,7</sup> The sulfur radical, anion, or cation then reacts with another disulfide, resulting in exchange reactions. These reactions, however, often require harsh reaction conditions such as high temperature and strong acids or bases, and are generally slow in solution, particularly in the case of alkyl disulfides. We considered that the disulfide exchange process, which proceeds rapidly under neutral conditions and at ambient temperature, would provide a new methodology, which has broad applications to biosciences and material sciences. Described here is a novel exchange reaction of disulfides catalyzed by a transition metal complex.<sup>8</sup> A rapid and mild disulfide exchange reaction takes place in the presence of RhH(PPh<sub>3</sub>)<sub>4</sub>, a phosphine, and trifluoromethanesulfonic acid, which can be used for the peptide disulfide exchange. In addition, the metal system catalyzes the disproportionation of diselenide or ditelluride with disulfide.

When bis(2-benzoyloxyethyl) disulfide **1** (0.5 mmol) was treated with dibutyl disulfide **2** (0.5 mmol) in the presence of RhH(PPh<sub>3</sub>)<sub>4</sub> (1.5 mol %), tris(*p*-tolyl)phosphine (6 mol %), and trifluoromethanesulfonic acid (3 mol %) in refluxing acetone for 15 min, 2-benzoyloxyethyl butyl disulfide **3** (0.49 mmol, 49% yield based on the BzO(CH<sub>2</sub>)<sub>2</sub>S group) was obtained with the recovery of **1** (0.21 mmol, 41%) and **2** (0.20 mmol, 40%), which is the statistical product ratio (Table 1, entry 1). When another portion of **2** (0.5 mmol) was added to the refluxing mixture, the yield of **3** increased to 71% after 10 min. Addition of **2** twice (1.0 and 1.0 mmol) at 10 min intervals further increased the yields to 86% and 92%, respectively. This catalyst system was found to be active after the equilibrium was attained. The exchange reaction of **1** and **2** in a 1:4 initial ratio gave **3** in 86% yield with recovery of **1** (12%) and **2** (76%) (entry 2). The rhodium complex, trifluoromethanesulfonic acid, and phosphine are essential; no reaction occurs in the absence of the rhodium complex (entry 3), and there was low conversion in the absence of the phosphine or acid (entries 4 and 5). The effect of added phosphine was examined by reacting for 3 min, and triarylphosphines, particularly those with electron-donating substituents were found to be effective (entries 6, 7, 8, and 9). Bidentate phosphines (entries 10 and 11) and tributylphosphine (entry 12) were less effective. This reaction was effectively promoted by trifluoromethanesulfonic acid, and the yield of **3** decreased with

**Table 1.** Rhodium-Catalyzed Exchange of Symmetrical Disulfides<sup>a</sup>


$(RS)_2 + (n-C_4H_9S)_2$ $\xrightarrow[\text{Acetone, refl., 15 min}]{\text{RhH(PPh}_3)_4 \text{ (3 mol\%)}, \text{CF}_3\text{SO}_3\text{H (6 mol\%)}, \text{p-tol}_3\text{P (12 mol\%)}}$		R-S-S-n-C <sub>4</sub> H <sub>9</sub>	
run	R	phosphine	yield/% <sup>b</sup>
1 <sup>c</sup>	BzO(CH <sub>2</sub> ) <sub>2</sub> S	<i>p</i> -tol <sub>3</sub> P	49
2		<i>p</i> -tol <sub>3</sub> P	86
3 <sup>c,d</sup>		<i>p</i> -tol <sub>3</sub> P	N.D. <sup>e</sup>
4 <sup>c</sup>		none	3
5 <sup>c,f</sup>		<i>p</i> -tol <sub>3</sub> P	8
6 <sup>c,g</sup>		( <i>p</i> -MeOC <sub>6</sub> H <sub>4</sub> ) <sub>3</sub> P	19
7 <sup>c,g</sup>		<i>p</i> -tol <sub>3</sub> P	24
8 <sup>c,g</sup>		PPh <sub>3</sub>	14
9 <sup>c,g</sup>		( <i>p</i> -ClC <sub>6</sub> H <sub>4</sub> ) <sub>3</sub> P	6
10 <sup>c,g</sup>		dppe <sup>h</sup>	N.D.
11 <sup>c,g</sup>		dppf <sup>i</sup>	10
12 <sup>c,g</sup>		Bu <sub>3</sub> P	5
13 <sup>j</sup>	<i>n</i> -C <sub>8</sub> H <sub>17</sub>	<i>p</i> -tol <sub>3</sub> P	77
14	PhCH <sub>2</sub>	<i>p</i> -tol <sub>3</sub> P	81
15	<i>t</i> -BuMe <sub>2</sub> SiO(CH <sub>2</sub> ) <sub>6</sub>	<i>p</i> -tol <sub>3</sub> P	84
16 <sup>k</sup>	Ph	<i>p</i> -tol <sub>3</sub> P	19
17 <sup>l</sup>		dppe	90
18 <sup>m</sup>	Ph	dppe	85

<sup>a</sup> See Supporting Information for reaction conditions. <sup>b</sup> Yield based on the RS group. <sup>c</sup> The reaction was conducted using **1** (0.5 mmol) and **2** (0.5 mmol) in the presence of RhH(PPh<sub>3</sub>)<sub>4</sub> (1.5 mol %), *p*-tol<sub>3</sub>P (6 mol %), and trifluoromethanesulfonic acid (3 mol %). <sup>d</sup> In the absence of Rh complex. <sup>e</sup> N.D.: not detected by <sup>1</sup>H NMR. <sup>f</sup> In the absence of CF<sub>3</sub>SO<sub>3</sub>H. <sup>g</sup> The reaction time: 3 min. <sup>h</sup> 1,2-Bis(diphenylphosphino)ethane. <sup>i</sup> 1,1'-Bis(diphenylphosphino)ferrocene. <sup>j</sup> The reaction was conducted using dioctyl disulfide (2.5 mmol) and **2** (12.5 mmol) in refluxing acetone for 30 min, and the product was isolated by distillation. <sup>k</sup> The reaction was conducted in acetone at room temperature for 5 min in the presence of RhH(PPh<sub>3</sub>)<sub>4</sub> (0.75 mol %) and *p*-tol<sub>3</sub>P (3 mol %). <sup>l</sup> The reaction was conducted in acetone at room temperature for 5 min in the presence of RhH(PPh<sub>3</sub>)<sub>4</sub> (0.75 mol %) and dppe (1.5 mol %). <sup>m</sup> The reaction was conducted using diphenyl disulfide **5** (1.0 mmol) and bis(*sec*-butyl) disulfide **4** (0.25 mmol) in refluxing acetone for 10 min in the presence of RhH(PPh<sub>3</sub>)<sub>4</sub> (1 mol %) and dppe (2 mol %).


methanesulfonic acid or *p*-toluenesulfonic acid. The rhodium-catalyzed exchange reactions of **2** (4 equiv) and several symmetrical dialkyl disulfides were rapid, completing within 15 min in refluxing acetone (entries 13, 14, and 15). Bis(*sec*-butyl) disulfide **4** and **2**, however, did not undergo the exchange reaction under these conditions. The reaction of a diaryl disulfide and a dialkyl disulfide took place more rapidly than that of dialkyl disulfides, when 1,2-bis(diphenylphosphino)ethane (dppe) was used instead of tris(*p*-tolyl)phosphine (entries 16, 17, and 18). The ligand effect is an interesting aspect of the transition metal-catalyzed exchange.

Since, the present disproportionation is considerably affected by the substituent on the disulfide, selective exchange of symmetrical disulfides can be conducted. Treatment of equimolar amounts (0.5 mmol) of **4**, diphenyl disulfide **5**, and di(*p*-tolyl) disulfide **6** gave phenyl *p*-tolyl disulfide **7** (0.49 mmol), **5** (0.25 mmol), and **6** (0.25 mmol) at room temperature for 1 min in the presence of 1 mol % of the rhodium complex and 2 mol % of dppe, with the unchanged


Scheme 1



Scheme 2



Scheme 3



**4** (0.48 mmol). Random disproportionation takes place giving statistical amounts of **7** (0.32 mmol), *sec*-butyl phenyl disulfide **8** (0.34 mmol), and *sec*-butyl *p*-tolyl disulfide **9** (0.34 mmol) at acetone reflux for 5 min with the recovery of **4** (0.14 mmol), **5** (0.16 mmol), and **6** (0.16 mmol). Such property may be used for the selective disulfide formation in polymeric compounds such as proteins.

A cysteine derivative **10** exchanged with **2** to give the unsymmetrical disulfide **11** without racemization (Scheme 1). It should be noted that a tripeptide glutathione derivative **12** also undergoes the reaction giving a disulfide **13** in 74% yield after recrystallization.

The exchange reactions of dialkyl disulfides and dialkyl diselenides are also catalyzed by the same complex (Scheme 2).<sup>9</sup> When dibutyl diselenide (0.25 mmol) was treated with **1** (1.5 mmol) in the presence of RhH(PPh<sub>3</sub>)<sub>4</sub> (5 mol %), tris(*p*-tolyl)phosphine (20 mol %), and trifluoromethanesulfonic acid (5 mol %) in refluxing acetone for 15 min, butylselenyl 2-benzoyloxyethyl sulfide was obtained (0.36 mmol, 72% yield based on the *n*-C<sub>4</sub>H<sub>9</sub>Se group). It was confirmed that the rhodium complex is essential for the reaction.

Exchange reaction of disulfides and ditellurides gives tellurino-sulfides (Scheme 3). No reaction takes place without the rhodium complex under the conditions.

We describe here a novel Rh-catalyzed exchange reaction of disulfides. Compared with the conventional exchange reaction, this

reaction takes place rapidly under mild conditions and is applicable to peptide disulfide exchange. In addition, the reaction can be controlled by changing the ligand or acid, and the application of this methodology to polysulfides including elemental sulfur is now under investigation. This is a novel combination of transition metal chemistry and organosulfur chemistry.

**Acknowledgment.** This work was supported by JSPS (Nos. 14207094 and 13771323). M.A. expresses her thanks for financial supports from New Energy and Industrial Technology Development Organization (NEDO) of Japan (No. 02A44003d), the Hayashi Memorial Foundation for Female Natural Scientists, the Fujisawa foundation, and the Sankio Chemical Award in Synthetic Organic Chemistry, Japan.

**Supporting Information Available:** Detailed experimental procedures, characterization data (PDF). This material is available free of charge via the Internet at <http://pubs.acs.org>.

## References

- Parker, A. J.; Kharasch, N. *Chem. Rev.* **1959**, *59*, 583. Also see the following for the metal-catalyzed S–S bond cleavage. Kuniyasu, H. In *Catalytic Heterofunctionalization*; Togni, A., Grutzmacher, H., Eds.; Wiley-VCH: Weinheim, 2001; p 271. Kondo, T.; Mitsudo, T. *Chem. Rev.* **2000**, *100*, 3205. Beletskaya, I.; Moberg, C. *Chem. Rev.* **1999**, *99*, 3435.
- Rosengren, K. *Acta Chem. Scand.* **1962**, *16*, 1401. Milligan, B.; Rivett, D. E.; Savage, W. E. *Aust. J. Chem.* **1963**, *16*, 1020. Sayamol, K.; Knight, A. R. *Can. J. Chem.* **1968**, *46*, 999.
- Nelander, B.; Sunner, S. *J. Am. Chem. Soc.* **1972**, *94*, 3576. Nagano, T.; Arakane, K.; Hirobe, M. *Tetrahedron Lett.* **1980**, 5021. Do, Q. T.; Elothmani, D.; Guillanton, G. L.; Simonet, J. *Tetrahedron Lett.* **1997**, *38*, 3383. Exchange reaction of alkyl disulfide in solution using NO–O<sub>2</sub> is reported. Itoh, T.; Tsutsumi, N.; Ohsawa, A. *Bioorg. Med. Chem. Lett.* **1999**, *9*, 2161.
- Leandri, G.; Tundo, A. *Ric. Sci.* **1953**, *23*, 1646; *Chem. Abstr.* **1954**, *48*, 12699t.
- McAllan, D. T.; Cullum, T. V.; Dean, R. A.; Fidler, F. A. *J. Am. Chem. Soc.* **1951**, *73*, 3627. Parker, A. J.; Kharasch, N. *J. Am. Chem. Soc.* **1960**, *82*, 3071. Haraldson, L.; Olander, C. J.; Sunner, S.; Varde, E. *Acta Chem. Scand.* **1960**, *14*, 1509. Dalman, G.; McDermed, J.; Gorin, G. *J. Org. Chem.* **1964**, *29*, 1480. Harpp, D. N.; Smith, R. A. *J. Am. Chem. Soc.* **1982**, *104*, 6045.
- Sanger, F. *Nature* **1953**, *171*, 1025. Kolthoff, I. M.; Stricks, W.; Kapoor, R. C. *J. Am. Chem. Soc.* **1955**, *77*, 4733. Benesch, R. E.; Benesch, R. *J. Am. Chem. Soc.* **1958**, *80*, 1666. Ryle, A. P.; Sanger, F.; Smith, L. F.; Kitai, R. *Biochem. J.* **1955**, *60*, 541. Kice, J. L.; Ekman, G. E. *J. Org. Chem.* **1975**, *40*, 711.
- Disulfide–thiol exchange reaction was reported. For example, Dalman, G.; McDermed, J.; Gorin, G. *J. Org. Chem.* **1964**, *29*, 1480. Eldjarn, L.; Pihl, A. *J. Am. Chem. Soc.* **1957**, *79*, 4589.
- Arisawa, M.; Yamaguchi, M. *J. Am. Chem. Soc.* **2000**, *122*, 2387. Arisawa, M.; Yamaguchi, M. *Adv. Synth. Cat.* **2001**, *343*, 27. Arisawa, M.; Yamaguchi, M. *Org. Lett.* **2001**, *3*, 311. Arisawa, M.; Yamaguchi, M. *Org. Lett.* **2001**, *3*, 763. Arisawa, M.; Momozuka, R.; Yamaguchi, M. *Chem. Lett.* **2002**, 272. Arisawa, M.; Suwa, A.; Fujimoto, K.; Yamaguchi, M. *Adv. Synth. Cat.* **2003**, *345*, 560.
- Examples of selenosulfide synthesis. Nakazaki, M. *J. Chem. Soc., Jpn., Pure Chem. Sect.* **1954**, 75, 338. Kostiner, E. S.; Reddy, M. N.; Urch, D. S.; Massay, A. G. *J. Organomet. Chem.* **1968**, *15*, 383. Sister, H. H.; Kotia, N. K. *J. Org. Chem.* **1971**, *36*, 1700. Guo, H.; Zhang, Y. *J. Chem. Res. (S)* **2000**, 374. Goto, K.; Nagahama, M.; Mizushima, T.; Shimada, K.; Kawashima, T.; Okazaki, R. *Org. Lett.* **2001**, *3*, 3569. Potapov, V. A.; Amosova, S. V.; Petrov, P. A.; Romanenko, L. S.; Keiko, V. V. *Sulfur Lett.* **1992**, *15*, 121.

JA035221U