


## Is It Possible To Estimate the Enantioselectivity of a Chiral Catalyst from Its Racemic Mixture?

Franz Lagasse, Masaki Tsukamoto, Christopher J. Welch, and Henri B. Kagan

*J. Am. Chem. Soc.*, **2003**, 125 (25), 7490-7491 • DOI: 10.1021/ja0300315 • Publication Date (Web): 31 May 2003

Downloaded from <http://pubs.acs.org> on March 29, 2009



### More About This Article

Additional resources and features associated with this article are available within the HTML version:

- Supporting Information
- Access to high resolution figures
- Links to articles and content related to this article
- Copyright permission to reproduce figures and/or text from this article

[View the Full Text HTML](#)



## Is It Possible To Estimate the Enantioselectivity of a Chiral Catalyst from Its Racemic Mixture?

Franz Lagasse,<sup>†</sup> Masaki Tsukamoto,<sup>†</sup> Christopher J. Welch,<sup>‡</sup> and Henri B. Kagan<sup>\*†</sup>

Laboratoire de Synthèse Asymétrique (UMR 8075), Institut de Chimie Moléculaire et des Matériaux d'Orsay, Université Paris-Sud, 91405 Orsay, France, and Process Research, Merck & Co., Inc., Rahway, New Jersey 07065

Received January 17, 2003; E-mail: kagan@icmo.u-psud.fr

In recent years, parallel synthesis and screening technologies have increasingly been applied to the discovery of chiral catalysts.<sup>1–3</sup> The evaluation of enantioselective catalysts in a parallel fashion is necessary to avoid the cancellation of opposing enantioselectivities that would result from combined screening. It would be extremely useful to directly measure the enantioselectivity of the component enantiomers of a racemic catalyst without first having to resolve or prepare the individual catalyst enantiomers.<sup>4–8</sup> In this Communication, we report and discuss our finding that the diastereoselectivity observed upon sequential transformation of an achiral substrate bearing two remote prochiral centers can sometimes be used to calculate catalyst enantioselectivity.

There are many reports of the reaction of an enantiopure reagent or catalyst with an achiral substrate bearing multiple prochiral centers.<sup>9–14</sup> Scheme 1 illustrates the simple case of a symmetrical diketone **1** with remote carbonyls that is quantitatively reduced in the presence of a chiral catalyst to afford a mixture of the three stereoisomers of diol **3**. For a reaction utilizing an enantiopure catalyst, it is easy to establish that if one defines  $ee_1$  as the enantioselectivity of the first step (formation of intermediate ketol, **2**) and  $de_2$  as the diastereoselectivities of the second step (assumed identical), then  $de_{\text{homo}}$ , the preference for formation of the homo-chiral versus meso diastereomer, is simply the product of  $ee_1$  and  $de_2$  (eq 1).<sup>15</sup> The observed diastereoselectivity,  $de_{\text{homo}}$ , is large when the stereoselectivities ( $ee_1$  and  $de_2$ ) are both high. In the special case where the stereoselectivities of the two steps are identical,  $de_{\text{homo}}$  is given simply as the square of  $ee_1$  (eq 2). Consequently, the easily measured  $de_{\text{homo}}$  can be used to calculate  $ee_1$ , the enantioselectivity of the first step.


$$ee_1 de_2 = de_{\text{homo}} \quad (1)$$

$$ee_1^2 = de_{\text{homo}} \quad (\text{if } ee_1 = de_2) \quad (2)$$


The situation becomes more complex for racemic catalysts, as the second reaction step may involve a catalyst molecule that is the enantiomer of the one used in the initial step. In such instances where substantial catalyst “scrambling” occurs,  $de_{\text{homo}}$  will be nil, whatever the  $ee_1$ . In the special case where the same enantiomer of the catalyst is involved in both steps, the above discussion shows that eqs 1 and 2 remain valid, and  $de_{\text{homo}} > 0$  will be observed. In other words, a racemic catalyst is expected to produce a low level of the meso product diastereomer whenever a single catalyst molecule with high enantioselectivity performs both steps on a bifunctional substrate. We report here on such a case involving a catalyst–substrate system in which the second reaction is fast relative to release of the initial product–catalyst complex.

To ensure that  $ee_1$  and  $de_2$  are the same, it is necessary to design a bifunctional substrate where the reactive centers are sufficiently

**Scheme 1.** Diagram of the Product Distribution in Two Consecutive Asymmetric Reactions with an Enantiopure Catalyst



separated so as to minimize substrate control in the second reaction. Otherwise, eq 2 becomes invalid, and one must use eq 1 where the desired information  $ee_1$  is obscured by the presence of  $de_2$  (a composite of catalyst and substrate control).<sup>15</sup> As a first check on the validity of the above ideas, we investigated diketone **1**, which afforded a nearly statistical 1:2:1 ratio of diol **3** stereoisomers (i.e., no diastereoselectivity) when reduced by a variety of achiral reagents.<sup>16–18</sup> Enantioselective borane reductions catalyzed by chiral oxazaborolidines (OABs) derived from 2,2-diphenylprolinol were initially considered. Catalysts such as **4–6** are often used in asymmetric catalysis.<sup>19</sup>



Ketone **7** was selected as a model for study of the enantioselectivity of the monoreduction of **1**. Its reduction to alcohol **8** by 0.75 equiv of  $BH_3 \cdot SME_2$  under various conditions in the presence of OAB catalysts gave enantioselectivities close to 90% ee. Reduction of diketone **1** by enantiopure catalysts **4–6** afforded the chiral diol **3** in very high ee (close to 99%). The diastereoselectivity  $de_{\text{homo}}$  is quite good, and the relationship between  $de_{\text{homo}}$  and  $ee_{\text{homo}}$  is accurately described by

$$ee_{\text{homo}} = 2(de_{\text{homo}})^{1/2}/(1 + de_{\text{homo}}) \quad (3)$$

established when  $ee_1 = de_2$ .<sup>15</sup>

We were pleased to observe that reduction of diketone **1** with the corresponding racemic catalysts **4–6** also afforded diol **3** with high diastereoselectivity ( $de_{\text{homo}}$ ).

As shown in Table 1, use of eq 2 to calculate  $ee_1$  on the basis of measured  $de_{\text{homo}}$  values gave results similar to those obtained by

<sup>†</sup> Université Paris-Sud.

<sup>‡</sup> Merck & Co., Inc.

**Table 1.** Diastereoselectivity and Enantioselectivity in the Reduction of Diketone **1** in THF<sup>a,b</sup>


catalyst	% de <sub>homo</sub> <sup>c</sup>	% ee <sub>1</sub> <sup>d</sup> (% ee of <b>8</b> ) <sup>c</sup>
( <i>R</i> )- <b>4</b> <sup>e</sup>	87	93 (82)
( <i>R</i> )- <b>4</b> <sup>f</sup>	86	93 (79)
<i>rac</i> - <b>4</b> <sup>e</sup>	56	75 (82)
<i>rac</i> - <b>4</b> <sup>f</sup>	86	93 (79)
( <i>S</i> )- <b>5</b> <sup>e</sup>	81	90 (87)
( <i>S</i> )- <b>5</b> <sup>f</sup>	72	85 (83)
<i>rac</i> - <b>5</b> <sup>f</sup>	83	91 (83)
( <i>R</i> )- <b>6</b> <sup>e</sup>	89	94 (–) <sup>g</sup>
( <i>R</i> )- <b>6</b> <sup>e</sup>	82	91 (–) <sup>g</sup>

<sup>a</sup> 10 mol % of catalyst unless otherwise stated. <sup>b</sup> Addition of a solution of **1** in 1 h to a solution of catalyst and BH<sub>3</sub>·SMe<sub>2</sub> (2.5 equiv). Reaction time: 1 h, unless otherwise stated. Quantitative conversion of **1**. More details in Supporting Information. <sup>c</sup> Measured by chiral SFC (Chiralcel OD-H). (*R*)-OAB catalysts give (*S,S*)-diol **3**. <sup>d</sup> Calculated from eq 2. <sup>e</sup> 0 °C. <sup>f</sup> 66 °C. <sup>g</sup> Not measured.

reduction of model ketone **7**. Comparable results were obtained using toluene as a reaction solvent (see Supporting Information).

Two additional catalytic systems were investigated, hydrosilylation ( $\alpha$ -naphthylphenylsilane) catalyzed by [RhCl(dip)(cod)]<sup>20</sup> and hydrogen transfer from 2-propanol catalyzed by a chiral ruthenium complex generated in situ from [{RuCl<sub>2</sub>(*p*-cymene)}]<sub>2</sub>, ephedrine, and KOH.<sup>21</sup> The enantio- and diastereoselectivities of these reactions were studied using the enantiopure catalysts and are in good agreement with eqs 3 and 2. However, no diastereoselectivity is observed with the racemic catalysts, suggesting that significant catalyst scrambling has occurred.

What then makes the OAB catalysts so special? To answer this question, it is helpful to consider the mechanism of the reaction.<sup>19</sup> It has been observed that two hydrogen atoms from BH<sub>3</sub> are used in ketone reductions, the second hydrogen being delivered in a faster reaction.<sup>22</sup> In the present case, the half-reduced species **9** contains an N-coordinated ROBH<sub>2</sub> moiety that can serve as an excellent hydride donor. Consequently, intramolecular hydride delivery to form **10** occurs rapidly, the large size of the chelation ring ensuring that the first stereocenter does not influence the stereochemical outcome of the second reduction.<sup>23</sup> Instead, product stereochemistry is controlled predominantly by the chiral oxazaborolidine system.



Additional support for this interpretation is provided by some experiments with (*S*)- or (*R*)-ketol **2** (98% ee), prepared by reduction of **1** by (+)- or (–)-Ipc<sub>2</sub>BBr, respectively. The reduction of (*S*)- or (*R*)-**2** by a stoichiometric amount of (*R*)-OAB **5** (in situ complexed by 1 mol equiv of borane)<sup>24</sup> gave a large amount of homochiral (*S,S*)-diol **3** (*S,S*:*meso* = 96:4) or *meso*-diol **3** (*meso*:*R,R*) = 93:7), respectively, clearly establishing that the stereochemical outcome of the second reduction step is under catalyst control, with the (*R*)-catalyst affording the (*S*)-stereocenter in both the matched and the mismatched cases.<sup>25</sup> This suggests that ee<sub>1</sub> and de<sub>2</sub> are indeed equivalent and that the use of eq 2 to calculate ee<sub>1</sub> is justified.

In conclusion, these results demonstrate that, at least in some specialized cases, it is possible to evaluate the enantioselectivity of racemic catalysts without the need for resolution. One can envision additional strategies to counteract the escape of the catalyst

before the second reaction has taken place. We are presently investigating these approaches in an effort to expand the generality of this method of using easily measured diastereoselectivity as an indicator of the enantioselectivity of the component enantiomers of a racemic catalyst.

**Acknowledgment.** We thank CNRS and Université Paris-Sud for their financial support. F.L. acknowledges the Servier Foundation and the Société des Amis des Sciences for fellowships. This work is dedicated to the memory of Professor Jean Jacques (Dec. 15, 1917–July 10, 2001).

**Supporting Information Available:** Experimental procedure for reduction of various diketones and ketones, results of OAB reductions of **1** in toluene, data concerning diols **3** and ketol **2**, equations and mathematical treatment, catalytic cycle and chromatograms of some ee measurements (PDF). This material is available free of charge via the Internet at <http://pubs.acs.org>.

## References

- (1) Kagan, H. B. *J. Organomet. Chem.* **1998**, *567*, 3–6.
- (2) Shimizu, K. D.; Snapper, M. L.; Hoveyda, H. *Chem.-Eur. J.* **1998**, *4*, 1885–1889.
- (3) (a) Reetz, M. T. *Angew. Chem., Int. Ed.* **2001**, *40*, 284–310. (b) Reetz, M. T. *Angew. Chem., Int. Ed.* **2002**, *41*, 1335–1338.
- (4) Some approaches rely on the addition of a suitable chiral additive.<sup>5–7</sup> The efficiency of a catalyst for kinetic resolution can be estimated from the racemic catalyst, as recently established.<sup>8</sup>
- (5) Brown, J. M.; Maddox, P. J. *Chirality* **1991**, *3*, 345–354.
- (6) Fallor, J. W.; Parr, J. J. *Am. Chem. Soc.* **1993**, *115*, 804–805.
- (7) Mikami, K.; Matsukawa, S. *Nature* **1997**, *385*, 613–617.
- (8) Dominguez, B.; Hodnett, N. S.; Lloyd-Jones, G. C. *Angew. Chem., Int. Ed.* **2001**, *40*, 4289–4291.
- (9) Rautenstrauch, V. *Bull. Soc. Chim. Fr.* **1994**, *131*, 515–524 and references therein.
- (10) El Baba, S.; Sartor, K.; Poulin, J.-C.; Kagan, H. B. *Bull. Soc. Chim. Fr.* **1994**, *131*, 525–533 and references therein.
- (11) Soai, K.; Hori, H.; Kawahara, M. *J. Chem. Soc., Chem. Commun.* **1992**, 106–108; *J. Chem. Soc., Chem. Commun.* **1992**, 1034.
- (12) Hoye, T. R.; Suhadolnik, J. C. *J. Am. Chem. Soc.* **1985**, *107*, 5312–5313.
- (13) Schreiber, S. L.; Schreiber, T. S.; Smith, D. B. *J. Am. Chem. Soc.* **1987**, *109*, 1525–1529.
- (14) Kuwano, R.; Sawamura, M.; Shirai, J.; Takahashi, M.; Ito, Y. *Bull. Chem. Soc. Jpn.* **2000**, *73*, 485–496.
- (15) ee<sub>1</sub>, de<sub>2</sub>, and de<sub>homo</sub> are taken to be between 0 and 1 for discussions here. Equations describing ee<sub>homo</sub> and the specialized case of two different stereoselectivities (de<sub>2</sub>, de<sub>3</sub>) in the second step are discussed in the Supporting Information.
- (16) Neudeck, H.; Schlögl, K. *Monatsh. Chem.* **1975**, *106*, 229–259.
- (17) The reduction of diketones PhCO(CH<sub>2</sub>)<sub>n</sub>COPh by LAH or BH<sub>3</sub>·SMe<sub>2</sub> showed that the diastereoselectivity is zero (racemic = meso) for *n* = 4–10. For *n* = 3, we found a product distribution of homo/*meso* 52/48 (LAH) and 56/44 (BH<sub>3</sub>·SMe<sub>2</sub> in toluene at 0 °C or 110 °C). The last value is in agreement with ref 18.
- (18) Quallich, G. J.; Keavey, K. N.; Woodall, T. M. *Tetrahedron Lett.* **1995**, *36*, 4729–4732.
- (19) Corey, E. J.; Helal, C. J. *Angew. Chem., Int. Ed.* **1998**, *37*, 1986–2012 and references therein.
- (20) Dumont, W.; Poulin, J.-C.; Dang, T.-P.; Kagan, H. B. *J. Am. Chem. Soc.* **1973**, *95*, 8295–8299.
- (21) Takehara, J.; Hashiguchi, S.; Fujii, A.; Inoue, S.-I.; Ikariya, T.; Noyori, R. *Chem. Commun.* **1996**, 233–234.
- (22) Mathre, D. J.; Thompson, A. S.; Douglas, A. W.; Hoogsteen, K.; Carroll, J. D.; Corley, E. G.; Grabowski, E. J. *J. Org. Chem.* **1993**, *58*, 2880–2888.
- (23) An achiral oxazaborolidine (10 mol % equiv prepared in situ from 2-aminoethanol and borane) used as a catalyst gives a low diastereoselectivity (% de<sub>homo</sub> = 28, THF, 66 °C), ruling out a strong substrate control in the second reduction (for details, see Supporting Information).
- (24) In the catalytic reaction of ketol (*R*)-**2** with (*S*)-OAB **5** in THF at 0 °C, the de<sub>homo</sub> is quite low (51%) because of the competitive direct formation of monoalkoxyborane, followed by uncatalyzed intramolecular reduction.
- (25) This was confirmed by the partial conversion (35%) of racemic ketol **2** in the presence of (*R*)-**5**, and BH<sub>3</sub> (THF, 0 °C). Ketol **2** was recovered (racemic), and a 1:1 mixture of (*S,S*)-**3** (92% ee) and *meso*-**3** was obtained.

JA0300315