


## Chiroptical Switching in a Bistable Molecular Shuttle

Giovanni Bottari, David A. Leigh, and Emilio M. Prez

*J. Am. Chem. Soc.*, **2003**, 125 (44), 13360-13361 • DOI: 10.1021/ja036665t • Publication Date (Web): 14 October 2003

Downloaded from <http://pubs.acs.org> on March 30, 2009



### More About This Article

Additional resources and features associated with this article are available within the HTML version:

- Supporting Information
- Links to the 23 articles that cite this article, as of the time of this article download
- Access to high resolution figures
- Links to articles and content related to this article
- Copyright permission to reproduce figures and/or text from this article

[View the Full Text HTML](#)

## Chiroptical Switching in a Bistable Molecular Shuttle

Giovanni Bottari, David A. Leigh,\* and Emilio M. Pérez


School of Chemistry, University of Edinburgh, The King's Buildings, West Mains Road, Edinburgh EH9 3JJ, U.K.

Received June 13, 2003; E-mail: david.leigh@ed.ac.uk


Although various methods for switching the positions of macrocycles in bistable rotaxane-based molecular shuttles have been developed,<sup>1</sup> exploiting such movements to trigger property changes has thus far received little attention.<sup>2,3</sup> Here we demonstrate one of the first examples of a property change achieved through a large amplitude translational motion in a rotaxane, a chiroptical switch in which light-induced shuttling of the macrocycle along the thread produces a strong induced circular dichroism (ICD) response when the macrocycle is hydrogen-bonded to a chiral peptide station.

Chiral dipeptides have previously been shown to induce an asymmetric response in the aromatic ring absorption bands of intrinsically achiral components of [2]rotaxanes through tight intercomponent binding in nonpolar solvents.<sup>4,5</sup> The effect can be "switched off" by adding a polar solvent (e.g., MeOH) to break the hydrogen-bonding interactions between macrocycle and thread. Although triggered changes in optical properties are currently utilized in optical data storage and processing, waveguides, and other photonics applications,<sup>6</sup> a solvent change is clearly unlikely to prove useful as a means of switching such real-world devices. However, the breaking (and making) of intercomponent interactions

### Scheme 1. Synthesis of Bistable Molecular Shuttles *E/Z-1*<sup>a</sup>



<sup>a</sup> Reaction conditions: i) *N*-Boc-L-leucine, 4-(dimethylamino)pyridine (4-DMAP), 1-(3-dimethylaminopropyl)-3-ethyl-carbodiimide hydrochloride (EDCI), CHCl<sub>3</sub>, 94%. ii) trifluoroacetic acid (TFA), CHCl<sub>3</sub>, quantitative. iii) *N*-Boc-glycine, 4-DMAP, EDCI, CHCl<sub>3</sub>, 91%. iv) TFA, CHCl<sub>3</sub>, quantitative. v) *N*-Boc-11-aminoundecanoic acid, 4-DMAP, EDCI, CHCl<sub>3</sub>, 82%. vi) TFA, CHCl<sub>3</sub>, quantitative. vii) fumaric acid monoethylester, 4-DMAP, EDCI, CHCl<sub>3</sub>, 85%. viii) EtOH, NaOH, 91%. ix) 4-DMAP, EDCI, DMF, 74%. x) *p*-xylylenediamine, isophthaloyl dichloride, Et<sub>3</sub>N, CHCl<sub>3</sub>, 58%. xi) EITHER 254 nm, CH<sub>2</sub>Cl<sub>2</sub>, 20 min, 56% OR 254 nm, CH<sub>3</sub>CN, 20 min, 49% OR 350 nm, benzophenone, CH<sub>2</sub>Cl<sub>2</sub>, 20 min, 70% OR 312 nm, CH<sub>2</sub>Cl<sub>2</sub> or CH<sub>3</sub>CN, 20 min, 62% OR 400–670 nm, cat. Br<sub>2</sub>, CH<sub>2</sub>Cl<sub>2</sub>, 2 min, >95% OR 130 °C, C<sub>2</sub>H<sub>2</sub>Cl<sub>4</sub>, 6 days, 95%.




**Figure 1.** <sup>1</sup>H NMR spectra (400 MHz in CDCl<sub>3</sub> at 298 K) of (a) fumaramide thread *E-2*, (b) fumaramide rotaxane *E-1*, (c) maleamide thread *Z-2*, (d) maleamide rotaxane *Z-1*. The assignments correspond to the lettering shown in Scheme 1.

is also a feature of positionally bistable molecular shuttles. Accordingly, it seemed possible that optical properties could be influenced solely by switching the position of a macrocycle in a molecular shuttle that incorporates a chiral peptide "station".

Such a bistable shuttle, *E/Z-1*, is shown in Scheme 1. The idea is that in *E-1* the macrocycle resides over the strongly macrocycle-binding fumaramide portion of the thread and the asymmetric center (*E-1*→*Z-1*), the ring moves to the glycy-L-leucine (Gly-Leu) unit, locking the molecule in a co-conformation where aromatic rings (principally those of the C-terminal stopper<sup>4</sup>) are held in a well-expressed chiral environment. The change in the position of the macrocycle should thus manifest itself in terms of a measurable change in the CD response.

Molecular shuttle *E-1* was synthesized from thread *E-2* in 58% yield (Scheme 1). Comparison of the <sup>1</sup>H NMR spectra of the



**Figure 2.** (a) CD spectra (0.1 mM in  $\text{CHCl}_3$ ) at 298 K of Z-1 (blue), E-1 (purple), E-2 (green), and Z-2 (red). (b) UV-vis spectra of E-1 (purple) and Z-1 (blue) in  $\text{CHCl}_3$  at 298 K,  $[\epsilon] = \text{mM}^{-1}\text{cm}^{-1}$ . Dotted line represents the Z-1/E-1 absorption ratio. (c) Percentage of E-1 in the photostationary state (from  $^1\text{H}$  NMR data, 400 MHz,  $\text{CD}_3\text{CN}$ , 298 K) after alternating irradiation at 254 nm (half integers) and 312 nm (integers) for five complete cycles. The right-hand Y axis shows the CD absorption at 246 nm.

E-rotaxane and thread (Figure 1, a and b) shows the excellent discrimination of the macrocycle toward the different stations. While the Gly-Leu protons are only slightly affected by the aromatic shielding effect of the macrocycle, the E-olefin protons are significantly shifted upfield ( $\sim 1.2$  ppm), confirming that the co-conformer having the macrocycle over the olefin station is the major translational isomer in E-1.

Light-induced E $\rightarrow$ Z isomerization<sup>7</sup> of the fumaramide unit reverses<sup>8</sup> the macrocycle-binding affinity of the two stations in the molecular shuttle because the cis-olefin (maleamide) is largely self-satisfying in terms of H-bonding and the amide carbonyl groups are no longer suitably orientated for binding to the macrocycle.<sup>7</sup> The  $^1\text{H}$  NMR spectra of Z-1 and Z-2 (Figure 1, c and d) reveal the concomitant change in the position of the macrocycle. The Z-olefin protons  $\text{H}_d$  and  $\text{H}_e$  occur at almost identical chemical shifts in rotaxane and thread, whereas the Gly methylene protons are now shielded by 0.9 ppm.

The shuttle design works remarkably well. When the macrocycle is tightly bound close to the Leu residue in Z-1, the aromatic rings of the rotaxane do, indeed, experience a well-expressed chiral environment as evidenced by CD spectroscopy. Of the two rotaxanes and two threads, only rotaxane Z-1 gives a CD response, which is both strong ( $-13\text{k deg cm}^2 \text{ dmol}^{-1}$ ) and, for the

L-enantiomer, negative (Figure 2a). The absence of any detectable signal for the E-rotaxane shows that the CD signal is genuinely only generated by controlling the position of the macrocycle in the shuttle.

The most efficient ways of interconverting the rotaxane diastereomers are photochemical: 350 nm, catalytic benzophenone sensitizer,  $\text{CH}_2\text{Cl}_2$ , 70% for E-1 $\rightarrow$ Z-1 and 400–670 nm, catalytic  $\text{Br}_2$ ,  $\text{CH}_2\text{Cl}_2$ , >95% for Z-1 $\rightarrow$ E-1 (Scheme 1). However, a modest difference in the electronic absorption spectra of E-1 and Z-1 (Figure 2b) results in different photostationary states for the isomers at 254 nm (56% Z-1 in  $\text{CH}_2\text{Cl}_2$ ; 49% Z-1 in  $\text{CH}_3\text{CN}$ ) and 312 nm (38% Z-1 in either  $\text{CH}_2\text{Cl}_2$  or  $\text{CH}_3\text{CN}$ ), meaning that even for this first-generation system a large net change ( $>1500 \text{ deg cm}^2 \text{ dmol}^{-1}$ ) in elliptical polarization can be achieved solely through irradiation with light of different wavelengths. The photoisomerizations are highly reproducible with five complete switching cycles (E-1 $\rightarrow$ Z-1 $\rightarrow$ E-1) carried out with no decomposition detectable by NMR or CD spectroscopy (Figure 2c).

In conclusion, we have demonstrated a system where a large amplitude displacement of a macrocycle along a thread elicits a chiral optical response. In addition to this being a novel mode of switching to be explored for possible photonic and data storage applications, control of the expression of chirality in switchable interlocked systems through hiding or revealing chiral subunits could find important applications in areas where chiral transmission from one chemical entity to another underpins a physical or chemical response, such as the seeding of supertwisted nematic liquid crystalline phases or asymmetric synthesis.

**Acknowledgment.** This work was supported by the European Union Future and Emerging Technology Program *MechMol* and the EPSRC. D.A.L. is an EPSRC Advanced Research Fellow (AF/982324).

**Supporting Information Available:** Experimental procedures and spectral data for all new compounds (PDF). This material is available free of charge via the Internet at <http://pubs.acs.org>.

## References

- (1) (a) Sauvage, J.-P.; Dietrich-Buchecker, C., Eds.; *Molecular Catenanes, Rotaxanes and Knots*; Wiley-VCH: Weinheim, 1999. (b) Balzani, V.; Credi, A.; Raymo, F. M.; Stoddart, J. F. *Angew. Chem., Int. Ed.* **2000**, *39*, 3348–3391.
- (2) For a recent example of a light-driven mechanical motion which elicits a chiral response, see: Muraoka, T.; Kinbara, K.; Kobayashi, Y.; Aida, T. *J. Am. Chem. Soc.* **2003**, *125*, 5612–5613.
- (3) For systems where a large amplitude translational motion in a rotaxane architecture brings about a property change, see: (a) size/shape variation in a molecular “muscle”: Jimenez-Molero, M. C.; Dietrich-Buchecker, C.; Sauvage, J. P. *Chem. Eur. J.* **2002**, *8*, 1456–1466. (b) changes in conductivity: Collier, C. P.; Wong, E. W.; Bělohorský, M.; Raymo, F. M.; Stoddart, J. F.; Kuekes, P. J.; Williams, R. S.; Heath, J. R. *Science* **1999**, *285*, 391–394. (c) color variation: Bissell, R. A.; Córdova, E.; Kaifer, A. E.; Stoddart, J. F. *Nature* **1994**, *369*, 133–137.
- (4) Asakawa, M.; Brancato, G.; Fanti, M.; Leigh, D. A.; Shimizu, T.; Slawin, A. M. Z.; Wong, J. K. Y.; Zerbetto, F.; Zhang, S. *J. Am. Chem. Soc.* **2002**, *124*, 2939–2950.
- (5) For an alternative chiral amino acid-containing rotaxane system see: (a) Smukste, I.; Smithrud, D. B. *J. Org. Chem.* **2003**, *68*, 2547–2558. (b) Smukste, I.; House, B. E.; Smithrud, D. B. *J. Org. Chem.* **2003**, *68*, 2559–2571.
- (6) (a) Irie, M., Ed. Photochromism: Memories and Switches (Special Issue). *Chem. Rev.* **2000**, *100*, 1638–1890. (b) Feringa, B. L., Ed. *Molecular Switches*; Wiley-VCH: Weinheim, 2001.
- (7) Gatti, F. G.; León, S.; Wong, J. K. Y.; Bottari, G.; Altieri, A.; Farran Morales, A. M.; Teat, S. J.; Frochot, C.; Leigh, D. A.; Brouwer, A. M.; Zerbetto, F. *Proc. Natl. Acad. Sci. U.S.A.* **2003**, *100*, 10–14.
- (8) Altieri, A.; Bottari, G.; Dehez, F.; Leigh, D. A.; Wong, J. K. Y.; Zerbetto, F. *Angew. Chem., Int. Ed.* **2003**, *42*, 2296–2300.

JA036665T