


Communication

Heterocyclic Peptide Backbone Modifications in an α -Helical Coiled Coil

W. Seth Horne, Maneesh K. Yadav, C. David Stout, and M. Reza Ghadiri

J. Am. Chem. Soc., **2004**, 126 (47), 15366-15367 • DOI: 10.1021/ja0450408 • Publication Date (Web): 05 November 2004

Downloaded from <http://pubs.acs.org> on April 5, 2009



More About This Article

Additional resources and features associated with this article are available within the HTML version:

- Supporting Information
- Links to the 10 articles that cite this article, as of the time of this article download
- Access to high resolution figures
- Links to articles and content related to this article
- Copyright permission to reproduce figures and/or text from this article

[View the Full Text HTML](#)



Heterocyclic Peptide Backbone Modifications in an α -Helical Coiled Coil

W. Seth Horne, Maneesh K. Yadav, C. David Stout, and M. Reza Ghadiri*

Departments of Chemistry, Molecular Biology, and The Skaggs Institute for Chemical Biology, The Scripps Research Institute, 10550 North Torrey Pines Road, La Jolla, California 92037

Received August 17, 2004; E-mail: ghadiri@scripps.edu

Among a number of elegant biomimetic approaches that have been developed for the modification of the peptide backbone,¹ only a few have been applied to protein structures.² Here, we report high-resolution structural consequences of amide backbone replacements in the context of a folded peptide architecture. We show that a triazole ϵ^2 -amino acid (Figure 1a)³ can be used as a dipeptide surrogate in α -helical coiled coils and report on the effects of the substitution on the thermodynamic stability, the helical secondary structure, and the four helix bundle quaternary organization.

The reactivity and functional group tolerance of Cu(I)-catalyzed 1,3-dipolar cycloaddition between azides and alkynes to give 1,2,3-triazoles⁴ have led to the increasing use of this reaction in bioconjugate chemistry.⁵ We reported recently the use of triazole ϵ^2 -amino acids as a dipeptide replacement in the design of self-assembling peptide nanotubes.⁶ We hypothesized that the structural and functional features of the triazole ϵ^2 -amino acids could also be potentially useful in the context peptide and protein secondary structures. The backbone of the ϵ -amino acid is one atom longer compared to that of a native dipeptide (Figure 1a), leading to a calculated increase in C_α - C_α spacing of 1.1 Å.⁷ In addition to the amide NH and carbonyl groups flanking the residue, the triazole ring possesses two nitrogen atoms (N² and N³) that might act as hydrogen bond acceptors. Furthermore, the triazole ring has a large dipole that could align with that of the other amides in a given peptide secondary structure.⁸

We selected the pLI mutant of the α -helical coiled coil GCN4 in order to test the utility of the ϵ^2 -amino acid substitution in the context of a peptide with well-defined secondary and quaternary structure in solution and in the solid state.⁹ In coiled coils, interhelical interactions between buried hydrophobic residues as well as exposed hydrophilic side chains lead to robust peptide self-assembly into α -helical bundles. We replaced dipeptides K₈L₉, K₁₅L₁₆, and E₂₂L₂₃ in the pLI-GCN4 sequence with an L-leucine-derived triazole ϵ -amino acid to give sequences **1**, **2**, and **3**, respectively (Figure 1b). In each case, the isobutyl side chain of the ϵ -residue was predicted to replace a leucine residue in the hydrophobic core of the native four-helix bundle. The amino acid employed was synthesized in two steps from L-leucine and used in standard solid-phase peptide synthesis conditions.

Circular dichroism (CD) spectra of peptides **1–3** (50 μ M in 10 mM MOPS, pH 7.0) showed minima at 208 and 222 nm, characteristic of an α -helical secondary structure (Figure S2 in the Supporting Information). Thermal denaturation, monitoring the molar ellipticity at 222 nm, indicated broad two-state transitions for peptides **2** and **3**, while peptide **1**, similar to the parent pLI-GCN4 sequence, was considerably more stable and did not fully denature up to 96 °C.⁹ Gel permeation chromatography indicated that peptides **1** and **3** adopt tetrameric oligomerization states in solution, while peptide **2** appears to exist primarily as a dimeric assembly (Table 1). These studies indicate that the modified peptides retain much of the native α -helical character, but that the position




Figure 1. (a) Native dipeptide and the L-leucine-derived triazole ϵ^2 -amino acid incorporated as a replacement. (b) Sequences for pLI-GCN4 and modified peptides **1–3**; X denotes incorporation of the ϵ^2 -residue.

Table 1. Biophysical Data and PDB IDs for Peptides **1–3**^a

peptide	$[\theta]_{222}$ (deg cm ² dmol ⁻¹) ^b	T_m (°C)	N_{agg} ^c	PDB ID
pLI-GCN4 ^d	-30600	>96	4	1GCL, 1UO2
1	-28000	>96	4	1U9G
2	-13100	36	2	1U9F
3	-21000	61	4	1U9H

^a See the Supporting Information for experimental details and full spectra. ^b At 4 °C. ^c Apparent aggregation state in solution as determined by gel permeation chromatography. ^d Values and structure 1GCL are from ref 9.




Figure 2. Schematic representation of the crystal structure of peptides **2** (a) and **3** (b) with atomic positions shown for the triazole residues. Each four-helix bundle superposes on a crystallographic 2-fold axis, and unique chains in each structure are indicated by different colors.¹⁰

of the ϵ^2 -amino acid substitution differentially influences thermodynamic stability.

We employed X-ray crystallography to ascertain the structural consequences of the ϵ^2 -amino acid substitutions in the context of the helical coiled coil architecture. Crystal structures were obtained for peptides **1–3** at 2.2 Å resolution. Although each peptide crystallized in a different unit cell and space group, all three exhibited the parallel tetrameric coiled coil structure of the parent




Figure 3. (a) Detail from the crystal structure of **2** showing participation of the triazole residue in main-chain hydrogen bonding; residues and nitrogen atoms of the triazole ring are numbered. (b) Top-down view of the crystal structure of **3** showing the hydrophobic plate formed by the triazole ϵ^2 -residues and interchain hydrogen bonds bridged by water. (c) Detail of two chains from the crystal structure of **3** showing the interhelical crossing. Dashed lines indicate hydrogen bonds. In panels a and c, H atoms are modeled based on N, C, and O coordinates.¹⁰

pLI-GCN4 sequence with a crystallographic 2-fold symmetry axis along the center of each bundle (Figure 2 and Figure S3 of the Supporting Information). In peptides **2** and **3**, a complete well-resolved structure similar to that of the parent pLI-GCN4 was obtained. However, in the case of **1**, the electron density for the portion of the peptide preceding the triazole ϵ -amino substitution (residues 1–8) was not observed, presumably due to chain disorder in the crystal (Figure S3 of the Supporting Information).

The crystal structure of **2** is similar to that of the parent pLI, with the hydrophobic side chain of the ϵ -residue projecting toward the core of the bundle (Figure S4 of the Supporting Information). Notably, the ϵ -residue in each chain fully participates in α -helical backbone hydrogen bonding (Figure 3a). The N² of the triazole accepts a hydrogen bond from the amide NH of Ile₁₈, and the triazole C⁵–H appears to participate in a CH–O hydrogen bond with the carbonyl oxygen of Ile₁₂.¹¹ This observation is supported by the geometry and 2.2 Å distance between the atoms and is in agreement with the large dipole of the triazole ring.⁸ Hence, the normal $i, i + 4$ hydrogen bonding that would have been provided to residues 12 and 18 in the parent pLI structure is replaced by the triazole ring. In addition, the amide NH of the ϵ -residue is hydrogen bonded to the carbonyl oxygen of Ile₁₂, while the N₆ of the imidazole ring of His₁₇ is hydrogen bonded to the carbonyl of Ser₁₄. These structural features give rise to an increase in the α -helical pitch of about 1.8 Å in the region of the ϵ -residue. This increase in the local pitch creates a shallow pocket adjacent to the triazole that is occupied by a water molecule providing bridging hydrogen bonds that may contribute to the stability of the helical structure (Figure S5 of the Supporting Information).

The ϵ^2 -amino acid substitution in peptide **3** creates an unusual right-handed interhelical crossover structure such that the helix formed by residues 1–22 of one chain is completed by residues 23–32 of another (Figure 3c and Figure S6 of the Supporting Information). Residues 20–24, which include the ϵ^2 -amino acid at position 22, act together as a template for the strand crossing, providing an overall helical chain register similar to that of the parent pLI fold. Interchain interactions consisting of a backbone hydrogen bond between the carbonyl oxygen of Glu₂₀ and the amide NH of Ala₂₃ and a water-bridged hydrogen bond between N³ of the triazole ring and the amide NH of Arg₂₄ further stabilize this intertwined fold (Figure 3c). The isobutyl side chains of the ϵ^2 -residue also form a core hydrophobic packing in the helix bundle similar to that in the parent pLI (Figure 3b).

In summary, we have shown that a non-natural 1,2,3-triazole ϵ^2 -amino acid can replace a dipeptide in an α -helical secondary structure. In light of these observations, we suggest that Cu(I)-catalyzed azide–alkyne coupling could be useful in the non-native chemical synthesis of peptides and proteins.¹²

Acknowledgment. We thank the National Institute of General Medical Sciences (GM57690) for financial support, and NSF for a fellowship to W.S.H.

Supporting Information Available: Figures S1–S6, experimental details, crystallographic data, and CD spectra. This material is available free of charge via the Internet at <http://pubs.acs.org>.

References

- (1) For reviews on the subject, see: (a) Spatola, A. F. *Chemistry and Biochemistry of Amino Acids, Peptides, and Proteins*; Dekker: New York, 1983; pp 267–357. (b) Ahn, J.-M.; Boyle, N. A.; MacDonald, M. T.; Janda, K. D. *Mini-Rev. Med. Chem.* **2002**, *2*, 463–473.
- (2) (a) Chapman, E.; Thorson, J. S.; Schultz, P. G. *J. Am. Chem. Soc.* **1997**, *119*, 7151–7152. (b) Belgere, G. S.; Dawson, P. E. *J. Am. Chem. Soc.* **2000**, *122*, 12079–12082. (c) Wales, T. E.; Fitzgerald, M. C. *J. Am. Chem. Soc.* **2001**, *123*, 7709–7710. (d) Deechongkit, S.; Nguyen, H.; Powers, E. T.; Dawson, P. E.; Gruebele, M.; Kelly, J. W. *Nature* **2004**, *430*, 101–105.
- (3) For a description of the ϵ^2 nomenclature, see Figure S1 of the Supporting Information.
- (4) (a) Rostovtsev, V. V.; Green, L. G.; Fokin, V. V.; Sharpless, K. B. *Angew. Chem., Int. Ed.* **2002**, *41*, 2596–2599. (b) Tornøe, C. W.; Christensen, C.; Meldal, M. *J. Org. Chem.* **2002**, *67*, 3057–3064.
- (5) (a) Wang, Q.; Chan, T. R.; Hilgraf, R.; Fokin, V. V.; Sharpless, K. B.; Finn, M. G. *J. Am. Chem. Soc.* **2003**, *125*, 3192–3193. (b) Link, A. J.; Tirrell, D. A. *J. Am. Chem. Soc.* **2003**, *125*, 11164–11165. (c) Deiters, A.; Cropp, T. A.; Mukherji, M.; Chin, J. W.; Anderson, J. C.; Schultz, P. G. *J. Am. Chem. Soc.* **2003**, *125*, 11782–11783. (d) Fazio, F.; Bryan, M. C.; Blixt, O.; Paulson, J. C.; Wong, C.-H. *J. Am. Chem. Soc.* **2002**, *124*, 14397–14402. (e) Speers, A. E.; Adam, G. C.; Cravatt, B. F. *J. Am. Chem. Soc.* **2003**, *125*, 4686–4687.
- (6) Horne, W. S.; Stout, C. D.; Ghadiri, M. R. *J. Am. Chem. Soc.* **2003**, *125*, 9372–9376.
- (7) Due to the planar aromatic ring structure of the triazole, this distance is fixed regardless of the backbone torsion angles.
- (8) Palmer, M. H.; Findlay, R. H.; Gaskell, A. J. *J. Chem. Soc., Perkin Trans. 2* **1974**, *4*, 420–428.
- (9) Harbury, P. B.; Zhang, T.; Kim, P. S.; Alber, T. *Science* **1993**, *262*, 1401–1407.
- (10) Figure prepared using PyMOL: DeLano, W. L. *The PyMOL Molecular Graphics System*; <http://www.pymol.org>.
- (11) For a review on CH–O hydrogen bonds, see: Derewenda, Z. S.; Lee, L.; Derewenda, U. *J. Mol. Biol.* **1995**, *252*, 248–262.
- (12) For a review of non-native peptide ligation strategies, see: Tam, J. P.; Xu, J.; Eom, K. D. *Biopolymers* **2001**, *60*, 194–205.

JA0450408