


Communication

**Dihydrogen Complexes of Electrophilic Metal Centers:  
 Observation of Cr(CO)(H), W(CO)(H) and [Re(CO)(H)]**

Steven L. Matthews, Vincent Pons, and D. Michael Heinekey

*J. Am. Chem. Soc.*, **2005**, 127 (3), 850-851 • DOI: 10.1021/ja0433370 • Publication Date (Web): 24 December 2004

Downloaded from <http://pubs.acs.org> on March 24, 2009



**More About This Article**

Additional resources and features associated with this article are available within the HTML version:

- Supporting Information
- Links to the 4 articles that cite this article, as of the time of this article download
- Access to high resolution figures
- Links to articles and content related to this article
- Copyright permission to reproduce figures and/or text from this article

[View the Full Text HTML](#)



**ACS Publications**  
 High quality. High impact.


## Dihydrogen Complexes of Electrophilic Metal Centers: Observation of $\text{Cr}(\text{CO})_5(\text{H}_2)$ , $\text{W}(\text{CO})_5(\text{H}_2)$ and $[\text{Re}(\text{CO})_5(\text{H}_2)]^+$

Steven L. Matthews, Vincent Pons, and D. Michael Heinekey\*

Department of Chemistry, University of Washington, Box 351700, Seattle, Washington 98195-1700

Received November 4, 2004; E-mail: heinekey@chem.washington.edu

The first examples of isolable transition metal dihydrogen complexes were reported by Kubas and co-workers 20 years ago.<sup>1</sup> These complexes are exemplified by tungsten complexes such as  $\text{W}(\text{CO})_3(\text{PR}_3)_2(\text{H}_2)$  ( $\text{R} = \text{Pr}$ ; Cy) and their Mo analogues, which bind hydrogen relatively weakly and are prepared by displacement of an agostic interaction with  $\text{H}_2$  gas. Subsequently, a Cr analogue was reported, which binds hydrogen more weakly.<sup>2</sup> Relatively weak binding of  $\text{H}_2$  in these complexes is signaled by only slight elongation of the H–H distance ( $d_{\text{HH}}$ ) compared to  $\text{H}_2$  gas. Values of  $d_{\text{HH}} = 0.85(1) \text{ \AA}$  ( $\text{Cr}$ ),<sup>2</sup>  $0.87(1) \text{ \AA}$  ( $\text{Mo}$ ),<sup>3</sup> and  $0.89(1) \text{ \AA}$  ( $\text{W}$ )<sup>3</sup> for these complexes have been determined by solid-state NMR measurements of dipolar coupling. A closely related class of dihydrogen complexes consists of molecules that are not isolable but have been studied in detail under matrix isolation conditions,<sup>4</sup> in liquid xenon,<sup>5</sup> or by fast spectroscopy methods.<sup>6</sup> For example, the prototypical molecules  $\text{M}(\text{CO})_5(\text{H}_2)$  ( $\text{M} = \text{Cr}$ ,  $\text{Mo}$ , and  $\text{W}$ ) have been observed in liquid Xe and in supercritical Xe, and are definitively characterized as dihydrogen complexes based on infrared spectroscopy, including studies of isotopically substituted derivatives. While vibrational data such as the detection in the infrared spectrum of the H–H stretching mode can confirm the presence of a dihydrogen ligand, quantitative structural information is lacking. The key structural parameter is the H–H distance ( $d_{\text{HH}}$ ), which is best determined by diffraction or NMR methods. Since the limited stability of these complexes precludes isolation, solution NMR spectroscopy becomes the method of choice. Hydrogen complexes with only CO as co-ligands are of fundamental interest in that these very electrophilic metal centers may bind hydrogen relatively weakly but provide substantial activation of the H–H bond toward heterolysis. These simple molecules have also been the focus of significant computational effort. The best computational results give  $d_{\text{HH}}$  of ca.  $0.8 \text{ \AA}$  for  $\text{M}(\text{CO})_5(\text{H}_2)$ , slightly shorter than that for the computed structures for  $\text{M}(\text{CO})_3(\text{PH}_3)_2(\text{H}_2)$ , which are models for the isolable phosphine-containing complexes. In this work, we present investigations of the structure and reactivity of such complexes in conventional solvents using NMR spectroscopy, which allows for a direct experimental test of the computed structures.



Photolysis (Hg lamp, 195 K) of  $\text{W}(\text{CO})_6$  solutions ( $\text{CD}_2\text{Cl}_2$ , 5 mM) under 20 psi of hydrogen gas affords a thermally labile dihydrogen complex (**1**). The  $^1\text{H}$  NMR spectrum of **1** in the hydride region is shown in Figure 1. A single resonance is observed at  $\delta =$





Figure 1. Partial (hydride region)  $^1\text{H}$  NMR spectra ( $\text{CD}_2\text{Cl}_2$ , 500 MHz, 240 K) of **1** (bottom) and **1-d<sub>1</sub>** (top).

–3.88 ppm with  $J_{\text{HW}} \approx 35 \text{ Hz}$ . Measurement of the relaxation time  $T_1$  of this resonance at temperatures between 190 and 220 K gives values between 25 and 60 ms, respectively (500 MHz). The relaxation time was still decreasing at the lowest temperature accessible, and the maximum rate of relaxation could not be measured. Although a quantitative analysis of these data is not possible, these observations are suggestive of a dihydrogen structure for complex **1**.<sup>7</sup> This was confirmed by carrying out the photolysis reaction under 15 psi of HD gas, which led to the spectrum shown in Figure 1. Complex **1** is formulated as a dihydrogen complex  $\text{W}(\text{CO})_5(\text{H}_2)$ .<sup>8</sup> The measured value of  $J_{\text{HD}}$  is 35.3 Hz, which is consistent with  $d_{\text{HH}} = 0.85 \text{ \AA}$ .<sup>9</sup> This coupling can be compared to the value of 34.0 Hz ( $d_{\text{HH}} = 0.87 \text{ \AA}$ ) reported for  $\text{W}(\text{CO})_3(\text{PiPr}_3)_2(\text{HD})$ .<sup>10</sup> Measurement of  $d_{\text{HH}}$  by determination of the dipolar coupling in the bound  $\text{H}_2$  ligand of  $\text{W}(\text{CO})_3(\text{PCy}_3)_2(\text{H}_2)$  by solid-state NMR gives a value of  $0.890(6) \text{ \AA}$ .<sup>3</sup> The HD coupling data for complex **1** indicates that the interaction of  $\text{H}_2$  with the metal center results in less activation of  $\text{H}_2$  than in the original Kubas complex. Due to the narrower line width exhibited by the hydride signal in **1-d<sub>1</sub>**, it was also possible to accurately measure  $J_{\text{HW}} = 38.2 \text{ Hz}$  (Figure 1), which is larger than the value of  $J_{\text{HW}} = 33.6 \text{ Hz}$  reported for  $\text{W}(\text{CO})_3(\text{PiPr}_3)_2(\text{H}_2)$ .<sup>10</sup>

The chromium analogue  $\text{Cr}(\text{CO})_5(\text{H}_2)$  (**2**) has been similarly prepared and exhibits a single hydride resonance at  $\delta = -7.45 \text{ ppm}$ . When the preparative reaction was carried out with HD gas, complex **2-d<sub>1</sub>** is obtained, with  $J_{\text{HD}} = 35.8 \text{ Hz}$ , only slightly different from that observed in **1-d<sub>1</sub>**. This is the largest value for  $J_{\text{HD}}$  reported to date, corresponding to  $d_{\text{HH}} = 0.84 \text{ \AA}$ , which is the shortest value of  $d_{\text{HH}}$  thus far observed for a dihydrogen complex. This observation



**Figure 2.** Partial (hydride region)  $^1\text{H}$  NMR spectrum (fluorobenzene- $d_5$ , 500 MHz, 260 K) of  $\mathbf{3-d_1}$ . The spectrum shown was obtained using a  $180^\circ$ - $\tau$ -90 pulse sequence with  $\tau = 16$  ms.

is consistent with a weak interaction of  $\text{H}_2$  with this highly electrophilic metal center. Complex  $\mathbf{2}$  had been previously prepared by photolysis of alkane solutions of  $\text{Cr}(\text{CO})_6$  under  $\text{H}_2$  pressure and found to decay with a half-life of ca. 25 s at ambient temperature.<sup>11</sup>

The experimentally determined values of  $d_{\text{HH}}$  for complexes  $\mathbf{1}$  and  $\mathbf{2}$  can be compared to the values reported from computational studies. A study using DFT methods gave values for  $d_{\text{HH}}$  of ca. 0.81–0.83 Å for  $\text{M}(\text{CO})_3(\text{PH}_3)_2(\text{H}_2)$ , which are computational models for the Cr, Mo, and W phosphine complexes of Kubas.<sup>12</sup> Using the same methodology,  $d_{\text{HH}}$  of ca. 0.79–0.80 Å was reported for  $\text{M}(\text{CO})_5(\text{H}_2)$ , slightly shorter than the values for the phosphine complexes, as expected for less basic metal centers. The calculations seem to significantly underestimate  $d_{\text{HH}}$  in the complexes of Kubas and also in the new complexes  $\mathbf{1}$  and  $\mathbf{2}$ .

The bound  $\text{H}_2$  ligand in complex  $\mathbf{1}$  is significantly activated with respect to heterolytic cleavage, with deprotonation to the known<sup>13</sup> anion  $[\text{HW}(\text{CO})_5]^-$  by one equivalent of  $\text{Et}_3\text{N}$  or excess (10 equiv) of  $\text{H}_2\text{O}$ . As expected, complex  $\mathbf{1}$  is significantly more acidic than the phosphine-containing analogues reported by Kubas and co-workers.<sup>14</sup>

We sought to evaluate the effect of a positive charge on the complexation of  $\text{H}_2$ , which we previously reported led to stronger hydrogen binding for cationic Re analogues of the neutral W complexes of Kubas.<sup>15,16</sup> Thus we examined the reaction of  $\text{Re}(\text{CO})_5\text{Cl}$  with  $[\text{Et}_3\text{Si}][\text{B}(\text{C}_6\text{F}_5)_4]^{17}$  in fluorobenzene solution under  $\text{H}_2$  gas. An immediate reaction occurs to give a new species  $\mathbf{3}$  with a single hydride signal at  $\delta = -4.60$  ppm. Complex  $\mathbf{3}$  can also be generated by reaction of  $\text{Re}(\text{CO})_5\text{H}$  with excess triflic acid. A relaxation time measurement at 260 K for the hydride resonance in complex  $\mathbf{3}$  gives a value of 34 ms, which diminishes slightly to 29 ms at 240 K (500 MHz). Since  $T_{1(\text{min})}$  could not be reached, it is not possible to deduce a value for  $d_{\text{HH}}$  from these data, but  $\mathbf{3}$  can be formulated as the dihydrogen complex  $[\text{Re}(\text{CO})_5(\text{H}_2)][\text{B}(\text{C}_6\text{F}_5)_4]$ . When the preparative reaction was carried out under HD gas, complex  $\mathbf{3-d_1}$  was obtained, with  $J_{\text{HD}} = 33.9$  Hz (Figure 2).

Isolation of  $\mathbf{3}$  as a solid has proven elusive due to the very high acidity of this complex (vide infra). The spectroscopic properties of complex  $\mathbf{3}$  are very similar to those of the phosphine-containing complex  $[\text{cis}-(\text{CO})_4(\text{PPh}_3)\text{Re}(\text{H}_2)]^+$ , previously reported by Kubas and co-workers.<sup>18</sup> In particular, the value for  $J_{\text{HD}}$ , and thus  $d_{\text{HH}}$ , are identical. Replacement of a  $\text{PPh}_3$  ligand with a CO presumably leads to much reduced electron density at the metal center but does

not affect  $d_{\text{HH}}$ . This confirms the crucial role of the trans CO ligand in controlling the degree of  $\text{H}_2$  activation, as noted previously by Kubas.<sup>19</sup>

The acidity of complex  $\mathbf{3}$  is greater than that of the neutral tungsten analogue. In contrast to complexes  $\mathbf{1}$  and  $\mathbf{2}$ , isotopically pure  $\mathbf{3-d_1}$  could not be cleanly prepared and invariably contained significant amounts of  $\mathbf{3}$  (Figure 2). We attribute this to the greater acidity of  $\mathbf{3}$  and surmise that  $\mathbf{3}$  arises from  $\mathbf{3-d_1}$  by intermolecular proton (deuteron) exchange catalyzed by traces of adventitious bases such as water, as previously demonstrated for cationic dihydrogen complexes of Re.<sup>20</sup> This acidity was confirmed by reaction of  $\mathbf{3}$  with excess  $\text{Et}_2\text{O}$ , which led to complete deprotonation to the neutral  $\text{Re}(\text{CO})_5\text{H}$ .

The preparation of these simple dihydrogen complexes demonstrates that binding of hydrogen to very electrophilic metal centers leads to dramatic activation of the bound hydrogen toward heterolysis. This is despite the H–H distance becoming only slightly elongated in comparison to the distance in free hydrogen gas, suggesting a relatively weak interaction with the metal center. We are continuing to investigate the preparation and reactivity of  $\sigma$ -bond complexes of highly electrophilic metal centers.

**Acknowledgment.** This research was supported by the National Science Foundation.

## References

- (1) Kubas, G. J.; Ryan, R. R.; Swanson, B. I.; Vergamini, P. J.; Wasserman, H. J. *J. Am. Chem. Soc.* **1984**, *106*, 451–452.
- (2) Kubas, G. J.; Nelson, J. E.; Bryan, J. C.; Eckert, J. E.; Wisniewski, L.; Zilm, K. *Inorg. Chem.* **1994**, *33*, 2954–2960.
- (3) Zilm, K. W.; Millar, J. M. *Adv. Magn. Opt. Res.* **1990**, *15*, 163–200.
- (4) Cf. Sweany, R. L. In *Transition Metal Hydrides*; Dedieu, A., Ed.; VCH: New York, 1992.
- (5) Upmacis, R. K.; Poliakoff, M.; Turner, J. J. *J. Am. Chem. Soc.* **1986**, *108*, 3645–3651.
- (6) Church, S. P.; Grevels, F. H.; Hermann, H.; Schaffner, K. *J. Chem. Soc., Chem. Commun.* **1985**, 30–32.
- (7) Desrosiers, P. J.; Cai, L.; Lin, Z.; Richards, R.; Halpern, J. *J. Am. Chem. Soc.* **1991**, *113*, 4173–4184.
- (8) We had previously reported that similar experimental conditions led to a dihydride species with a chemical shift of  $-2$  ppm and no HD coupling in the monodeuterated derivative (see Heinekey, D. M.; Law, J. K.; Shultz, S. M. *J. Am. Chem. Soc.* **2001**, *123*, 12728–12729). We now believe that the previously observed NMR signal may be due to an impurity.
- (9) Heinekey, D. M.; Luther, T. A. *Inorg. Chem.* **1996**, *35*, 4396–4399.
- (10) This coupling was originally reported as  $J_{\text{HD}} = 33.6$  Hz (Kubas, G. J.; Unkefer, C. J.; Swanson, B. I.; Fukuihshima, E. J. *J. Am. Chem. Soc.* **1986**, *107*, 7000–7006), but subsequent higher resolution data gives  $J_{\text{HD}} = 34.0$  and  $J_{\text{HW}} = 33.6$  Hz. (Gusev, D. G.; Kuhlman, R. L.; Renkema, K. B.; Eisenstein, O.; Caulton, K. G. *Inorg. Chem.* **1996**, *35*, 6775–6783.)
- (11) Upmacis, R. K.; Gadd, G. E.; Poliakoff, M.; Simpson, M. B.; Turner, J. J.; Whyman, R.; Simpson, A. F. *J. Chem. Soc., Chem. Commun.* **1985**, 27–30.
- (12) Tomas, J. Lledós, A.; Jean, Y. *Organometallics* **1998**, *17*, 4932–4939.
- (13) Darensbourg, M. Y.; Slater, S. *J. Am. Chem. Soc.* **1981**, *103*, 5914–5915.
- (14) Van Der Sluys, L. S.; Miller, M. M.; Kubas, G. J.; Caulton, K. G. *J. Am. Chem. Soc.* **1991**, *113*, 2513–2520.
- (15) Heinekey, D. M.; Radzewich, C. E.; Voges, M. H.; Schomber, B. M. *J. Am. Chem. Soc.* **1997**, *119*, 4172–4181.
- (16) Kubas, G. J. *Adv. Inorg. Chem.* **2004**, *56*, 127–177.
- (17) Gelabert, R.; Moreno, M.; Lluch, J. M.; Lledós, A.; Pons, V.; Heinekey, D. M. *J. Am. Chem. Soc.* **2004**, *126*, 8813–8822.
- (18) Huhmann-Vincent, J.; Scott, B. L.; Kubas, G. J. *Inorg. Chim. Acta* **1999**, *294*, 240–254.
- (19) Kubas, G. J. *Metal Dihydrogen and  $\sigma$ -Bond Complexes: Structure, Theory and Reactivity*; Kluwer: New York, 2001; p 147.
- (20) Heinekey, D. M.; Voges, M. H.; Barnhart, D. M. *J. Am. Chem. Soc.* **1996**, *118*, 10792–10802.

JA0433370