


Sulfoxide Carbon–Sulfur Bond Activation

Joseph M. O'Connor, Kevin D. Bunker, Arnold L. Rheingold, and Lev Zakharov

J. Am. Chem. Soc., **2005**, 127 (12), 4180-4181 • DOI: 10.1021/ja050160g • Publication Date (Web): 08 March 2005

Downloaded from <http://pubs.acs.org> on March 24, 2009



More About This Article

Additional resources and features associated with this article are available within the HTML version:

- Supporting Information
- Links to the 4 articles that cite this article, as of the time of this article download
- Access to high resolution figures
- Links to articles and content related to this article
- Copyright permission to reproduce figures and/or text from this article

[View the Full Text HTML](#)





Sulfoxide Carbon–Sulfur Bond Activation

Joseph M. O'Connor,* Kevin D. Bunker, Arnold L. Rheingold,* and Lev Zakharov

Department of Chemistry and Biochemistry, University of California, San Diego, La Jolla, California 92093


Received January 10, 2005; E-mail: jmoconnor@ucsd.edu

Transition metal insertion into the carbon–sulfur bond of sulfides to give well-characterized oxidative addition products has been achieved for cases involving sp-, sp²-, and sp³-hybridized carbon atoms.¹ The widespread use of sulfoxides in synthesis makes carbon–sulfur insertion reactions particularly desirable for higher oxidation state sulfur. Despite the existence of numerous metal–sulfoxide complexes,² carbon–sulfur bond activation has not previously been directly observed for sulfoxides. However, indirect evidence for transition metal insertion into the carbon–sulfur bond of sulfoxides is found in the sulfanylzincation reactions of alkynoates (eq 1)³ and the conversion of phenylsulfinyl ethene and Fe₂(CO)₉ to [Fe₂(μ-CH=CH₂)(μ-SPh)(CO)₆] (eq 2).⁴



We previously reported that (η⁵-C₅H₅)Co(PPh₃)₂ (**1**) undergoes reaction with alkynyl sulfone **2-Ph** at room temperature to give the η²-alkyne complex **3-Ph** in 75–80% yield (Scheme 1).⁵ There was

Scheme 1



no evidence for conversion of either **3-Ph** or **3-Tol** to a cobalto-sulfone complex (e.g., **4**). We now report that reaction of **1** and an alkynylsulfoxide leads to the first direct observation of carbon–sulfur bond activation involving a sulfoxide.

When the reaction of **1** (0.013 mmol) and alkynyl sulfoxide **5** (0.013 mmol) was monitored by ¹H NMR spectroscopy in benzene-*d*₆, the cobaltosulfoxide complex **6** was observed to form in 72% yield.⁶ In a larger scale reaction, **5** (0.30 mmol) was added to a dry benzene solution (20 mL) of **1** (0.30 mmol), and the mixture was allowed to stir at room temperature under a nitrogen atmosphere for 18 h. Removal of the volatiles and recrystallization of the residue led to isolation of **6** as dark purple, slightly air-sensitive crystals (54% yield). In the IR spectrum (NaCl) of **6**, ν(C≡C) is observed at 2030 cm⁻¹ and a strong band at 1064 cm⁻¹ is assigned to ν(S=O).⁷ The carbon–carbon triple bond stretch occurs 264 cm⁻¹ wavenumbers lower than that for noncoordinated **5**, and this compares

with a ν(C≡C) shift of 338 cm⁻¹ upon going from **2-Ph** to **3-Ph**. In the ¹³C{¹H} NMR spectrum (CDCl₃) of **6**, the alkynyl sp-carbons resonate at 90.2 (d, J_{CP} = 62.0 Hz) and 113.1 (d, J_{PC} = 40.4 Hz) ppm, shifted from 90.1 and 100.1 ppm in **5**. For comparison, the sp-carbons of the alkyne ligand in **3-Ph** are observed at 105.5 (d, J_{PC} = 3.5 Hz) and 118.1 (d, J_{PC} = 11.6 Hz) ppm. The observation of an sp-carbon resonance at 90 ppm and the large J_{PC} coupling constants indicated that the expected η²-alkyne complex (**7**), if formed, had not been isolated. The spectroscopic data and the known insertion of metals into the silicon–carbon bond of trimethylsilyl acetylenes⁸ raised the possibility of a (Cp)Co(PPh₃)-(TMS)(C≡CSOTol) structure for **6**. However, in the mass spectrum (FAB) of **6**, the base peak at 483 *m/z* [M⁺ – SOTol] and a fragment at 386 *m/z* [483 – CCTMS] suggested that cobalt had selectively inserted into the (sp)carbon–sulfur bond of **5**.

The metallosulfoxide structure of **6** was conclusively established by an X-ray crystallographic analysis (Figure 1, Table 1). Data col-




Figure 1. Solid-state molecular structures of **6-P1** (bottom) and **4** (top). Ellipsoids at the 30% probability level.

Table 1. Selected Bond Lengths (Å) and Angles (deg) for **4** and **6**

complex	6-P2(1)/n	6-P1	4-P2(1)/n
Co(1)–S(1)	2.2196(7)	2.2438(9)	2.2014(8)
Co(1)–P(1)	2.2108(7)	2.2047(9)	2.2391(9)
S(1)–O(1)	1.508(2)	1.513(2)	1.4588(17)
S(1)–O(2)			1.4698(16)
C(1)–Co(1)–S(1)	88.30(8)	88.91(10)	93.54(7)
Co(1)–S(1)–O(1)	108.61(8)	109.54(9)	104.11(11)
Co(1)–S(1)–O(2)			105.29(10)
Co(1)–S(1)–C(6)	106.33(8)	103.92(10)	107.34(8)
O(1)–S(1)–C(6)	105.03(12)	104.43(13)	114.27(7)
O(2)–S(1)–C(6)			109.47(7)
S(1)–Co(1)–P(1)	91.97(3)	90.73(3)	95.40(3)
Co(1)–C(1)–C(2)	176.1(2)	176.1(3)	174.5(2)
C(1)–Co(1)–P(1)	87.30(8)	91.98(9)	89.84(7)

lected on a needle of **6**, obtained as described above, refined in the centrosymmetric triclinic space group *P1*, indicating that the sample

was a racemate.⁹ Structural analysis of a second crystal, grown by slow evaporation of a benzene solution containing **6**, belonged to the centrosymmetric monoclinic space group $P2(1)/n$. Both polymorphs exhibited the same $S_{Co}, R_S/R_{Co}, S_S$ relative configurations. The sulfur atom in both structures of **6** exhibits a trigonal pyramidal geometry and is situated an average of 0.686 Å from the plane defined by Co(1), C(6), and O(1). This displacement is typical of other sulfenato ligands and organic sulfoxides (0.7–0.8 Å range). The average S(1)–O(1) [1.510(2) Å] and S(1)–C(6) [1.804(3) Å] distances in **6** are remarkably similar to those found in MeS(=O)–Tol [1.493(6) and 1.797(6) Å]. However, the Co(1)–S(1)–C(6) angle of 105.1° in **6** is 7.5° larger than the Me–S–Tol angle for MeS(=O)Tol,¹⁰ which may be attributed to greater steric congestion in **6**.


There are four structurally characterized cobaltosulfoxide complexes, all of which involve chelating [η^2 -(*N,S*)-H₂N(CHR)(CH₂)_{*n*}S(O)] sulfenato ligands.^{11–13} The average Co–S and S–O distances in this set of structures is 2.232 and 1.540 Å, respectively. More specifically, (NH₂CH₂CH₂NH₂)₂Co(NH₂CH₂CH₂SO)[SCN]₂ (**8**) exhibits a 96.51° Co–S–C angle, a S–O distance of 1.552(3) Å, and ν (S=O) of 986 cm⁻¹.^{11a} The sulfur–oxygen bond in **6** is significantly less polarized than that in **8**, based on the shorter S–O distance (1.510 Å) and higher frequency ν (S=O) stretch (1064 cm⁻¹) observed for **6**.

Metallosulfoxide **6** is capable of acting as both an oxygen atom acceptor and an oxygen atom donor. Exposure of a benzene solution of **6** to oxygen gas led to the formation of the metallosulfone **4**, which was isolated as bright-red crystals in 18% yield (Scheme 1).¹⁴ Ground state O₂ is known to oxidize both thiolato and *cis*-disulfenato ligands;^{12b,15} however, we are unaware of a precedent for conversion of a single sulfenato ligand to a sulfinato ligand upon reaction with O₂.¹⁶ Although it would be attractive to assume direct O₂ oxidation of sulfur, the current data do not allow for such a conclusion to be made.¹⁷

A comparison of the solid-state structure of **4** (Figure 1) to that for **6** reveals slightly larger S(1)–Co(1)–P(1) and S(1)–Co(1)–C(1) angles in the more congested metallosulfone. The sulfur ligands adopt very similar conformations with Cp–Co(1)–S(1)–O(1) and C(7)–C(6)–S(1)–O(1) torsion angles of –59.6 and –20.5° for **6** versus Cp–Co(1)–S(1)–O(2) and C(7)–C(6)–S(1)–O(2) torsion angles of +51.2 and –12.4° in **4**.

In benzene-*d*₆, **6** is slowly (months) transformed into a 4:1 ratio of thiolato-bridged cobalt dimers, **9-*eq,ax*** and **9-*eq,eq***, in 85% combined yield (eq 3).¹⁸ In addition, OPPh₃ (77%) and TMS-C≡C (43%) are observed as byproducts. Chromatographic workup of a large-scale reaction led to the collection of two distinct fractions (orange and green), but with the orange fraction rapidly turning green over the course of 15 min. ¹H NMR spectroscopic analysis of the two fractions indicated the same 4:1 ratio of dimers. Dark-green crystals of **9-*eq,eq*** were grown by slow evaporation of solvent, and an X-ray crystallographic analysis provided the first structural characterization of an *eq,eq*-[CpM(SR)]₂ complex. In benzene-*d*₆ at 70 °C, the two isomers of **9** interconvert rapidly on the NMR time scale, and the E_{act} for conversion of **9-*eq,ax*** to **9-*eq,eq*** was determined to be 74.9 ± 0.5 kJ/mol. For comparison, both at low temperature and in the solid state, [CpCo(SBu^t)₂ is

observed only as an *eq,ax*-isomer, which is reported to rapidly interconvert with the *ax,eq*-isomer at higher temperature (E_{act} = 60 kJ/mol).¹⁹



Acknowledgment. Support of the National Science Foundation (CHE-9975939) is gratefully acknowledged.

Supporting Information Available: Experimental procedures, characterization data for all new compounds, and crystallographic data for **4**, **6**, and **9-*eq,eq***. This material is available free of charge via the Internet at <http://pubs.acs.org>.

References

- (1) For leading references, see: (a) Matsunaga, P. T.; Hillhouse, G. L. *Angew. Chem., Int. Ed. Engl.* **1994**, *33*, 1748. (b) Vicic, D. A.; Jones, W. D. *J. Am. Chem. Soc.* **1999**, *121*, 7606. (c) Chen, J.; Angelici, R. J. *Coord. Chem. Rev.* **2000**, *206–207*, 63. (d) Savarin, C.; Srogl, J.; Liebeskind, L. S. *Org. Lett.* **2001**, *3*, 91.
- (2) Alessio, E. *Chem. Rev.* **2004**, *104*, 4203.
- (3) Maezaki, N.; Yagi, S.; Ohsawa, S.; Ohishi, H.; Tanaka, T. *Tetrahedron* **2003**, *59*, 9895 and references therein.
- (4) Ibbotson, A.; Reduto dos Reis, A. C.; Saberi, S. P.; Slawin, A. M. Z.; Thomas, S. E.; Tustin, G. J.; Williams, D. J. *J. Chem. Soc., Perkin Trans. I* **1992**, 1251.
- (5) O'Connor, J. M.; Ji, H.-L.; Iranpour, M.; Rheingold, A. L. *J. Am. Chem. Soc.* **1993**, *115*, 1586.
- (6) We are aware of only two organometallic sulfenato complexes: (a) George, T. A.; Watkins, D. D. *Inorg. Chem.* **1973**, *12*, 398. (b) Heinrich, L.; Mary-Verla, A.; Li, Y.; Vaissermann, J.; Chottard, J.-C. *Eur. J. Inorg. Chem.* **2001**, 2203.
- (7) The SO stretch assignment is supported by the absence of the 1064 cm⁻¹ band in the spectrum of metallosulfone **4** (vide infra).
- (8) Müller, C.; Lachicotte, R. J.; Jones, W. D. *Organometallics* **2002**, *21*, 1190 and references therein.
- (9) For racemization of sulfenato ligands, see: Kita, M.; Yamanari, K.; Shimura, Y. *Bull. Chem. Soc. Jpn.* **1982**, *55*, 2873.
- (10) De la Camp, U.; Hope, H. *Acta Crystallogr.* **1970**, *B26*, 846.
- (11) (a) Adzami, I. K.; Libson, K.; Lydon, J. D.; Elder, R. C.; Deutsch, E. *Inorg. Chem.* **1979**, *18*, 303. (b) Jackson, W. G.; Sargeson, A. M.; Whimp, P. O. *J. Chem. Soc., Chem. Commun.* **1976**, 934. (c) Kita, M.; Yamanari, K.; Kitahama, K.; Shimura, Y. *Bull. Chem. Soc. Jpn.* **1981**, *54*, 2995. (d) Okamoto, K.; Konno, T.; Einaga, H.; Hidaka, J. *Bull. Chem. Soc. Jpn.* **1987**, *60*, 393.
- (12) For metallosulfoxide structures of other metals, see: (a) Weigand, W.; Wunsch, R. *Chem. Ber.* **1996**, *129*, 1409. (b) Grapperhaus, C. A.; Darensbourg, M. Y. *Acc. Chem. Res.* **1998**, *31*, 451. (c) Kovacs, J. A. *Chem. Rev.* **2004**, *104*, 825.
- (13) Reaction of **1** with **5** may involve precoordination at the alkyne, the sulfoxide oxygen, or the sulfoxide sulfur. The observation that **4** does not undergo carbon–sulfur bond activation suggests one of the latter two possibilities. For a structurally characterized cobalt η^2 -sulfenato complex, see: Kung, I.; Schweitzer, D.; Shearer, J.; Taylor, W. D.; Jackson, H. L.; Lovell, S.; Kovacs, J. A. *J. Am. Chem. Soc.* **2000**, *122*, 8299.
- (14) For organometallic metallosulfones, see: (a) Wojcicki, A. *Adv. Organomet. Chem.* **1974**, *12*, 31. (b) Lefort, L.; Lachicotte, R. J.; Jones, W. D. *Organometallics* **1998**, *17*, 1420 and references therein.
- (15) Buonomo, R. M.; Font, I.; Maguire, M. J.; Reibenspies, J. H.; Tuntulani, T.; Darensbourg, M. Y. *J. Am. Chem. Soc.* **1995**, *117*, 963.
- (16) Conversion of a bis(sulfenato) complex to a bis(sulfinato) upon reaction with triplet oxygen has been suggested to involve addition of a single O₂ across the two *cis*-sulfur atoms (see ref 15).
- (17) Nickel sulfenates serve as O atom donors/acceptors: Farmer, P. J.; Verpeaux, J.-N.; Amatore, C.; Darensbourg, M. Y.; Musie, G. *J. Am. Chem. Soc.* **1994**, *116*, 9355.
- (18) Complex **9** has been previously reported, but with no characterization data or comment on the existence of two isomers: Frisch, P. D.; Lloyd, M. K.; McCleverty, J. A.; Seddon, D. J. *Chem. Soc., Dalton Trans.* **1973**, 2268.
- (19) Shaver, A.; Morris, S.; Turrin, R.; Day, V. W. *Inorg. Chem.* **1990**, *29*, 3622.

JA050160G