

Addition/Correction

**Fabrication of Colloidal Crystals with Tubular-like Packings [J. Am. Chem. Soc. 2005, 127, 3268–3269].**

Feng Li, Xavier Badel, Jan Linnros, and John B. Wiley

*J. Am. Chem. Soc.*, **2005**, 127 (19), 7262-7262 • DOI: 10.1021/ja052158v • Publication Date (Web): 21 April 2005

Downloaded from <http://pubs.acs.org> on March 25, 2009

**More About This Article**

---

Additional resources and features associated with this article are available within the HTML version:

- Supporting Information
- Access to high resolution figures
- Links to articles and content related to this article
- Copyright permission to reproduce figures and/or text from this article

[View the Full Text HTML](#)




**ACS Publications**  
High quality. High impact.

**Fabrication of Colloidal Crystals with Tubular-like Packings** [*J. Am. Chem. Soc.* **2005**, *127*, 3268–3269].

Feng Li, Xavier Badel, Jan Linnros, and John B. Wiley\*

Page 3268: The scale bar in Figure 1f was mislabeled. The corrected version of Figure 1 is shown below.



**Figure 1.**

JA052158V

10.1021/ja052158v

Published on Web 04/21/2005

**Total Synthesis of Cribrostatin IV: Fine-Tuning the Character of an Amide Bond by Remote Control** [*J. Am. Chem. Soc.* **2005**, *127*, 4596–4598].

Collin Chan, Richard Heid, Shengping Zheng, Jinsong Guo, Bishan Zhou, Takeshi Furuuchi, and Samuel J. Danishefsky\*

Page 4596. In paragraph 5, reference is made to a “DEF” ring precursor. In actuality, the statement refers to the “CDE” ring system.

Page 4597. (1*R*,2*R*)-(–)-*N*-*p*-Tosyl-1,2-diphenyl-ethylenediamine is the ligand used in the Noyori transfer hydrogenation. We incorrectly disclosed the opposite (*S,S*)-diamine ligand. We would like to thank Mr. Zhigang Wang for informing us of this error.

JA059909D

10.1021/ja059909d

Published on Web 04/22/2005