


Communication

Synthesis, Structure, and Spectroscopy of an Oxodiiron(II) Complex

Nathan A. Eckert, Sebastian Stoian, Jeremy M. Smith,
Emile L. Bominaar, Eckard Mnck, and Patrick L. Holland

J. Am. Chem. Soc., **2005**, 127 (26), 9344-9345 • DOI: 10.1021/ja0436704 • Publication Date (Web): 09 June 2005

Downloaded from <http://pubs.acs.org> on March 25, 2009



More About This Article

Additional resources and features associated with this article are available within the HTML version:

- Supporting Information
- Links to the 11 articles that cite this article, as of the time of this article download
- Access to high resolution figures
- Links to articles and content related to this article
- Copyright permission to reproduce figures and/or text from this article

[View the Full Text HTML](#)



ACS Publications
High quality. High impact.

Synthesis, Structure, and Spectroscopy of an Oxodiiron(II) Complex

Nathan A. Eckert,[†] Sebastian Stoian,[‡] Jeremy M. Smith,[†] Emile L. Bominaar,^{*,‡} Eckard Münck,^{*,‡} and Patrick L. Holland^{*,†}

Departments of Chemistry, University of Rochester, Rochester, New York 14627, and Carnegie Mellon University, Pittsburgh, Pennsylvania 15213

Received October 18, 2004; Revised Manuscript Received May 23, 2005; E-mail: holland@chem.rochester.edu

Diiron units with bridging oxygen ligands (i.e. O^{2-} , OH^- , or H_2O) are present in metalloenzymes such as methane monooxygenase, Δ^9 -desaturase, and ribonucleotide reductase.¹ In these enzymes, the Fe_2 unit is bridged by one or two oxygen-based ligands.¹ As these enzymes activate O_2 and transfer oxygen atoms to various substrates, the iron centers pass through several oxidation states from $Fe^{II}Fe^{II}$ to $Fe^{IV}Fe^{IV}$.¹

To better understand the mechanisms of these enzymes, chemists have synthesized oxo-bridged diiron complexes in a variety of oxidation states.^{2,3} Oxodiiron(III) complexes are very common, while their reduced analogues (i.e. $Fe^{II}Fe^{III}$ or $Fe^{II}Fe^{II}$) are rare.² For example, there are only two isolable and fully characterized $Fe^{II}Fe^{III}$ oxo compounds.⁴ Others have been inferred from electrochemical studies.⁵ There is no published oxodiiron(II) compound that is stable under ambient conditions.⁶ Here we report the synthesis, structure, and spectroscopic properties of $[L^{tBu}Fe]_2O$ ($L^{tBu} = ArNC(tBu)CHC(tBu)Ar^-$, where $Ar = 2,6$ -diisopropylphenyl), an isolable oxo-bridged diferrous compound. Although the trigonal planar coordination of the iron atoms in this compound is different from the geometries in iron enzymes,¹ the ability to stabilize this unusual oxidation level is notable because it demonstrates a new direction in oxodiiron chemistry.

The reaction of equimolar amounts of $[L^{tBu}FeH]_2$ ⁷ and H_2O in THF gives orange-red $[L^{tBu}Fe]_2O$ in 71% yield.⁸ The same compound can also be accessed in 53% yield by adding 1 equiv of Me_3NO to $L^{tBu}FeNNFeL^{tBu}$.⁹ The solid-state structure of $[L^{tBu}Fe]_2O$ is shown in Figure 1.⁸ There is a crystallographically imposed C_2 axis that renders the two $L^{tBu}Fe$ units equivalent. The molecule is planar at iron (sum of bond angles = $359.4(1)^\circ$), as found in other three-coordinate iron diketiminate compounds.^{7,9,10} The $Fe-O-Fe$ angle of $167.55(14)^\circ$ is only slightly bent, and the two diketiminate planes are nearly perpendicular (dihedral angle of $70.09(4)^\circ$).

The $Fe-O$ distance of $1.7503(4)$ Å is difficult to put into context because the only iron(II) compounds with oxo ligands are part of mixed-valence dimers that do not have localized valence on the crystallographic time scale.⁴ However, there are many examples of crystallographically characterized oxodiiron(III) compounds ($Fe-O$, av 1.774 Å, std dev 0.029 Å).¹¹ Because the ionic radius of Fe^{II} (0.77 Å) is substantially larger than that of Fe^{III} (0.63 Å),¹² it is noteworthy that the two $Fe-O$ bonds in $[L^{tBu}Fe]_2O$ are as short as those observed for $Fe(III)$ -oxo bridges. They are substantially shorter than the $Fe-O$ distances in iron(II) hydroxides.^{4b,c,13} Density-functional calculations show that reduction from the hypothetical diiron(III) complex $[L^{tBu}Fe]_2O^{2+}$ to diiron(II) places two electrons in d_z^2 orbitals, where the z axes are oriented normal to the diketiminate planes. As these orbitals have nonbonding character with respect to the ligands, the $Fe-O$ distances calculated for the diiron(II) species ($Fe-O = 1.76$ Å) are very close to those calculated for diiron(III) ($Fe-O = 1.77$ Å).




Figure 1. Thermal ellipsoid plot of $[L^{tBu}Fe]_2O \cdot OEt_2$ (50% probability ellipsoids), viewed along the C_2 axis. Hydrogen atoms and unassociated diethyl ether are omitted for clarity. Relevant distances (Å) and angles (deg): $Fe-O$ 1.7503(4), $Fe-N1$ 2.005(2), $Fe-N2$ 1.966(2), $Fe-O-Fe$ $167.55(14)$, $N1-Fe-N2$ $98.22(7)$.

Solid-state FTIR spectra of $[L^{tBu}Fe]_2O$ show an intense band at 868 cm^{-1} that shifts to 833 cm^{-1} when $H_2^{18}O$ is used to prepare the oxo complex (Figure 2). The 35 cm^{-1} shift is consistent with that expected for an $Fe-O$ harmonic oscillator. We assign this peak to the asymmetric stretching vibration of the $Fe-O-Fe$ core, on the basis of the correlation between $Fe-O-Fe$ stretching frequencies and $Fe-O-Fe$ bond angles in oxodiiron(III) complexes.¹⁴ No isotope-sensitive bands are observed in solution resonance Raman spectra, despite excitation into electronic absorptions at 380 and 560 nm.

The zero-field Mössbauer spectrum of solid $[L^{tBu}Fe]_2O$ at 4.2 K (Figure 3A) exhibits a quadrupole doublet with $\Delta E_Q = 1.42(2)$ mm/s and $\delta = 0.64(2)$ mm/s (vs Fe metal at 298 K). In frozen toluene solution we observe $\Delta E_Q = 1.44(2)$ mm/s and $\delta = 0.79(2)$ mm/s.¹⁵ Given the low coordination numbers, the iron(II) sites are expected to be high-spin, like the three-coordinate $Fe(II)$ diketiminate complexes reported previously.¹⁶




Figure 2. Infrared spectra of KBr pellets of $[L^{tBu}Fe]_2O$ and its ^{18}O isotopomer. Vertical scaling of the subtraction has been increased for clarity.

[†] University of Rochester.

[‡] Carnegie Mellon University.




Figure 3. Mössbauer spectra of solid $[L^{\text{tBu}}\text{Fe}]_2\text{O}$ at 4.2 K, recorded in zero field (A) and a parallel applied field of 4.0 T (B). The solid line in (B) is a computer simulation for $\Delta E_Q = (eQV_{zz}/12)[(1 + (\eta^2/3))]^{1/2} = +1.42$ mm/s, $\eta = 0$, $\delta = 0.64$ mm/s, and $B_{\text{eff},x} = (1 + \alpha_x)B$ for $\alpha_x = 0.2$; details are given in the Supporting Information. Absorption from a contaminant with broad, unresolved features (ca. 20%) is indicated by arrows.¹⁷

We have recorded Mössbauer spectra for two solid samples and for a toluene sample at temperatures from 4.2 to 150 K and applied fields, **B**, up to 7.0 T. The applied-field spectra have shapes very similar to those observed for diamagnetic compounds.¹⁸ However, at all temperatures and for all fields, in the solid as well as in the frozen solution, the magnetic splitting of the low-energy features is slightly larger than accounted for by the applied field (see Figure S-1, Supporting Information). The applied-field spectra yield $\Delta E_Q > 0$ and a small asymmetry parameter, $0 \leq \eta \leq 0.4$. Within the resolution, the splitting of the low-field features can be modeled by adding a *positive* magnetic hyperfine field, \mathbf{B}_{int} , perpendicular to the largest component of the electric field gradient, V_{zz} . Our simulations reveal that \mathbf{B}_{int} is proportional to the applied field, $\mathbf{B}_{\text{int}} = \alpha\mathbf{B}$, but independent of temperature up to 150 K. The simulation in Figure 3B assumes that the effective field along *x* is given by $B_{\text{eff},x} = (1 + \alpha_x)B$, with $\alpha_x \approx +0.2$ and $\alpha_y = \alpha_z = 0$ (see Supporting Information for details). The observation of a positive and temperature-independent α_x suggests substantial unquenched orbital angular momentum, as observed by us in other three-coordinate iron(II) complexes.¹⁶

Room-temperature ^1H NMR spectra of $[L^{\text{tBu}}\text{Fe}]_2\text{O}$ in C_6D_6 exhibit only seven paramagnetically shifted resonances, with relative intensities as expected for the coordinated diketiminate ligands.¹⁹ This shows that the diketiminate ligands are equivalent on the NMR time scale at room temperature, and the molecule has averaged D_{2h} or D_{2d} symmetry. The observation of equivalent Fe sites agrees with the single high-spin iron(II) environment seen by Mössbauer spectroscopy in frozen toluene solution.

In conclusion, we have synthesized and characterized a stable oxodiiron(II) complex. It contains two identical high-spin, three-coordinate Fe(II) centers supported by bulky diketiminate ligands. Further studies on the reactivity, electronic structure, and magnetic properties of this interesting complex are underway.

Acknowledgment. The authors thank the National Science Foundation (CHE-0134658, P.L.H.; MCB-0424494, E.M.), the University of Rochester (Weissberger Memorial Fellowship, N.A.E.), and the A. P. Sloan Foundation (P.L.H.) for funding, and Profs. K. R. Rodgers and G. Lukat-Rodgers for resonance Raman measurements.

Supporting Information Available: Synthetic, spectroscopic, and crystallographic data, and computational details (PDF, CIF). This material is available free of charge via the Internet at <http://pubs.acs.org>.

References

- (1) (a) Wallar, B. J.; Lipscomb, J. D. *Chem. Rev.* **1996**, *96*, 2625. (b) Solomon, E. I.; Brunold, T. C.; Davis, M. I.; Kemsley, J. N.; Lee, S. K.; Lehnert, N.; Neese, F.; Skulan, A. J.; Yang, Y. S.; Zhou, J. *Chem. Rev.* **2000**, *100*, 235.
- (2) (a) Kurtz, D. M. *Chem. Rev.* **1990**, *90*, 585 and references therein. (b) Tshuva, E. Y.; Lippard, S. J. *Chem. Rev.* **2004**, *104*, 987.
- (3) Reviews and recent examples: (a) Fontecave, M.; Ménage, S.; Duboc-Toia, C. *Coord. Chem. Rev.* **1998**, *178–180*, 1555. (b) Du Bois, J.; Mizoguchi, T. J.; Lippard, S. J. *Coord. Chem. Rev.* **2000**, *200–202*, 443. (c) Costas, M.; Chen, K. C.; Que, L. *Coord. Chem. Rev.* **2000**, *200–202*, 517. (d) Liu, C.; Yu, S.; Li, D.; Liao, Z.; Sun, X.; Xu, H. *Inorg. Chem.* **2002**, *41*, 913. (e) Tshuva, E. Y.; Lee, D.; Bu, W.; Lippard, S. J. *J. Am. Chem. Soc.* **2002**, *124*, 2416. (f) Tolman, W. B.; Que, L. *J. Chem. Soc., Dalton Trans.* **2002**, *5*, 653. (g) Raffard-Pons y Moll, N.; Banse, F.; Miki, K.; Nierlich, M.; Girerd, J. *Eur. J. Inorg. Chem.* **2002**, *8*, 1941. (h) Marlin, D. S.; Olmstead, M. M.; Mascharak, P. K. *Inorg. Chem.* **2003**, *42*, 1681.
- (4) (a) Arena, F.; Floriani, C.; Chiesi-Villa, A.; Guastini, C. *J. Chem. Soc., Chem. Commun.* **1986**, 1369. (b) Cohen, J. D.; Payne, S. C.; Hagen, K. C.; Sanders-Loehr, J. *J. Am. Chem. Soc.* **1997**, *119*, 2960. (c) Payne, S. C.; Hagen, K. S. *J. Am. Chem. Soc.* **2000**, *122*, 6399.
- (5) (a) Hartman, J. R.; Rardin, R. L.; Chaudhuri, P.; Pohl, K.; Wiegardt, K.; Nuber, B.; Weiss, J.; Papaefthymiou, G. C.; Frankel, R. B.; Lippard, S. J. *J. Am. Chem. Soc.* **1987**, *109*, 7387. (b) Feig, A. L.; Masschelein, A.; Bakac, A.; Lippard, S. J. *J. Am. Chem. Soc.* **1997**, *119*, 334. (c) Davydov, R. M.; Ménage, S.; Fontecave, M.; Gräslund; Ehrenberg, A. *J. Biol. Inorg. Chem.* **1997**, *2*, 242. (d) Musie, G.; Lai, C.; Reibenspies, J. H.; Sumner, L. W.; Darensbourg, M. Y. *Inorg. Chem.* **1998**, *37*, 4086. (e) Kiani, S.; Tapper, A.; Staples, R. J.; Stavropoulos, P. *J. Am. Chem. Soc.* **2000**, *122*, 7503.
- (6) In one case, electrochemical experiments imply transient stability of an oxoiron(II) compound. See ref 5d.
- (7) Smith, J. M.; Lachicotte, R. J.; Holland, P. L. *J. Am. Chem. Soc.* **2003**, *125*, 15752.
- (8) Synthetic and crystallographic details are in the Supporting Information.
- (9) Smith, J. M.; Lachicotte, R. J.; Pittard, K. A.; Cundari, T. R.; Lukat-Rodgers, G.; Rodgers, K. R.; Holland, P. L. *J. Am. Chem. Soc.* **2001**, *123*, 9222.
- (10) (a) Smith, J. M.; Lachicotte, R. J.; Holland, P. L. *Chem. Commun.* **2001**, 1542. (b) Smith, J. M.; Lachicotte, R. J.; Holland, P. L. *Organometallics* **2002**, *21*, 4808. (c) Sciarone, T. J.; Meetsma, A.; Hessen, B.; Teuben, J. H. *Chem. Commun.* **2002**, 1580. (d) Vela, J.; Smith, J. M.; Lachicotte, R. J.; Holland, P. L. *Chem. Commun.* **2002**, 2886. (e) Gibson, V. C.; Marshall, E. L.; Navarro-Llobet, D.; White, A. J. P.; Williams, D. J. *Dalton Trans.* **2002**, *23*, 4321. (f) Panda, A.; Stender, M.; Wright, R. J.; Olmstead, M. M.; Klavins, P.; Power, P. P. *Inorg. Chem.* **2002**, *41*, 3909. (g) Eckert, N. A.; Smith, J. M.; Lachicotte, R. J.; Holland, P. L. *Inorg. Chem.* **2004**, *43*, 3306. (h) Vela, J.; Stoian, S.; Flaschenriem, C. J.; Münck, E.; Holland, P. L. *J. Am. Chem. Soc.* **2004**, *126*, 4522.
- (11) Allen, F. H. *Acta Crystallogr.* **2002**, *B58*, 380. A search of the Cambridge Structural Database provided 194 examples of crystallographically characterized compounds containing an unsupported Fe–O–Fe unit (i.e. no other bridging ligands). Only diiron compounds were included, and all the compounds found in the search were diiron(III). Histograms of the search results are included in the Supporting Information.
- (12) Values given are for four-coordinate iron, the lowest coordination number for which values are available. Shannon, R. D. *Acta Crystallogr.* **1976**, *A32*, 751.
- (13) (a) Kitajima, N.; Tamura, N.; Tanaka, M.; Moro-oka, Y. *Inorg. Chem.* **1992**, *31*, 3342. (b) Lachicotte, R. J.; Kitaygorodskiy, A.; Hagen, K. S. *J. Am. Chem. Soc.* **1993**, *115*, 8883. (c) Lee, D.; Pierce, B.; Krebs, C.; Hendrich, M. P.; Huynh, B. H.; Lippard, S. J. *J. Am. Chem. Soc.* **2002**, *124*, 3993. (d) Stubna, A.; Jo, D.-H.; Costas, M.; Brennessel, W. W.; Andres, H.; Bominaar, E. L.; Münck, E.; Que, L. *Inorg. Chem.* **2004**, *43*, 3067.
- (14) Sanders-Loehr, J.; Wheeler, W. D.; Shiemke, A. K.; Averill, B. A.; Loehr, T. M. *J. Am. Chem. Soc.* **1989**, *111*, 8084.
- (15) This surprising difference in isomer shift (δ) implies that **1** has a different conformation in solution. See Supporting Information for details.
- (16) Andres, H.; Bominaar, E. L.; Smith, J. M.; Eckert, N. A.; Holland, P. L.; Münck, E. *J. Am. Chem. Soc.* **2002**, *124*, 3012.
- (17) Our samples are prepared in a glovebox, and the sample holders are sealed with a shrink fit cap (Delrin over brass). However, the intensity of the contaminant signals increases over weeks during storage in liquid N_2 .
- (18) Antiferromagnetic coupling of the ferrous ions in $[L^{\text{tBu}}\text{Fe}]_2\text{O}$ is supported by density functional theory calculations (see Supporting Information) that indicate an exchange coupling $J \approx 200\text{--}250\text{ cm}^{-1}$ (with $\mathbf{H} = \mathbf{J}\mathbf{S}_1 \cdot \mathbf{S}_2$), in accord with a nearly linear Fe–O–Fe superexchange pathway.
- (19) The two methyl groups of the ligand isopropyl groups are inequivalent due to hindered rotation of the aryl moiety. See ref 10f and Supporting Information.

JA0436704