


Communication

**Oxygen Binding to Tyrosinase from *Streptomyces antibioticus* Studied by Laser Flash Photolysis**

Shun Hirota, Takumi Kawahara, Emanuela Lonardi, Ellen de Waal, Noriaki Funasaki, and Gerard W. Canters

*J. Am. Chem. Soc.*, **2005**, 127 (51), 17966-17967 • DOI: 10.1021/ja0541128 • Publication Date (Web): 03 December 2005

Downloaded from <http://pubs.acs.org> on March 25, 2009



**More About This Article**

Additional resources and features associated with this article are available within the HTML version:

- Supporting Information
- Links to the 3 articles that cite this article, as of the time of this article download
- Access to high resolution figures
- Links to articles and content related to this article
- Copyright permission to reproduce figures and/or text from this article

[View the Full Text HTML](#)

## Oxygen Binding to Tyrosinase from *Streptomyces antibioticus* Studied by Laser Flash Photolysis

Shun Hirota,<sup>\*,†,‡</sup> Takumi Kawahara,<sup>†</sup> Emanuela Lonardi,<sup>§</sup> Ellen de Waal,<sup>§</sup> Noriaki Funasaki,<sup>†</sup> and Gerard W. Canters<sup>§</sup>


Department of Physical Chemistry, 21st Century COE Program, Kyoto Pharmaceutical University, 5 Nakauchi-cho, Misasagi, Yamashina-ku, Kyoto 607-8414, Japan, PRESTO, JST, Kawaguchi, Saitama 332-0012, Japan, and Leiden Institute of Chemistry, Gorlaeus Laboratories, Leiden University, Einsteinweg 55, 2333 CC Leiden, The Netherlands

Received June 22, 2005; E-mail: hirota@mb.kyoto-phu.ac.jp


Tyrosinases (EC 1.14.18.1) are involved in the synthesis of melanin pigments in bacteria, plants, and mammals. They catalyze the *o*-hydroxylation of monophenols (monophenolase activity) and the oxidation of *o*-diphenols to *o*-quinones (diphenolase activity).<sup>1</sup> Tyrosinases possess a dinuclear copper active site similar to that of hemocyanin (Hc), which is an oxygen carrier protein in mollusks and arthropods, and to that of catechol oxidase.<sup>2</sup> Crystallographic studies have shown that Hc binds oxygen in a  $\mu$ - $\eta^2$ : $\eta^2$ -peroxodicycopper(II) binding mode,<sup>3</sup> while optical studies have shown that oxytyrosinase (Ty<sub>oxy</sub>) and oxyhemocyanin exhibit similar absorption bands at about 345 and 600 nm.<sup>4</sup> They also exhibit similar resonance Raman bands at about 750 cm<sup>-1</sup>, which are assigned to the peroxo stretching mode of the peroxodicycopper site.<sup>5</sup> From these results, as confirmed by NMR spectroscopy,<sup>6</sup> the active site structure of tyrosinase (Ty) is thought to be similar to that of Hc. Model compound studies have shown that a  $\mu$ -oxodicycopper(III) species may be in equilibrium with the  $\mu$ - $\eta^2$ : $\eta^2$ -peroxodicycopper(II) species.<sup>7</sup> Binding of O<sub>2</sub> to a copper complex has been observed by photorelease of carbon monoxide from the CO complex in a mixture of CO and O<sub>2</sub>.<sup>8</sup> Although Ty has attracted much interest because of its intriguing oxygen activation capability,<sup>1</sup> the information on the O<sub>2</sub>-binding character of Ty is limited. Therefore, we have studied by flash photolysis how small molecules, such as carbon monoxide and *p*-nitrophenol (a substrate-analogue inhibitor<sup>9</sup>), may affect O<sub>2</sub> binding kinetics.

After photodissociation (Scheme 1, A→B), O<sub>2</sub> can stay in the active site pocket of the protein and perform geminate rebinding (Scheme 1, B→A), or leave the protein pocket, especially at higher temperature (Scheme 1, B→C). The amount of O<sub>2</sub> dissociation from

**Scheme 1.** Photodissociation of O<sub>2</sub> from Tyrosinase



the protein may depend on the accessibility of the active site and the protein flexibility. When the O<sub>2</sub> molecule leaves the copper site of Ty after flash photolysis and rebinds to it, the absorbance change can be observed in the microsecond to millisecond time scale. This behavior is well-known for heme proteins. For example, above 200 K, equilibrium fluctuations among conformational substates of myoglobin (Mb) open pathways for the ligand through the protein matrix.<sup>10</sup>



**Figure 1.** Flash-photolysis measurements of Ty<sub>oxy</sub> by 355 nm pulse irradiation. Absorbance changes at 345 nm under (A) 100% O<sub>2</sub> and (B) O<sub>2</sub>:CO = 1:1 (v/v). (A) Single and (B) double exponential best-fitted curves are shown in red. Experimental conditions: Ty, 20  $\mu$ M; O<sub>2</sub>, (A) 1.54 mM and (B) 0.77 mM; 15 °C; 50 mM potassium phosphate buffer, pH 7.2; laser pulse, 30 mJ, 10 Hz. The traces are an average of (A) 128 and (B) 256 shots. (Inset) A: Overlap of wavelength dependence of the initial absorbance changes (empty circles) with the difference absorption spectrum (red, oxy minus deoxy). B: Absorption spectra of Ty under 100% O<sub>2</sub> (red) and O<sub>2</sub>:CO = 1:1 (v/v) (blue).


Purified met-tyrosinase was reduced with 1–10 mM hydroxylamine and dialyzed under anaerobic conditions.<sup>11</sup> The obtained reduced tyrosinase (deoxytyrosinase, Ty<sub>red</sub>) was converted into Ty<sub>oxy</sub> by aeration, and the solution was filtered before each measurement. The absorbance change of Ty<sub>oxy</sub> at 345 nm under O<sub>2</sub> by 355 nm pulse irradiation is shown in Figure 1A. The absorbance decreased immediately after the pulse and recovered on a microsecond time scale. The initial absorbance decrease corresponds to dissociation of O<sub>2</sub> from the dicopper center, whereas the intensity recovery indicates rebinding of O<sub>2</sub> to the deoxy enzyme. Geminate recombination was not observed, although it may occur within the dead time of the instrument.<sup>12</sup> A first-order rate constant was obtained by fitting the absorbance change with an exponential curve ( $(2.0 \pm 0.5) \times 10^4$  s<sup>-1</sup>).<sup>13,14</sup> The O<sub>2</sub> binding rate constant ( $k_{O_2}$ ) was then calculated as  $k_{O_2} = 13 \pm 3$   $\mu$ M<sup>-1</sup> s<sup>-1</sup> by taking the oxygen concentration (1.54 mM at 15 °C) into account.<sup>15</sup> The largest absorbance change was observed at 345 nm, and the wavelength dependence of the initial intensity change in the 305–405 nm range showed a good correlation with the oxy 345 nm absorption band (Figure 1A, inset), which indicates that the absorbance change is due to O<sub>2</sub> dissociation and rebinding. The O<sub>2</sub> rebinding kinetics did not change significantly on increasing the pH to 8.1, which suggests that there is no significant conformational change in Ty between pH 7.1 and 8.1.

To observe the effect of other molecules on O<sub>2</sub> binding, flash-photolysis experiments were performed with O<sub>2</sub> mixed with carbon monoxide (O<sub>2</sub>:CO = 1:1 (v/v)) (Figure 1B). O<sub>2</sub> and CO binding to the dinuclear copper active site are mutually exclusive,<sup>16</sup> and CO caused formation of Ty<sub>oxy</sub> to decrease to about 40% according to the absorption spectra (Figure 1B, inset).<sup>17</sup> Photodissociation of the

<sup>†</sup> Kyoto Pharmaceutical University.

<sup>‡</sup> PRESTO, JST.

<sup>§</sup> Leiden University.



**Figure 2.** Arrhenius plot of oxygen binding rate constant of  $Ty_{oxy}$  obtained by 355 nm pulse irradiation at different temperatures (5–25 °C). Other experimental conditions are the same as those listed in Figure 1.

**Table 1.** Kinetic and Thermodynamic Parameters for Oxygen Binding to Tyrosinase and Hemocyanin

	$k_{O_2}$ ( $\mu M^{-1} s^{-1}$ )	$k_{-O_2}$ ( $s^{-1}$ )	$K_D$ ( $\mu M$ )
Ty <i>Streptomyces antibioticus</i>	19 <sup>a</sup>	310 <sup>b</sup>	16.5
Ty <i>Agaricus bisporus</i> <sup>22</sup>	23	1100 <sup>b</sup>	46.6
Hc <i>Panulirus interruptus</i> <sup>123</sup>	31–154	150–470	2.0–9.1
Hc <i>Octopus vulgaris</i> <sup>24</sup>			90

<sup>a</sup> At 21 °C. <sup>b</sup> Calculated from  $k_{O_2}$  and  $K_D$ .

$O_2$  molecules from the active site by pulse irradiation also decreased to about 40%, whereas the recovery exhibited two phases (Figure 1B). The faster phase showed a rate similar to the binding rate observed in Figure 1A, and it is ascribed to  $O_2$  binding to  $Ty_{red}$ . The  $O_2$ -rebinding rate constant of the slower phase amounted to  $0.7 \pm 0.1 \text{ ms}^{-1}$ . This phase is attributed to  $Ty_{oxy}$  formation by dissociation of CO from the active site of the CO-bound species, the concentration of which increased transiently by  $O_2$  photodissociation.<sup>18</sup> The rate constant of the slower phase was similar to that obtained by stopped-flow mixing of a  $Ty_{red}$  solution under CO atmosphere with  $O_2$ -saturated buffer (results not shown). The intensity of the initial absorption change by pulse irradiation decreased significantly when 100  $\mu M$  *p*-nitrophenol was added, although  $Ty_{oxy}$  was generated as judged by the absorption spectrum. These results show that small molecules may occupy the active site and affect the  $O_2$  binding kinetics of Ty.

The  $O_2$ -binding rate constant and the initial intensity change increased with temperature (Figure 2). The larger intensity changes for higher temperatures may correspond to an enhancement in protein fluctuations, which increase the dissociation of the  $O_2$  molecule from the protein. It has been observed in flash-photolysis experiments on  $O_2$  binding to Mb that the geminate fraction decreases by raising the temperature, which has been interpreted as enhancement of structural fluctuations at higher temperatures.<sup>19</sup> Using the reported  $O_2$  dissociation constant for Ty from *Streptomyces antibioticus* ( $K_D = 16.5 \mu M$ ) at 21 °C and estimating  $k_{O_2}$  at 21 °C from Figure 2,<sup>20</sup> the  $O_2$  dissociation rate constant is calculated as  $310 \text{ s}^{-1}$  (Table 1). The activation enthalpy of the rate-limiting step of  $O_2$  binding is calculated by the temperature dependence of  $k_{O_2}$  to be  $12.8 \pm 2.6 \text{ kcal/mol}$ . It has been proposed that the rate-limiting step in CO binding to Mb involves some conformational change, and that conformational relaxation of Mb when going from

the CO-bound state toward the deoxy state increases the barrier for CO binding.<sup>21</sup> The activation enthalpy for  $O_2$  binding to Ty may also reflect a protein conformational change connected with binding of  $O_2$  since the Cu–Cu distance has been shown to decrease from 4.6 to 3.6 Å by oxygenation for Hc.<sup>3</sup> The present study observes  $O_2$  rebinding by flash photolysis and shows that the method would be a powerful tool to study the  $O_2$  binding kinetics in Ty.

**Acknowledgment.** This work was partially supported by Grants-in-Aid for Scientific Research from MEXT (Priority Areas, Water and Biomolecules, No. 164041242, S.H.), JSPS (Category C, No. 16550149, and the 21st COE Program (S.H.)), and JST (S.H.).

## References

- (1) Solomon, E. I.; Sundaram, U. M.; Machonkin, T. E. *Chem. Rev.* **1996**, *96*, 2563–2605.
- (2) Klabunde, T.; Eicken, C.; Sacchettini, J. C.; Krebs, B. *Nat. Struct. Biol.* **1998**, *5*, 1084–1090.
- (3) (a) Magnus, K. A.; Hazes, B.; Ton-That, H.; Bonaventura, C.; Bonaventura, J.; Hol, W. G. J. *Proteins* **1994**, *19*, 302–309. (b) Magnus, K. A.; Ton-That, H.; Carpenter, J. E. *Chem. Rev.* **1994**, *94*, 727–735. (c) Cuff, M. E.; Miller, K. I.; Holde, K. E. v.; Hendrickson, W. A. *J. Mol. Biol.* **1998**, *278*, 855–870.
- (4) (a) Jolley, R. L., Jr.; Evans, L. H.; Makino, N.; Mason, H. S. *J. Biol. Chem.* **1974**, *249*, 335–345. (b) Heirwegh, K.; Borginon, H.; Lontie, R. *Biochim. Biophys. Acta* **1961**, *48*, 517–526. (c) Van Holde, K. E. *Biochemistry* **1967**, *6*, 93–99.
- (5) (a) Ling, J.; Nestor, L. P.; Czernuszewicz, R. S.; Spiro, T. G.; Fraczkiewicz, R.; Sharma, K. D.; Loehr, T. M.; Sanders-Loehr, J. *J. Am. Chem. Soc.* **1994**, *116*, 7682–7691. (b) Eickman, N. C.; Solomon, E. I.; Larrabee, J. A.; Spiro, T. G.; Lerch, K. *J. Am. Chem. Soc.* **1978**, *100*, 6529–6531.
- (6) Bubacco, L.; Salgado, J.; Tepper, A. W. J. W.; Vijgenboom, E.; Canters, G. W. *FEBS Lett.* **1999**, *442*, 215–220.
- (7) (a) Halfen, J. A.; Mahapatra, S.; Wilkinson, E. C.; Kaderli, S.; Young, V. G., Jr.; Que, L., Jr.; Zuberbuhler, A. D.; Tolman, W. B. *Science* **1996**, *271*, 1397–1400. (b) Mahadevan, V.; Hou, Z.; Cole, A. P.; Root, D. E.; Lal, T. K.; Solomon, E. I.; Stack, T. D. P. *J. Am. Chem. Soc.* **1997**, *119*, 11996–11997.
- (8) Fry, H. C.; Scaltrito, D. V.; Karlin, K. D.; Meyer, G. J. *J. Am. Chem. Soc.* **2003**, *125*, 11866–11871.
- (9) Tepper, A. W. J. W.; Bubacco, L.; Canters, G. W. *J. Am. Chem. Soc.* **2005**, *127*, 567–575.
- (10) (a) Austin, R. H.; Beeson, K. W.; Eisenstein, L.; Frauenfelder, H.; Gunsalus, I. C. *Biochemistry* **1975**, *14*, 5355–5373. (b) Steinbach, P. J.; Ansari, A.; Berendzen, J.; Braunschweig, D.; Chu, K.; Cowen, B. R.; Ehrenstein, D.; Frauenfelder, H.; Johnson, J. B.; Lamb, D. C.; Luck, S.; Mourant, J. R.; Nienhaus, G. U.; Ormos, P.; Philipp, R.; Xie, A.; Young, R. D. *Biochemistry* **1991**, *30*, 3988–4001.
- (11) Tepper, A. W. J. W.; Bubacco, L.; Canters, G. W. *J. Biol. Chem.* **2002**, *277*, 30436–30444.
- (12) Geminate rebinding is difficult to observe due the difficulty in obtaining 355 nm picosecond pulses.
- (13) Within experimental error, the rate varied linearly with  $O_2$  concentration, although at less than 50%  $O_2$  saturation, low S/N precluded reliable determination of the rate.
- (14) Disappearance of the reduced species was also observed by monitoring the fluorescence change ( $Ty_{red}$  fluorescence is roughly halved upon binding of  $O_2$ ).<sup>9</sup> It occurred with a similar rate as observed in Figure 1.
- (15) Since  $k_{obs} = k_{O_2}[O_2] + k_{-O_2}$ , or  $k_{obs}/k_{-O_2} \approx (1/K_D)[O_2] + 1$ , it follows from the literature value of  $K_D$  and the value of  $k_{obs}$  that  $k_{obs} \approx k_{O_2}[O_2] \gg k_{-O_2}$ .
- (16) Decker, H.; Connelly, P. R.; Robert, C. H.; Gill, S. *Biochemistry* **1988**, *27*, 6901–6908.
- (17) It follows that the constant for the equilibrium  $Ty_{oxy} + CO \xrightleftharpoons[k_2]{k_1} CO\text{-bound Ty} + O_2$  is close to 1 and that  $k_1 \approx k_2$ .
- (18) The ratio of the amplitude of the slower phase against that of the faster phase increased with the  $[CO]/[O_2]$  ratio. The rate constant of the slower phase was constant, in agreement with  $k_1 \approx k_2$  (see ref 17).
- (19) Sato, F.; Shiro, Y.; Sakaguchi, Y.; Iizuka, T.; Hayashi, H. *J. Biol. Chem.* **1990**, *265*, 18823–18828.
- (20) Tepper, A. W. J. W.; Bubacco, L.; Canters, G. W. *J. Biol. Chem.* **2004**, *279*, 13425–13434.
- (21) Nienhaus, G. U.; Mourant, J. R.; Chu, K.; Frauenfelder, H. *Biochemistry* **1994**, *33*, 13413–13430.
- (22) Rodriguez-Lopez, J. N.; Fenoll, L. G.; Garcia-Ruiz, P. A.; Varon, R.; Tudela, J.; Thorneley, R. N. F.; Garcia-Canovas, F. *Biochemistry* **2000**, *39*, 10497–10506.
- (23) Andrew, C. R.; McKillop, K. P.; Sykes, A. G. *Biochim. Biophys. Acta* **1993**, *1162*, 105–114.
- (24) Salvato, B.; Santamaria, M.; Beltrami, M.; Alzuet, G.; Casella, L. *Biochemistry* **1998**, *37*, 14065–14077.

JA0541128