

Linkage Effects on Binding Affinity and Activation of GPR30 and Estrogen Receptors ER α/β with Tridentate Pyridin-2-yl Hydrazine Tricarbonyl–Re/^{99m}Tc(I) Chelates

Chinnasamy Ramesh,[†] Bj Bryant,[†] Tapan Nayak,^{‡,§} Chetana M. Revankar,[§] Tamara Anderson,[‡] Kathryn E. Carlson,[#] John A. Katzenellenbogen,[#] Larry A. Sklar,[¶] Jeffrey P. Norenberg,[‡] Eric R. Prossnitz,[§] and Jeffrey B. Arterburn^{*†}

Department of Chemistry and Biochemistry MSC 3C, New Mexico State University, P.O. Box 30001, Las Cruces, New Mexico 88003, Radiopharmaceutical Sciences Program, College of Pharmacy, University of New Mexico Health Sciences Center, Albuquerque, New Mexico 87131, Department of Cell Biology and Physiology and Department of Pathology, University of New Mexico, and Cancer Research and Treatment Center, Albuquerque, New Mexico 87131, and Department of Chemistry, University of Illinois, 600 South Mathews Avenue, Urbana, Illinois 61801

Received September 1, 2006; E-mail: jarterbu@nmsu.edu

The hormone, 17 β -estradiol (E2), mediates diverse physiological and regulatory control of reproduction, growth, development, immune response, bone mass, and neurological and cardiovascular health. Carcinomas of the breast, endometrium, and ovary are common malignancies in female reproductive tissues, and assessment of their estrogen responsiveness is crucial for prognosis and choice of therapeutic regimen. The classical model of estrogen activation describes ligand binding to receptors ER α/β initiating conformational changes, which result in interactions with transcriptional co-regulators and promoter DNA sequences of target genes. The resulting gene transcription provides a physiological response within hours following estrogen exposure. Additional non-genomic estrogen-induced rapid cell signaling pathways have been recognized as important contributors to the overall biological response.¹ A new mediator of estrogen-dependent signal transduction, the 7-transmembrane G-protein-coupled receptor GPR30, was recently identified.² This receptor can be coexpressed in cells with ER α/β or individually in cells lacking classical estrogen receptors. Revankar et al. demonstrated that GPR30 is an intracellular protein, in the endoplasmic reticulum, that binds estrogen with high affinity ($K_d \sim 6$ nM) and mediates rapid cellular responses, including calcium mobilization and phosphatidylinositol 3,4,5-trisphosphate production in the nucleus. The role of GPR30 in the growth and proliferation of estrogen-responsive tumors is a focus of investigation as a new biomarker and target for cancer diagnostics and therapeutics and could represent an important imaging target in ER α/β -negative cells.³

Noninvasive imaging technology offers great promise for the in vivo characterization of estrogen-dependent tumors. The intracellular location of ER α/β and GPR30 requires neutral, cell-permeable imaging agents. Organometallic tricarbonyl–^{99m}Tc- and ⁹⁴Tc-labeled estrogens have potential for diagnostic imaging using single photon emission computed tomography (SPECT) or positron emission tomography (PET).⁴ Complexes with promising in vitro ER α/β receptor binding affinity have been identified, but poor target tissue uptake and complicated radiolabeling procedures have inhibited further development.⁴ Lipophilicity and chelate instability are recognized problems, and other significant parameters may be identified through in vitro characterization of estrogen derivatives




Figure 1. X-ray ORTEP rendition of complex 1.

in cell culture. Herein, we report a new class of tridentate pyridin-2-yl hydrazine Re/^{99m}Tc chelates that exhibit strong binding affinity to GPR30 and ER α/β and evaluate intracellular binding with a functional cell-based receptor-mediated signaling assay.

Alkylation of the di-*t*-Boc-5-bromopyridin-2-yl hydrazine^{4c} with *t*-butylbromoacetate followed by deprotection with trifluoroacetic acid gave the desired chelate in 90% yield. This ligand formed the neutral tridentate complex **1** in 87% yield from tricarbonyl–Re(I) in aqueous ethanol.⁵ Single crystals of **1** suitable for X-ray crystallography were grown by slow evaporation from ethyl acetate. The structure shown in Figure 1 exhibits an undistorted octahedral coordination geometry with tridentate facial orientation of pyridyl, hydrazine-*N'*, and carboxylate groups.

Sonogashira coupling of the protected chelate with 17 α -ethynyl-estradiol, followed by deprotection, and Re(CO)₃⁺ labeling gave **2** in 95% yield. The complex was characterized by HPLC-MS ESI-negative detection as two chelate diastereomers (*m/z* 730, 9.55, and 10.50 min) as expected from the facial coordination geometry, but they were not isolated separately. Stereospecific hydrogenation of **2** using Lindlar's catalyst produced complex **3** containing a (*Z*)-ethene linkage in 70% yield. Hydrogenation of **2** with Pd/C gave the saturated alkane-linked complex **4** in 98% yield.

The receptor binding affinities of **2–4** for full length human ER α and ER β were determined by competitive radiometric assays with [³H]estradiol and expressed as relative binding affinity (RBA) compared with 17 β -estradiol (100%). The connecting linkage affected the binding affinity and selectivity of the complexes for ER α/β . The linear alkyne complex **2** exhibited high affinity and 4-fold greater selectivity for ER α over ER β . The (*Z*)-alkene

[†] New Mexico State University.

[‡] University of New Mexico Health Sciences Center.

[§] Department of Cell Biology and Physiology, University of New Mexico.

[#] Department of Pathology, University of New Mexico.

[¶] University of Illinois.




Figure 2. Fluorescence assays for GPR30 binding and activation with **2** and **3**. (a) GPR30 competitive binding of E2, 17 α -E2, **2**, and **3** with fluorescent estradiol derivative E2-Alexa using permeabilized COS7 cells. (b) Calcium mobilization assay for GPR30 and ER α in transfected COS7 cells versus concentration of E2, and complexes **2** and **3** (each at 10, 100, and 1000 nM).

Table 1. Structure and Relative Binding Affinity of Estradiol Complexes **2**, **3**, and **4** for ER α/β and GPR30

L	—C=C— 2	(Z)—CH=CH— 3	—CH ₂ CH ₂ — 4
ER α	20	10	1
ER β	5	24	0.5
GPR30	42	64	no data

complex **3** had greater affinity and 2-fold selectivity for ER β . These RBA values compare favorably with the best examples of reported estradiol tricarboxylate—Re(I) complexes.⁴ Complex **4** had low affinity for both ER α/β that can be attributed to unfavorable steric and entropic effects of the flexible alkyl linkage with receptor binding. These results indicate that relatively minor changes in the linkage can affect binding affinity and selectivity and could be optimized to increase ER subtype selectivity as demonstrated for selective estrogen receptor modulators.⁶

The GPR30 binding affinity of **2** and **3** was determined using a fluorescent Alexa633-labeled conjugate of estradiol that binds to GPR30 and ER α/β .^{2c} Binding affinity was assessed by flow cytometry using GPR30-transfected COS7 cells that do not express endogenous GPR30 or ER α/β . The cells were permeabilized with saponin to enable access of the charged fluorescent estrogen. Competitive binding was evaluated using 17 β -estradiol as a positive control for specific binding of the fluorescent estrogen to GPR30 (Figure 2a), and the relative K_i values of **2** and **3** are reported in Table 1. The nonbinding diastereomer 17 α -estradiol was included as a negative control. Both complexes **2** and **3** exhibited strong binding to GPR30 and demonstrate the potential for targeting this receptor.

To assess receptor binding in whole cells, we employed a functional assay based on the rapid receptor-mediated mobilization of intracellular calcium elicited by estrogen ligand binding to ER α and GPR30. Transfected COS7 cells expressing ER α or GPR30 were incubated at room temperature with the calcium-responsive fluorescent dye indo1-AM. Dose-dependent fluorescence was measured at 10, 100, and 1000 nM extracellular concentrations of E2, **2**, and **3** (Figure 2b). Alkyne complex **2** initiated a rapid increase in intracellular calcium concentrations with both ER α and GPR30. The (Z)-alkene complex **3** produced lower calcium levels during the same time period. These results demonstrate that complexes **2** and **3** are able to penetrate the cell membrane, bind, and activate the targeted estrogen receptors. Cell permeability, a critical

parameter affecting the kinetics of tissue uptake and overall biodistribution, can be assessed using this type of functional assay.

Convenient methodology for labeling with [^{99m}Tc(CO)₃(H₂O)₃]⁺ has been developed, and the IsoLink kit from Mallinckrodt reliably produced the aquo complex with radiochemical purity $\geq 98\%$ ($n \geq 20$).⁷ Due to the acid sensitivity of the tertiary propargylic 17- β -alcohol of the estradiol chelate, the alkaline mixture was neutralized with acetic acid rather than HCl. ^{99m}Tc-Labeling was conducted at ambient temperature for 2 h. Solid-phase extraction using a C-18 Sep-Pak with 40% EtOH/H₂O effectively removed excess ligand, and the ^{99m}Tc complex **2** eluted with EtOH in radiochemical yields ranging from 90 to 95% of the total loaded radioactivity. The radiochemical purity assessed by HPLC was $\geq 85\%$ ($n \geq 10$), and chromatographic resolution of the two diastereoisomeric chelates was observed by radiometric detection. The specific activity determined post-purification was 40 mCi/ μ g. HPLC and ITLC analyses demonstrated no significant degradation, reoxidation, or “leak” of ^{99m}Tc from the chelate after 48 h of storage at 4, 25, and 37 °C. The complex exhibited a log *P* octanol/water coefficient of 3.87 ± 0.49 ($n = 4$) determined by shake flask method that corresponds closely with estradiol log *P*^{ow} = 4.01. The stability of ^{99m}Tc-**2** was also evaluated in the presence of a 100-fold excess of histidine (1 mM). The radiochemical purity accessed via HPLC after 24 h incubation at 37 °C was $76.32 \pm 2.81\%$ ($n = 3$) with less than 1% reoxidation of the ^{99m}Tc species.

These results demonstrate that chelates **2** and **3** interact with receptors GPR30 and ER α/β in whole cells and suggest increased use of in vitro assays may facilitate the development of targeted imaging agents for intracellular receptors.

Acknowledgment. NIH/SCORE GM08136 (J.B.A.), CA116662, and UNM CRTC Translational Research Award (E.R.P.), CA25836 (J.A.K.), Cowboys for Cancer Research Foundation. WM Keck Foundation and NIH Roadmap U54MH074425.

Supporting Information Available: Experimental and spectral data provided as pdf files, X-ray structure of **2** provided as a cif file. This material is available free of charge via the Internet at <http://pubs.acs.org>.

References

- (a) Losel, R. M.; Falkenstein, E.; Feuring, M.; Schultz, A.; Tillmann, H.-C.; Rossol-Haseroth, K.; Wehling, M. *Physiol. Rev.* **2003**, *83*, 965–1016. (b) Osborne, C. K.; Schiff, R. *J. Clin. Oncol.* **2005**, *23*, 1616–1622. (c) Ariazi, E. A.; Ariazi, J. L.; Cordera, F.; Jordan, V. C. *Curr. Top. Med. Chem.* **2006**, *6*, 181–202.
- (a) Filardo, E. J.; Quinn, J. A.; Bland, K. I.; Frackelton, A. R. *Mol. Endocrinol.* **2000**, *14*, 1649–1660. (b) Thomas, P.; Pang, Y.; Filardo, E. J.; Dong, J. *Endocrinology* **2005**, *146*, 624–632. (c) Revankar, C. M.; Cimino, D. F.; Sklar, L. A.; Arterburn, J. B.; Prossnitz, E. R. *Science* **2005**, *307*, 1625–1630.
- Prossnitz, E. R.; Arterburn, J. B.; Edwards, B. S.; Sklar, L. A.; Oprea, T. I. *Exp. Opin. Drug Discovery* **2006**, *1*, 137–150.
- (a) Top, S.; El Hafa, H.; Vessieres, A.; Quivy, J.; Vaissermann, J.; Hughes, D. W.; Mcglinchey, M. J.; Mornon, J. P.; Thoreau, E.; Jaouen, G. *J. Am. Chem. Soc.* **1995**, *117*, 8372–8380. (b) Luyt, L. G.; Bigott, H. M.; Welch, M. J.; Katzenellenbogen, J. A. *Bioorg. Med. Chem. Lett.* **2003**, *11*, 4977–4989. (c) Arterburn, J. B.; Corona, C.; Rao, K. V. *J. Org. Chem.* **2003**, *68*, 7063–7070. (d) Top, S.; Boubekeur, L.; Jaouen, G.; Mundwiler, S.; Spingler, B.; Alberto, R. *Eur. J. Inorg. Chem.* **2004**, 2013–2017. (e) Bigott, H. M.; Parent, E.; Luyt, L. G.; Katzenellenbogen, J. A.; Welch, M. J. *Bioconjugate Chem.* **2005**, *16*, 255–264.
- Lazarova, N.; James, S.; Babich, J.; Zubieta, J. *Inorg. Chem. Commun.* **2004**, *7*, 1023–1026.
- Manas, E. S.; Unwalla, R. J.; Xu, Z. B.; Malamas, M. S.; Miller, C. P.; Harris, H. A.; Hsiao, C.; Akopian, T.; Hum, W. T.; Malakian, K.; Wolfson, S.; Bapat, A.; Bhat, R. A.; Stahl, M. L.; Somers, W. S.; Alvarez, J. C. *J. Am. Chem. Soc.* **2004**, *126*, 15106–15119.
- (a) Alberto, R.; Schibli, R.; Egli, A.; Schubiger, A. P.; Abram, U.; Kaden, T. A. *J. Am. Chem. Soc.* **1998**, *120*, 7987–7988. (b) Schibli, R.; Schubiger, P. A. *Eur. J. Nucl. Med. Mol. Imaging* **2002**, *29*, 1529–1542. (c) Stichelberger, A.; Waibel, R.; Dumas, C.; Schubiger, P. A.; Schibli, R. *Nucl. Med. Biol.* **2003**, *30*, 465–470.

JA066360P