


## Quantitative Microarray Profiling of DNA-Binding Molecules

James W. Puckett, Katy A. Muzikar, Josh Tietjen, Christopher L. Warren, Aseem Z. Ansari, and Peter B. Dervan

*J. Am. Chem. Soc.*, **2007**, 129 (40), 12310-12319 • DOI: 10.1021/ja0744899 • Publication Date (Web): 19 September 2007

Downloaded from <http://pubs.acs.org> on February 14, 2009



### More About This Article

Additional resources and features associated with this article are available within the HTML version:

- Supporting Information
- Links to the 2 articles that cite this article, as of the time of this article download
- Access to high resolution figures
- Links to articles and content related to this article
- Copyright permission to reproduce figures and/or text from this article

[View the Full Text HTML](#)

## Quantitative Microarray Profiling of DNA-Binding Molecules

James W. Puckett,<sup>†</sup> Katy A. Muzikar,<sup>†</sup> Josh Tietjen,<sup>‡</sup> Christopher L. Warren,<sup>‡</sup>  
Aseem Z. Ansari,<sup>\*,‡</sup> and Peter B. Dervan<sup>\*,†</sup>

Contribution from the Division of Chemistry and Chemical Engineering, California Institute of Technology, Pasadena, California 91125, and Department of Biochemistry and Genome Center, University of Wisconsin, Madison, Wisconsin 53706

Received July 3, 2007; E-mail: dervan@caltech.edu; ansari@biochem.wisc.edu

**Abstract:** A high-throughput Cognate Site Identity (CSI) microarray platform interrogating all 524 800 10-base pair variable sites is correlated to quantitative DNase I footprinting data of DNA binding pyrrole-imidazole polyamides. An eight-ring hairpin polyamide programmed to target the 5 bp sequence 5'-TACGT-3' within the hypoxia response element (HRE) yielded a CSI microarray-derived sequence motif of 5'-WWACGT-3' (W = A,T). A linear  $\beta$ -linked polyamide programmed to target a (GAA)<sub>3</sub> repeat yielded a CSI microarray-derived sequence motif of 5'-AARAARWWG-3' (R = G,A). Quantitative DNase I footprinting of selected sequences from each microarray experiment enabled quantitative prediction of  $K_a$  values across the microarray intensity spectrum.

### Introduction

Cell-permeable small molecules which bind specific DNA sequences and are able to interfere with protein–DNA interfaces would be useful in modulating eukaryotic gene expression. For targeting the regulatory elements of eukaryotic genes, knowledge of the preferred binding landscape of the ligand and the energetics of each site would guide gene regulation studies. Pyrrole–imidazole polyamides are a class of cell permeable oligomers which can be programmed, based on simple aromatic amino acid pairing rules, to bind a broad repertoire of DNA sequences.<sup>1</sup> Knowledge of polyamide match sites has allowed us to pursue the characterization of the equilibrium association constants and, hence, free energies of hairpin polyamides for cognate DNA sites by quantitative footprint titration methods. Despite the predictive power of simple pairing rules, the sequence dependent variability of DNA minor groove shape affords significant variability in the range of affinities for match as well as all formal single and double base pair mismatch sites.<sup>1</sup>

**Quantitative Footprint Titrations.** Characterization of polyamide binding preferences has been studied using quantitative DNase I footprinting titrations, affording binding isotherms that enable rigorous determination of the equilibrium association constant,  $K_a$ .<sup>2</sup> The resolution of footprinting is conservatively limited to association constants of 2-fold difference or greater. Polyamide binding preferences have frequently been interrogated using DNA fragments roughly 100 bp in size containing as many as four 6–10 bp binding sites, which are identical with the

exception of a single position that iteratively exhibits A•T, T•A, C•G, and G•C base pairs. Each binding site is interspersed with an 8 or more base pair spacer region to prevent interaction between the binding sites.<sup>3</sup> Obtaining high quality data limits a <sup>32</sup>P end-labeled DNA fragment to four unique binding sites due to the resolving power of a polyacrylamide gel in a quantitative footprint titration. While DNase I footprinting has enabled the elucidation of a binding code for hairpin polyamides, a relatively limited set of binding sites has been studied. To comprehensively interrogate all four encoded positions of an eight-ring hairpin polyamide, one would need 136 unique binding sites. In addition, interrogation of the base pairs flanking the polyamide core would necessitate 2080 (for 6 bp total) or 32 896 (for 8 bp total) binding sites.

**CSI Microarray Platform.** Several high-throughput platforms have been developed to characterize the binding properties of ligand–DNA interactions.<sup>4</sup> Of these, two have been used to explicitly study the binding preferences of polyamides. The fluorescence intercalator displacement assay has interrogated polyamide binding to 512 unique 5 bp sequences in a microplate format.<sup>4b</sup> The more recently developed cognate site identifier (CSI) microarray platform presents all 32 896 unique eight-mers (scalable to all unique ten-mers) to fluorescently labeled polyamides, enabling an unbiased interrogation of binding preference.<sup>4c</sup> By coupling DNase I footprinting with the CSI microarray data, the binding affinities ( $K_a$  values) of DNA-


<sup>†</sup> California Institute of Technology.

<sup>‡</sup> University of Wisconsin, Madison.

(1) Im/Py targets G•C; Py/Py targets A•T and T•A; and Ct/Py targets T•A. (a) Dervan, P. B.; Edelson, B. S. *Curr. Opin. Struct. Biol.* **2003**, *13*, 284–299. (b) Hsu, C. F.; Phillips, J. W.; Trauger, J. W.; Farkas, M. E.; Belitsky, J. M.; Heckel, A.; Olenyuk, B. Z.; Puckett, J. W.; Wang, C. C. C.; Dervan, P. B. *Tetrahedron* **2007**, *63*, 6146–6151.  
(2) Trauger, J. W.; Dervan, P. B. *Methods Enzymol.* **2001**, *340*, 450–466.

(3) (a) Doss, R. M.; Marques, M. A.; Foister, S.; Chenoweth, D. M.; Dervan, P. B. *J. Am. Chem. Soc.* **2006**, *128*, 9074–9079. (b) Marques, M. A.; Doss, R. M.; Foister, S.; Dervan, P. B. *J. Am. Chem. Soc.* **2004**, *126*, 10339–10349. (c) Foister, S.; Marques, M. A.; Doss, R. M.; Dervan, P. B. *Bioorg. Med. Chem.* **2003**, *11*, 4333–4340.

(4) (a) For a review on Protein Binding Microarrays (PBMs), see: Bulyk, M. L. *Methods Enzymol.* **2006**, *410*, 279–299. (b) For a review on fluorescence intercalator displacement (FID) assays, see: Tse, W. C.; Boger, D. L. *Acc. Chem. Res.* **2004**, *37*, 61–69. (c) For the initial report of cognate site identifier (CSI) microarrays, see: Warren, C. L.; Kratochvil, N. C. S.; Hauschild, K. E.; Foister, S.; Brezinski, M. L.; Dervan, P. B.; Phillips, G. N.; Ansari, A. Z. *Proc. Natl. Acad. Sci. U.S.A.* **2006**, *103*, 867–872.



**Figure 1.** (a) Quantitative DNase I footprinting gives rise to a defined equilibrium association constant at a specified binding site for a given DNA binding molecule. (b) The CSI microarray platform gives rise to relative binding preferences of an entire sequence space for the same molecule with a sequence logo as a standard summary output.

binding molecules for a significantly larger number of DNA sequences could be determined (Figure 1). To date, CSI microarray intensities of hairpin polyamide–Cy3 conjugates have been linearly correlated to the  $K_a$  values of unlabeled polyamides.<sup>4c</sup> We will examine whether this relationship between DNase I footprint titration-derived  $K_a$  values for Cy3-labeled polyamides and the corresponding microarray data remains true for additional polyamide binding architectures. Because the Cy3–polyamide conjugate may alter sequence specificity when compared with its biologically active counterpart, the sequence specificities of fluorophore-labeled polyamide and the biologically relevant polyamide will also be determined.

A CSI microarray harbors immense sequence specificity data; determining how to best represent this data is critical. The first reported CSI work<sup>4c</sup> represented binding preferences as a sequence logo<sup>5</sup> derived from several motif-finding algorithms<sup>6</sup> that searched the highest Z-score bins (the ~300 highest intensities on the array), assigning equal weight to each sequence. It also examined the relative abundance of each sequence motif mutation within its respective Z-score bin.<sup>4c</sup> In this paper we observe that  $K_a$ -weighting sequence motifs does not alter the sequence logo appreciably. In addition, a comprehensive single base pair mutational analysis is performed, which quantifies the specificities encoded by the polyamide at each position the polyamide interacts with DNA.

Two Cy3-labeled polyamides of biological interest<sup>7</sup> are examined on a CSI microarray that displays all unique 10 base pair DNA sequences. These polyamides include a hairpin

structure whose sequence specificities can be predicted from the extensive DNase I footprinting data characterizing other pyrrole–imidazole polyamides<sup>1</sup> and a linear  $\beta$ -linked structure whose sequence specificity is less well understood.<sup>8</sup> In order to correlate the CSI relative affinities (intensities) to absolute affinities ( $K_a$  values), DNase I footprinting was performed on a subset of these sequences for both the Cy3–polyamide conjugates and the related, unlabeled polyamides of known biological activity.

## Results and Discussion


**Polyamide Design.** Two polyamide core sequences have been chosen as representative of both hairpin and linear  $\beta$ -linked polyamide architectures. These core recognition sequences exhibit biologically significant roles, modulating transcription in cell culture experiments.<sup>7</sup> Hairpin polyamides **1** and **2** (Figure 2) were selected based on results from a project in which a polyamide–fluorescein conjugate, Ct-Py-Py-Im-(R)-<sup>H2N</sup>- $\gamma$ -Py-Im-Py-Py-Dp-FITC (**1**) displaced hypoxia inducible factor-1 $\alpha$  (HIF-1 $\alpha$ ) from the hypoxia response element (HRE) of the vascular endothelial growth factor (VEGF) gene, downregulating VEGF expression 60% in cell culture experiments.<sup>7a,b</sup> This eight-ring hairpin was programmed to bind the sequence 5'-WTWCGW-3' (W = A,T).<sup>1,3c</sup> In particular, polyamide **1** was shown to bind the HRE sequence, 5'-TACGTG-3', on the VEGF promoter by footprint titration.<sup>7a,b</sup> The Cy3 moiety was conjugated (**2**) at the same position as fluorescein for **1** to best mimic the binding properties between the two polyamides.

As with polyamide **1**, polyamide **3** (Figure 2) is known to bind its biologically relevant target. Polyamide **3**, Im-Py- $\beta$ -Im-

(5) Schneider, T. D.; Stephens, R. M. *Nucleic Acids Res.* **1990**, *18*, 6097–6100.

(6) (a) Bailey, T. L.; Elkan, C. In *Proceedings of the Second International Conference on Intelligent Systems for Molecular Biology*; Altman, R., Brutlag, D., Karp, P., Lathrop, R., Searls, D., Eds.; AAAI Press: Menlo Park, 1994, pp 28–36. (b) Liu, X. S.; Brutlag, D. L.; Liu, J. S. *Nat. Biotechnol.* **2002**, *20*, 835–839. (c) Hughes, J. D.; Estep, P. W.; Tavazoie, S.; Church, G. M. *J. Mol. Biol.* **2000**, *296*, 1205–1214.

(7) (a) Olenyuk, B. Z.; Zhang, G. J.; Klco, J. M.; Nickols, N. G.; Kaelin, W. G.; Dervan, P. B. *Proc. Natl. Acad. Sci. U.S.A.* **2004**, *101*, 16768–16773. (b) Nickols, N. G.; Jacobs, C. S.; Farkas, M. E.; Dervan, P. B. *Nucleic Acids Res.* **2007**, *35*, 363–370. (c) Burnett, R.; Melander, C.; Puckett, J. W.; Son, L. S.; Wells, R. D.; Dervan, P. B.; Gottesfeld, J. M. *Proc. Natl. Acad. Sci. U.S.A.* **2006**, *103*, 11497–11502.



**Figure 2.** Hairpin polyamides **1** and **2** targeted to the hypoxia response element (HRE), 5'-TACGTG-3'. Linear  $\beta$ -linked polyamides **3** and **4** targeted to GAA repeats in Friedrich's Ataxia.

Py- $\beta$ -Im- $\beta$ -Dp, targets an intronic 5'-(GAA)<sub>n</sub>-3' repeat hyperexpansion, enabling 2.5-fold upregulation of the *frataxin* gene, whose deficiency causes the neurodegenerative disorder Friedrich's Ataxia.<sup>7c</sup> Limited knowledge about the linear  $\beta$ -linked class of polyamides<sup>8</sup> precludes the existence of binding rules. The linear  $\beta$ -linked architecture has the added complexity of binding in 1:1 and 2:1 ligand/DNA stoichiometries, and we would anticipate that this class will be generally less useful due to sequence promiscuity resulting from multiple binding modes. Its 1:1 binding preferences for purine tracts, such as (GAA)<sub>n</sub>, likely reflect shape selectivity for sequences with narrow DNA minor groove conformations.<sup>8c</sup> In a 2:1 binding stoichiometry, polyamide **3** would be predicted to target 5'-WGCWGCWGCW-3'.<sup>8a</sup> Remarkably, relatively few genes are affected from cell culture studies of **3** suggesting that this polyamide may be specific for 5'-AAGAAGAAG-3'.<sup>7c</sup> The Cy3 fluorophore has been conjugated to the C-terminal 3,3'-diamino-*N*-methylpropylamine tail (polyamide **4**).

**CSI Microarray Design and Results.** CSI microarrays were synthesized using maskless array synthesis (MAS) technology<sup>9</sup> to display all 524 800 unique 10-base pair sites in quadruplicate across six microarrays. Replicates of individual hairpins occur on separate microarrays. Each hairpin on the chip consists of a self-complementary palindromic sequence interrupted by a central 5'-GGA-3' sequence to facilitate hairpin formation: 5'-GCGC-N<sup>1</sup>N<sup>2</sup>N<sup>3</sup>N<sup>4</sup>N<sup>5</sup>N<sup>6</sup>N<sup>7</sup>N<sup>8</sup>N<sup>9</sup>N<sup>10</sup>-GCGC-GGA-GCGC-N<sup>10</sup>N<sup>9</sup>N<sup>8</sup>N<sup>7</sup>N<sup>6</sup>N<sup>5</sup>N<sup>4</sup>N<sup>3</sup>N<sup>2</sup>N<sup>1</sup>-GCGC-3' (N = A,T,C,G). Previous experiments have found that 95% of the oligonucleotides on the array form duplexes.<sup>4c</sup>

Polyamides **2** and **4** were slowly titrated onto the arrays and imaged at each concentration until saturation of the highest

intensity binding sites was observed, 10 nM and 175 nM concentrations, respectively, for **2** and **4**. After each small addition of polyamide, the arrays were washed prior to imaging. The data for each of the arrays were then normalized as previously described<sup>4c</sup> to give averaged sequence intensities of the 524 800 10-base pair sites for **2** and **4**. As found with previously reported CSI arrays,<sup>4c</sup> histograms of the probe intensities for **2** and **4** display a strong right-handed tail (Supporting Information Figure 1). The fractional standard deviations among probe replicates (standard deviation of replicates/average normalized intensity) average  $0.15 \pm 0.09$  (polyamides **2** and **4**), for intensities exceeding  $1 \times 10^3$  (Supporting Information Figure 2).


**Plasmid Design.** Three plasmids have been designed based on output from the CSI microarray intensities (Figure 3). Because of our interest in testing the dynamic range of the CSI assay in terms of the representative  $K_a$  values measured by a broad range of intensities, plasmids pKAM3 and pJWP17 were constructed to harbor binding sites of equal intensity spacing across a broad portion of each array's intensities, between highest and lowest intensities. The  $K_a$  values found using pKAM3 were clustered across the three highest intensities, necessitating further interrogation. Plasmid pKAM4 was designed to probe three additional intensities. A single binding site (**IIIa** and **IIb**) was held constant between pKAM3 and pKAM4 to enable interplasmid comparison of binding affinities. Because pJWP17 afforded  $K_a$  values broadly spaced across the intensity spectrum, no further study was pursued.

Since our goal is to directly compare footprinting-derived  $K_a$  values with CSI-array derived intensities, each plasmid binding site mimics the full 10 base pair binding site from the array in addition to two flanking base pairs on either side of the binding site: 5'-GC-(N)<sub>10</sub>-GC-3' (N = A,T,C,G). Attempts to fully replicate the 5'-GCGC-(N)<sub>10</sub>-GCGC-3' binding site from the array exhibited secondary structure formation when the respective amplicons were sequenced and separated by denaturing gel electrophoresis.

(8) (a) Dervan, P. B.; Urbach, A. R. In *Essays in Contemporary Chemistry*; Quinkert, G., Kisakürek, M. V., Eds.; Verlag Helvetica Chimica Acta: Zurich, 2000; pp 327–339. (b) Urbach, A. R.; Dervan, P. B. *Proc. Natl. Acad. Sci. U.S.A.* **2001**, *98*, 4343–4348. (c) Urbach, A. R.; Love, J. J.; Ross, S. A.; Dervan, P. B. *J. Mol. Biol.* **2002**, *320*, 55–71. (d) Marques, M. A.; Doss, R. M.; Urbach, A. R.; Dervan, P. B. *Helv. Chim. Acta* **2002**, *85*, 4485–4517.

(9) Singh-Gasson, S.; Green, R. D.; Yue, Y. J.; Nelson, C.; Blattner, F.; Sussman, M. R.; Cerrina, F. *Nat. Biotechnol.* **1999**, *17*, 974–978.





**Figure 3.** Insert sequences utilized for plasmids, with binding sites boxed, labeled with their corresponding CSI array intensity, and numbered. (a) pKAM3 is shown, in addition to a microarray schematic demonstrating the relationship between the plasmid and a selected microarray sequence. (b) pKAM4. (c) pJWP17.

**Table 1.** Quantitative DNase I Footprinting Derived  $K_a$  Values ( $M^{-1}$ ) for Polyamides 1 and 2, Their 10 Base Pair Binding Sites, and the Corresponding CSI Microarray Intensity<sup>a</sup>

a) pKAM-3		Ia	IIa	IIIa	IVa
Polyamide		TTTTACGTAA	TTTTACGTAG	TTTTACGTGA	TTTTACGGAA
1		$7.5 (\pm 1.8) \times 10^9$ [1]	$5.1 (\pm 0.6) \times 10^9$ [1.5]	$4.2 (\pm 0.6) \times 10^9$ [1.8]	$1.5 (\pm 0.8) \times 10^8$ [50]
2		$4.5 (\pm 1.0) \times 10^9$ [1]	$3.0 (\pm 0.6) \times 10^9$ [1.5]	$2.1 (\pm 0.3) \times 10^9$ [2.1]	$6.2 (\pm 2.0) \times 10^7$ [73]
CSI Intensity ( $\times 10^3$ )		75.6 ( $\pm 9.9$ )	51.4 ( $\pm 7.4$ )	31.3 ( $\pm 4.8$ )	4.2 ( $\pm 1.4$ )
b) pKAM-4		Ib	IIb	IIIb	IVb
Polyamide		TTTTACGTGA	AATTTTCGTGT	GCTTTTCGTCC	ACCTTCGTGA
1		$5.4 (\pm 0.9) \times 10^9$ [1.3]	$2.3 (\pm 0.1) \times 10^9$ [3.2]	$2.8 (\pm 0.2) \times 10^9$ [2.6]	$1.5 (\pm 0.2) \times 10^9$ [5]
2		$1.6 (\pm 0.2) \times 10^9$ [2.8]	$4.0 (\pm 0.9) \times 10^8$ [11]	$5.8 (\pm 0.7) \times 10^8$ [7.8]	$1.3 (\pm 0.2) \times 10^8$ [34]
CSI Intensity ( $\times 10^3$ )		31.3 ( $\pm 4.8$ )	20.0 ( $\pm 2.8$ )	12.0 ( $\pm 1.5$ )	6.0 ( $\pm 0.4$ )

<sup>a</sup> All footprinting incubations were conducted at a minimum in triplicate at 23 °C for 14 h. Standard deviations are shown in parentheses. The bracketed numbers are  $K_{a-max}/K_{a-current}$  to compare  $K_a$  values within each polyamide series.

**Quantitative DNase I Footprint Titrations: Affinity and Specificity Determination.** Hairpin polyamides 1 and 2 were incubated each for 14 h with pKAM3 or pKAM4 prior to DNase I cleavage. These two polyamides were found to bind each of seven unique 10-base pair binding sites in the same rank order, preferentially binding 5'-TTTTACGTAA-3' with affinities of  $7.5 \times 10^9 M^{-1}$  (1) and  $4.5 \times 10^9 M^{-1}$  (2) (Figure 4 and Table 1).


Replacing the fluorescein dye on polyamide 1 with Cy3 (polyamide 2) introduced an energetic penalty that ranged from 1.5- to 10-fold, with the minimum penalty occurring at the two highest CSI intensity binding sites (Table 1). Polyamide 2 differentiated the highest and lowest affinity binding sites by

70-fold, slightly more than the 50-fold differentiation found for the fluorescein-labeled polyamide 1.

Linear  $\beta$ -linked polyamides 3 and 4 were each incubated for 14 h with pJWP17 prior to DNase I cleavage. They bound four unique 10-base pair sites in the same rank order, preferentially binding 5'-AAGAAGAAGT-3' (Table 2 and Figure 5).

Appending the Cy3 dye to polyamide 3 had either no effect on affinity or reduced binding affinity as much as 30-fold (Table 2). Polyamide 3 bound all four binding sites over a 2400-fold range in affinity, eight times broader than that for polyamide 4.

**Calibrating Microarrays for  $K_a$  Prediction.** Because DNase I footprinting enables the calculation of  $K_a$  and the direct



**Figure 4.** DNase I footprinting gels and corresponding isotherms of polyamides **1** and **2** on pKAM3 and pKAM4. (a) Polyamide **1** on pKAM3. (b) Polyamide **2** on pKAM3. (c) Polyamide **1** on pKAM4. (d) Polyamide **2** on pKAM4.

**Table 2.** Quantitative DNase I Footprinting Derived  $K_a$  Values ( $M^{-1}$ ) for Polyamides **3** and **4**, Their 10 Base Pair Binding Sites, and the Corresponding CSI Microarray Intensity<sup>a</sup>

pJWP-17		Ic	IIc	IIIc	IVc
Polyamide		AAGAAGAAGT	AAGAAGTCA	ATGTTGTTGA	ATGAAGACGA
<b>3</b>	→◇●◇◇○◇◇●	$2.4 (\pm 0.6) \times 10^{10}$ [1]	$9.3 (\pm 2.3) \times 10^9$ [3]	$2.9 (\pm 0.7) \times 10^8$ [80]	$1.0 (\pm 0.4) \times 10^7$ [2400]
<b>4</b>	Cy3(+)-◇◇●◇◇○◇◇●	$3.3 (\pm 0.7) \times 10^9$ [1]	$2.7 (\pm 0.8) \times 10^8$ [10]	$1.1 (\pm 0.4) \times 10^8$ [30]	$1.0 (\pm 0.2) \times 10^7$ [330]
CSI Intensity ( $\times 10^3$ )		75.2 ( $\pm 9.2$ )	51.2 ( $\pm 6.2$ )	26.8 ( $\pm 7.3$ )	4.1 ( $\pm 0.4$ )

<sup>a</sup> All footprinting incubations were conducted at a minimum in triplicate at 23 °C for 14 h. Standard deviations are shown in parentheses. The bracketed numbers are  $K_{a-max}/K_{a-current}$  to compare  $K_a$  values within each row.

comparison of four binding sites in a single assay, determining energetics data from CSI microarrays is crucial for understanding the global binding specificity of a polyamide. An eight-ring hairpin polyamide targeting 5'-WGWWCW-3' (W = A,T) and characterized by quantitative DNase I footprinting, Im-Py-Py-Py- $\gamma$ -Im-Py-Py- $\beta$ -Dp,<sup>3c,10</sup> has been compared to its Cy3-labeled counterpart studied on the CSI-array platform, demonstrating a linear relationship between intensity and  $K_a$ .<sup>4c</sup>

Because microarray intensity at a specific microarray feature should be proportional to the fractional occupancy of DNA at


that feature, the relationship between equilibrium association constant ( $K_a$ ) or dissociation constant ( $K_d$ ) and background-normalized microarray intensity should be<sup>11</sup>

$$\text{Intensity} = c \times \Theta = c \times \frac{K_a[\text{PA}]}{1 + K_a[\text{PA}]} = c \times \frac{[\text{PA}]}{K_d + [\text{PA}]} \quad (1)$$

In this relationship,  $\Theta$  represents the fractional occupancy of DNA at a specific feature,  $c$ , a scalar to reflect that microarray intensity can vary with incident laser intensity, and  $[\text{PA}]$ , the

(10) (a) Trauger, J. W.; Baird, E. E.; Dervan, P. B. *Nature* **1996**, *382*, 559–561. (b) Trauger, J. W. Ph.D. Thesis, California Institute of Technology, 1999.

(11) For derivation of these equations, see the Supporting Information for Bulyk, M. L.; Huang, X. H.; Choo, Y.; Church, G. M. *Proc. Natl. Acad. Sci. U.S.A.* **2001**, *98*, 7158–7163.



**Figure 5.** DNase I footprinting gels and corresponding isotherms of polyamides **3** and **4** on pJWP17.


free polyamide concentration on the CSI array. The terms  $c$  and  $[PA]$  are solved for a curve fit to eq 1 using  $K_a$  values derived from DNase I footprint titrations and CSI microarray intensity data. Examining the limiting case where  $[PA] \ll K_d$  one observes a simplification to eq 1:

$$\text{Intensity} = c \times \frac{[PA]}{K_d} = c \times [PA] \times K_a \quad (2)$$

Equation 2 represents the linear subset of the more general CSI intensity –  $K_a$  relationship described in eq 1. Fitting the footprinting data of polyamide **2** to its corresponding microarray

(12) (a)  $c \times [PA]$  was  $1.7 \times 10^{-5}$  for polyamide **2**. (b)  $c$  was  $80.2 \times 10^3$  and  $[PA]$  was  $5.5 \times 10^{-9}$  M for polyamide **4**. (c) To view plots reflecting the same curve fits of Figure 6 on a log–log scale, please see Supporting Information Figure 3.

(13) Although the data for polyamide **2** (Figure 6a) maps intensity and  $K_a$  values using the linearized equation 2, this fit is distinct from that obtained by fitting the data to a line of the form  $y = mx + b$ , which includes an intensity-axis intercept term. While very small in this case, the differences in the slopes and intercepts of the lines may indicate error both in the background correction of the microarray and in the DNase I footprinting data. To correct for this possibility, we propose the use of an error term,  $\epsilon$ , that would modify eqs 1 and 2 to the following: Intensity =  $c \times \Theta + \epsilon = c \times \{K_a[PA]\} / \{1 + K_a[PA]\} + \epsilon = c \times \{[PA]\} / \{K_d + [PA]\} + \epsilon$  (eq 1e) and Intensity =  $c \times [PA] / K_d + \epsilon = c \times [PA] \times K_a + \epsilon$  (eq 2e). When fitting the intensity and  $K_a$  data for polyamide **2** to the modified equation 2e, one finds a marginally improved fit ( $R^2 = 0.97$ ), although the curve fit for polyamide **4** using equation 1e is unimproved ( $R^2 = 0.99$ ). For polyamide **2**,  $c \times [PA] = 1.5 \times 10^{-5}$  and  $\epsilon = 5.5 \times 10^3$ . For polyamide **4**,  $c = 81.2 \times 10^3$ ,  $[PA] = 5.7 \times 10^{-9}$  M, and  $\epsilon = -1.1 \times 10^3$ .



**Figure 6.** CSI array intensities correlate well with DNase I footprinting-determined  $K_a$  values. (a) Polyamide **2** vs CSI array fit to eq 2. (b) Polyamide **4** vs CSI array fit to eq 1.

intensities (Table 1) using eq 2 fits well ( $R^2 = 0.94$ ). The linearized eq 2 does not, however, map intensity and  $K_a$  with high correlation for polyamide **4**. Fitting the data to eq 1 affords a significantly better fit ( $R^2 = 0.99$ ), indicating that  $[PA]$  is not insignificant relative to the  $K_d$  of the highest intensity microarray data (Figure 6).<sup>12,13</sup>

The  $K_a$ -calibrated microarrays can subsequently be used to interpolate  $K_a$  values from normalized sequence intensities.  $K_a$  values are derived by rearranging eq 1 to present  $K_a$  as a function of microarray intensity:

$$K_a = \frac{\text{Intensity}}{[PA] \times (c - \text{Intensity})} \quad (3)$$


In the case where  $[PA] \ll K_d$ , eq 2 is rearranged to

$$K_a = \frac{\text{Intensity}}{[PA] \times c} \quad (4)$$


**Correlating Binding Between Cy3-Labeled and Biologically Relevant Polyamides.** While establishing a general  $K_a$ –intensity relationship for Cy3-labeled polyamides is a crucial first step toward global sequence interrogation of a core polyamide motif, it is equally important that the biologically relevant polyamide has sequence preferences that correlate with its Cy3-labeled counterpart. Scatter plots of polyamide **1** vs **2** and polyamide **3** vs **4** are best fit by a power relationship of  $y = ax^n$ , where  $(x,y)$  denotes the  $K_a$  values for (**1**, **2**) or (**3**, **4**) (Figure 7).<sup>14</sup> The  $R^2$  between **1** and **2** is 0.87, and that between **3** and **4** is 0.78.

(14) For the relationship between polyamides **1** and **2**,  $a = 0.0253$  and  $n = 1.115$ . For the relationship between polyamides **3** and **4**,  $a = 349.83$  and  $n = 0.637$ .






**Figure 7.** (a) Correlation of  $K_a$  values for polyamide 1 (fluorescein labeled) and polyamide 2 (Cy3 labeled). (b) Correlation of  $K_a$  values for polyamide 3 (unlabeled) and polyamide 4 (Cy3 labeled).



**Figure 8.** Sequence logo for polyamide 2.



**Figure 9.** Sequence logo for polyamide 4.

**Sequence Analysis.** To graphically represent the binding preferences of polyamides 2 and 4, sequence logos have been generated (Figures 8 and 9).

In all cases, the motif finding program MEME<sup>6a</sup> was utilized to extract sequence motifs from the CSI binding intensities. The position specific probability matrices output by MEME were used as inputs to enoLOGOS<sup>15</sup> to generate a sequence logo.<sup>16</sup> The logo for polyamide 2 was created by searching the ~2500

highest sequence intensities of the CSI microarray.<sup>17</sup> These data points span approximately a 3-fold range in  $K_a$ . The logo for polyamide 4 interrogated the 48 highest intensity sequences (a 7-fold range in  $K_a$ ) of the CSI microarray.<sup>18</sup> We examined  $K_a$ -weighted sequence logos for both polyamides 2 and 4 and found minimal differences in the resulting logos (Supporting Information Figure 3).

The motif for polyamide 2 has the most information at a site width of six – 5'-WWACGT-3' (Figure 8; W = A,T). The chlorothiophene/pyrrole pair (Ct/Py) specificity cannot be globally elucidated using polyamide 2 because of the palindromic nature of the ACGT binding site core. It is evident that the core does specify 5'-ACG-3' using Py/Py, Py/Im, and Im/Py pairings, respectively. Polyamide 3 specifies 9 base pairs based on MPE footprinting data (unpublished). Polyamide 4 elicits a 9 bp motif that is best represented as 5'-AARAARWWG-3' (Figure 9; R = G,A and W = A,T). Previous work would suggest that Im may have no sequence preferences within linear  $\beta$ -linked polyamides,<sup>8</sup> although this selection of 9 bp high affinity binding sites for 4 suggests at least G•C or A•T specificity, consistent with microarray data from Friedrich's Ataxia cell culture work.<sup>7c</sup>

#### Quantitative Profiling of Single Base Pair Mismatches.

While sequence logos provide a visual representation of sequence specificity, traditional studies on polyamides quantitate the specificity of a ring pairing at a selected base pair. We have examined a comprehensive single base pair mutational analysis of both polyamides 2 and 4 using  $K_a$  values interpolated from the calibrated CSI microarrays (Tables 3 and 4).<sup>19</sup>

Because the motif finding algorithm MEME found 5'-WWACGT-3' (W = A,T) as a preferred binding sequence for polyamide 2, we utilized this core sequence for mutational studies. Additionally, because of the 5'-ACGT-3' palindromic element of this binding site, we have isolated only binding sites containing 5'-WWWWWWACGT-3' and their mutant counterparts to preclude analyzing variants where the polyamide may be rotated 180° from the presumed orientation. To determine a  $K_a$  for 5'-WWWWWWACGT-3' (for example), the geometric mean of all microarray binding sites containing this motif was found. Walking from 5' to 3' on 5'-W<sup>1</sup>W<sup>2</sup>A<sup>3</sup>C<sup>4</sup>G<sup>5</sup>T<sup>6</sup>-3', we observe that there is 3-fold specificity for W versus S (S = C,G) at position 1 (occupied by the linker). At position 2 (Ct/Py pair), there is 20-fold specificity for W versus S but minimal for T•A versus A•T. The previous study of Ct/Py specificity noted only modest specificity for T•A versus A•T.<sup>3c</sup> Position 3 (a Py/Py pair) confirms the previously observed W over S specificity.<sup>1</sup> At position 4 (a Py/Im pair) the polyamide encodes the greatest specificity with preference for C•G versus A•T, T•

(16) Figure 8 utilized 10 variable bases and contained a background GC content of 50%; Figure 9 utilized 10 variable bases and two fixed bases, each flanking the 5' and 3' portion of the variable region, and contained a background GC content of 58%. These background GC content corrections were utilized in the motif searching parameters.

(17) There are 1258 occurrences of a full 6 bp match sequence, TTAGCT. Double this number of highest intensity sequences was also searched yielding only modest changes in the data. The sequence logo is reported for the 2516 highest intensity sequences.







(18) There are 24 occurrences of a full 9 bp match sequence, AAGAAGAAG on the microarray. Double this number of highest intensity sequences was searched in addition to searching only the 24 highest intensities, yielding only small changes in the data. The sequence logo reported contains the 48 highest intensity sequences.

(19) To convert intensity to  $K_a$ , we have included the error term  $\epsilon$  in our calculations. This gives modified versions of eqs 3 and 4,  $K_a = (\text{Intensity} - \epsilon)/\{[\text{PA}] \times (c - \text{Intensity} + \epsilon)\}$  and  $K_a = (\text{Intensity} - \epsilon)/\{[\text{PA}] \times c\}$ , respectively.

(15) Workman, C. T.; Yin, Y. T.; Corcoran, D. L.; Ideker, T.; Stormo, G. D.; Benos, P. V. *Nucleic Acids Res.* **2005**, *33*, W389–W392.



**Table 3.** Microarray-Derived Binding Affinities and Specificities of All Single Base Pair Mismatch Sites for Polyamide 2<sup>a</sup>










Polyamide 2	X:Z	$K_a$ (M <sup>-1</sup> )
5'-W W W W <b>X</b> A C G T-3'  Cy3-(+)	A:T T:A C:G G:C	2.0 (1.4) × 10 <sup>9</sup> 2.5 (1.2) × 10 <sup>9</sup> 6.9 (1.4) × 10 <sup>8</sup> 6.8 (1.6) × 10 <sup>8</sup>
5'-W W W W W <b>X</b> A C G T-3'  Cy3-(+)	A:T T:A C:G G:C	1.8 (1.3) × 10 <sup>9</sup> 2.7 (1.2) × 10 <sup>9</sup> 1.0 (2.0) × 10 <sup>8</sup> 1.3 (2.2) × 10 <sup>8</sup>
5'-W W W W W W <b>X</b> C G T-3'  Cy3-(+)	A:T T:A C:G* G:C	2.2 (1.3) × 10 <sup>9</sup> 1.1 (1.6) × 10 <sup>9</sup> ≤ 10 <sup>8</sup> 1.3 (2.5) × 10 <sup>8</sup>
5'-W W W W W W A <b>X</b> G T-3'  Cy3-(+)	A:T* T:A* C:G G:C*	≤ 10 <sup>8</sup> ≤ 10 <sup>8</sup> 2.2 (1.3) × 10 <sup>9</sup> ≤ 10 <sup>8</sup>
5'-W W W W W W A C <b>X</b> T-3'  Cy3-(+)	A:T T:A C:G G:C	1.2 (1.4) × 10 <sup>9</sup> 2.9 (1.8) × 10 <sup>8</sup> 2.4 (1.8) × 10 <sup>8</sup> 2.2 (1.3) × 10 <sup>9</sup>
5'-W W W W W W A C G <b>X</b> -3'  Cy3-(+)	A:T T:A C:G* G:C*	1.3 (1.4) × 10 <sup>9</sup> 2.2 (1.3) × 10 <sup>9</sup> ≤ 10 <sup>8</sup> ≤ 10 <sup>8</sup>

<sup>a</sup> All  $K_a$  values are derived from the geometric average of all CSI binding site intensities on the array containing a specified sequence, converted to a  $K_a$  value using eq 4, corrected to include an error term  $\epsilon$ .<sup>13</sup> The values in parentheses are the geometric standard deviations for each  $K_a$  value. X:Z entries marked with a superscripted "\*" contain averaged intensities below  $\epsilon$ . For these entries, an upper bound on the  $K_a$  is estimated based on the log-log plot of  $K_a$  versus intensity found in Supporting Information Figure 3.

A, or G:C. It is likely that this preference is at least 20-fold. At position 5, polyamide 2 appears to exhibit less specificity than would be predicted for an Im/Py ring pair, binding almost as well to A:T as to G:C.<sup>1</sup> The polyamide "turn unit," position 6, confirms a strong preference for W over S.<sup>1</sup> Through this quantitative study, we observe four strongly encoded binding positions, italicized in 5'-WWWCGW-3'. The discrepancy between the observed sequence logo, as found by MEME, and the suggested specificity by a single base pair mutation study likely stems from (i) the examination of all sequences in the single base pair mutation as compared to only a subset for the sequence logo, (ii) the assumption by the logo of independence of base pair-polyamide interaction at each position, and (iii) the examination in the single base pair mutation of the average  $K_a$  of a group of sequences containing a specified motif.

In conjunction with the sequence logo for polyamide 2, the CSI array analysis validates the sequence specificity programmed by the aromatic amino acid ring pairs. The extensive DNase I footprinting data on eight-ring and six-ring hairpin polyamides, while limited on the scale of a CSI microarray, enabled the creation of pairing rules that are remarkably general.<sup>1</sup> It is evident from the microarray that Im/Py and Py/Im ring pairs offer the greatest specificity for a single base pair, while Py/Py, Ct/Py, and the "turn unit" afford general W specificity. While the Ct/Py ring pair conferred minimal specificity for T:A versus A:T, its W specificity is likely an improvement over the use of a Py/Py ring pair, which at the N-terminus of an eight-ring hairpin polyamide exhibits specificity for A:T, T:A,

**Table 4.** Microarray-Derived Binding Affinities and Specificities of All Single Base Pair Mismatch Sites for Polyamide 4<sup>a</sup>

Polyamide 4	X:Z	$K_a$ (M <sup>-1</sup> )
5'- <b>X</b> A R A A R W W G-3'  Cy3-(+)	A:T T:A C:G G:C	1.6 (2.2) × 10 <sup>8</sup> 8.7 (2.0) × 10 <sup>7</sup> 4.3 (1.8) × 10 <sup>7</sup> 4.7 (1.8) × 10 <sup>7</sup>
5'-A <b>X</b> R A A R W W G-3'  Cy3-(+)	A:T T:A C:G G:C	1.6 (2.2) × 10 <sup>8</sup> 7.7 (2.2) × 10 <sup>7</sup> 2.1 (1.8) × 10 <sup>7</sup> 3.2 (1.9) × 10 <sup>7</sup>
5'-A A <b>X</b> A A R W W G-3'  Cy3-(+)	A:T T:A C:G G:C	1.3 (2.0) × 10 <sup>8</sup> 3.5 (1.7) × 10 <sup>7</sup> 8.4 (1.6) × 10 <sup>7</sup> 2.0 (2.2) × 10 <sup>8</sup>
5'-A A R <b>X</b> A R W W G-3'  Cy3-(+)	A:T T:A C:G G:C	1.6 (2.2) × 10 <sup>8</sup> 4.3 (1.8) × 10 <sup>7</sup> 9.9 (1.9) × 10 <sup>6</sup> 7.5 (2.6) × 10 <sup>6</sup>
5'-A A R A <b>X</b> R W W G-3'  Cy3-(+)	A:T T:A C:G G:C	1.6 (2.2) × 10 <sup>8</sup> 8.3 (2.2) × 10 <sup>7</sup> 9.1 (2.1) × 10 <sup>6</sup> 1.1 (2.0) × 10 <sup>7</sup>
5'-A A R A A <b>X</b> W W G-3'  Cy3-(+)	A:T T:A C:G G:C	1.5 (2.1) × 10 <sup>8</sup> 5.5 (1.7) × 10 <sup>7</sup> 8.5 (1.7) × 10 <sup>7</sup> 1.7 (2.3) × 10 <sup>8</sup>
5'-A A R A A R <b>X</b> W G-3'  Cy3-(+)	A:T T:A C:G G:C	1.7 (2.2) × 10 <sup>8</sup> 1.5 (2.2) × 10 <sup>8</sup> 2.2 (2.2) × 10 <sup>7</sup> 2.0 (2.4) × 10 <sup>7</sup>
5'-A A R A A R W <b>X</b> G-3'  Cy3-(+)	A:T T:A C:G G:C	1.6 (2.5) × 10 <sup>8</sup> 1.5 (2.0) × 10 <sup>8</sup> 3.5 (2.0) × 10 <sup>7</sup> 3.9 (2.0) × 10 <sup>7</sup>
5'-A A R A A R W W <b>X</b> -3'  Cy3-(+)	A:T T:A C:G G:C	1.2 (1.9) × 10 <sup>8</sup> 1.1 (2.0) × 10 <sup>8</sup> 1.1 (1.7) × 10 <sup>8</sup> 1.6 (2.2) × 10 <sup>8</sup>

<sup>a</sup> All  $K_a$  values are derived from the geometric average of all CSI binding site intensities on the array containing a specified sequence, converted to a  $K_a$  value using eq 3, corrected to include an error term  $\epsilon$ .<sup>13</sup> The values in parentheses are the geometric standard deviations for each  $K_a$  value.

and G:C versus C:G.<sup>3c</sup> The sequence specificity of 2 correlates remarkably well with the 5'-ACGT-3' specificity of echinomyacin,<sup>20</sup> also known to affect VEGF expression in cell culture.<sup>21</sup>

The examination of polyamide 4 marks the most comprehensive sequence specificity study of a linear  $\beta$ -linked polyamide since the original examination of the binding specificity for Im- $\beta$ -Im-Py- $\beta$ -Im- $\beta$ -Im-Py- $\beta$ -Dp.<sup>8a,b</sup> In the 5'-A<sup>1</sup>A<sup>2</sup>R<sup>3</sup>A<sup>4</sup>A<sup>5</sup>R<sup>6</sup>W<sup>7</sup>W<sup>8</sup>G<sup>9</sup>-3' sequence (R = G,A; W = A,T), positions 4, 5, and 7, each containing either a Py or a  $\beta$ , exhibit the greatest specificity for W over S (S = C,G). Intriguingly, the  $\beta$  at position 4 prefers A:T over T:A, an unexpected specificity. The sequence logo for polyamide 4 indicates that Im has a modest preference for G:C or A:T over other base pairings; in this mutational study, however, imidazole is generally degenerate. The wide range of  $K_a$  values comprising

(20) Van Dyke, M. M.; Dervan, P. B. *Science* **1984**, 225, 1122–1127.

(21) Kong, D.; Park, E. J.; Stephen, A. G.; Calvani, M.; Cardellina, J. H.; Monks, A.; Fisher, R. J.; Shoemaker, R. H.; Melillo, G. *Cancer Res.* **2005**, 65, 9047–9055.

each motif (high geometric standard deviation) make the statistical significance of any specificities under 4 relatively small. In general, the geometric standard deviations for polyamide **4** were higher than those for polyamide **2**, when including only those table entries for polyamide **2** in which each  $K_a$  value was composed of all instances of the motif. One potential source of the increased standard deviation in binding affinities is the single variable base flanking the nine base pair binding site for polyamide **4**. Because the minor groove width is a potentially important contributor to binding affinity and specificity for the linear  $\beta$ -linked class of polyamides,<sup>8c</sup> a single variable, flanking base is unlikely to enable comprehensive interrogation of the global set of sequence-dependent DNA microstructures. As with polyamide **2**, the discrepancies observed between the sequence logo of polyamide **4** and the comprehensive single base pair mutational analysis likely stem from similar causes.

With the sequence logo (approximated as 5'-AARAAR-WWG-3') as a snapshot of the highest affinity binding sites for polyamide **4** ( $K_a \approx 5 \times 10^8$  to  $3.3 \times 10^9$  M<sup>-1</sup>) and the footprint titration binding isotherms for determining DNA binding mode, we confirm a preference for the 1:1 binding stoichiometry. Previous data characterizing the linear  $\beta$ -linked polyamide Im- $\beta$ -Im-Py- $\beta$ -Im- $\beta$ -Im-Py- $\beta$ -Dp demonstrated a 30-fold energetic preference for the 1:1 versus 2:1 binding stoichiometry, presumably due to the increased entropic cost of the 2:1 binding mode.<sup>8a</sup> It is remarkable that polyamide **3** exhibited specificity for upregulation of the *frataxin* gene in cell culture,<sup>7c</sup> since the sequence preference for **4** was not overwhelmingly 5'-AAGAA-GAAG-3'. Two possible explanations for this observation are (i) that multiple binding events in the genome have marginal effects on transcription and that the specificity is amplified by the GAA repeat expansion in Friedreich's Ataxia or (ii) that many of the sequences described by 5'-AARAARWWG-3' exist in higher order chromosomal structures that cannot be targeted by polyamide **3**.

**Suggestions for Microarray Usage.** In the case where the free ligand concentration is small relative to the  $K_d$  for each binding site on the CSI microarray, a linear  $K_a$ -intensity relationship is observed. The binding profiles examined for polyamide **2** and for previously studied molecules are examples of linear  $K_a$ -intensity relationships.<sup>4c</sup> For the highest intensity sites also studied by DNase I footprinting (Figure 6a), the CSI microarray experiment contains greater resolving power and can differentiate  $K_a$  values that are indistinguishable by quantitative DNase I footprint titrations. In this example, as CSI intensity data approach  $\epsilon$ , small changes in intensity yield large changes in predicted  $K_a$ . Because the characterization of DNA-binding ligands is most concerned with defining a perfect match site, this limitation is minor. CSI data for polyamide **2** conservatively enables distinguishing a 50-fold range of  $K_a$  values, thus encompassing the majority of single base pair mismatch specificities.

In the case where the free ligand concentration is comparable to the  $K_d$ , a nonlinear  $K_a$ -intensity relationship is observed. The binding profile for polyamide **4** marks an example of a CSI microarray studied compound that occurs outside the linear range of eq 1. In this case, clustered high-intensity data points can span a broad range of  $K_a$  values (Figure 6b). The error inherent to the CSI microarray analysis is thus amplified when  $K_a$  values in this high CSI intensity region are interpolated.

Because of the gradual polyamide titration onto the array, it should be possible to capture snapshots of both polyamide saturation within the linear  $K_a$ -intensity region for the highest affinity binding sites on the microarray and binding site saturation enabling lower intensity data points to fall within the higher precision linear  $K_a$ -intensity region. Such titration may enable high precision  $K_a$  data to be extracted from all intensities of the microarray.

The sequence logos presented in this paper represent a snapshot of a binding profile for the highest affinity binding sites by a dye-labeled ligand. The polyamide core dictates the majority of the binding specificity revealed by CSI microarray analysis — the presence of a Cy3 label may reduce affinity to a binding site relative to its unlabeled counterpart but does not alter the rank order of binding preferences. Complementing the graphical image of a sequence logo, the comprehensive single base pair mutational analysis afforded by the extensive microarray data quantitates one's understanding of the polyamide sequence preferences.

## Conclusion

Correlating the sequence preference landscape present on the CSI microarray to quantitative footprinting enables energetic studies using global binding information. This capacity marks a significant forward step for the field of small molecule-DNA recognition and enables the comprehensive interrogation of DNA binding small molecules to be better understood. The elucidation of 5'-WWACGT-3' as the binding site for **2** confirmed the previously established pairing code for hairpin polyamides, and the determination of 5'-AARAARWWG-3' for **4** helps explain the specificity it exhibited in cell culture. The correlation between a Cy3-labeled polyamide and an unlabeled polyamide of biological interest means that these motifs well approximate the binding profiles for **1** and **3**, respectively. DNase I footprinting-calibrated CSI microarrays have been shown to be an effective technique for determining the binding affinities of DNA-binding ligands for a vastly expanded repertoire of DNA sequences, and we envision them to be a critical tool for reliably determining sequence specificity for other ligands in the future.

## Experimental Section

Materials, methods, synthesis, plasmid preparation, and DNase I footprinting procedures are found in the Supporting Information.

**Microarray Procedures.** Microarrays were synthesized by using a Maskless Array Synthesizer (NimbleGen Systems, Madison, WI). Homopolymer (T<sub>10</sub>) linkers were covalently attached to monohydroxysilane glass slides. Oligonucleotides were then synthesized on the homopolymers to create a high-density oligonucleotide microarray. The array surface was derivatized such that the density of oligonucleotides was sufficiently low within the same feature so that no one oligonucleotide would hybridize with its neighbors. Four copies of each hairpin containing a unique 10 bp site (5'-GCGC-N<sup>1</sup>N<sup>2</sup>N<sup>3</sup>N<sup>4</sup>N<sup>5</sup>N<sup>6</sup>N<sup>7</sup>N<sup>8</sup>N<sup>9</sup>N<sup>10</sup>-GCGC-GGA-GCGC-N<sup>10</sup>N<sup>9</sup>N<sup>8</sup>N<sup>7</sup>N<sup>6</sup>N<sup>5</sup>N<sup>4</sup>N<sup>3</sup>N<sup>2</sup>N<sup>1</sup>-GCGC-3') required a total of 2 099 200 features, divided among six microarrays.

**Binding Assay.** Microarray slides were immersed in 1x PBS and placed in a 90 °C water bath for 30 min to induce hairpin formation of the oligonucleotides. Slides were then transferred to a tube of nonstringent wash buffer (saline/sodium phosphate/EDTA buffer, pH 7.5/0.01% Tween 20) and scanned to check for low background (<200 intensity). Microarrays were scanned by using an Axon 4000B,

and the image files were extracted with GENEPIX PRO Version 3.0 (Axon Instruments, Foster City, CA).

**Polyamide Binding.** Microarrays prepared as above were placed in the microarray hybridization chamber and washed twice with nonstringent wash buffer. Polyamide was diluted to 10 nM (for **2**) or 175 nM (for **4**) in Hyb buffer (100 mM Mes/1 M NaCl/20 mM EDTA, pH 7.5/0.01% Tween 20). Polyamide was then added to the hybridization chamber and incubated at room temperature for 1 h. Finally, the microarrays were washed twice with nonstringent wash buffer and scanned.

**Data Processing.** For each replicate, global mean normalization was used to ensure the mean intensity of each microarray was the same. Local mean normalization<sup>22</sup> was then used to ensure that the intensity was evenly distributed throughout each sector of the microarray surface. Outliers between replicate features were detected by using the  $Q$  test at 90% confidence and filtered out. The replicates were then quantile-normalized<sup>23</sup> to account for any possible nonlinearity between arrays.

Duplicate features were then averaged together. The median of the averaged features was subtracted to account for background.

**Acknowledgment.** This work was supported by the National Institutes of Health (GM27681 to P.B.D. and A.Z.A.). We thank the Beckman Institute Sequence Analysis Facility for DNA sequencing.

**Supporting Information Available:** Experimental materials and methods, polyamide synthesis, plasmid preparation, and DNase I footprint titration details. Histograms of the frequencies of microarray intensities and fractional standard deviations. Additional  $K_a$  vs CSI microarray intensity plots and  $K_a$ -weighted sequence logos of polyamides **2** and **4**. This material is available free of charge via the Internet at <http://pubs.acs.org/>.

JA0744899

(22) Colantuoni, C.; Henry, G.; Zeger, S.; Pevsner, J. *Bioinformatics* **2002**, *18*, 1540–1541.

(23) Bolstad, B. M.; Irizarry, R. A.; Astrand, M.; Speed, T. P. *Bioinformatics* **2003**, *19*, 185–193.