


Communication

Nucleophilic Carbene and HOAt Relay Catalysis in an Amide Bond Coupling: An Orthogonal Peptide Bond Forming Reaction

Harit U. Vora, and Tomislav Rovis

J. Am. Chem. Soc., **2007**, 129 (45), 13796-13797 • DOI: 10.1021/ja0764052 • Publication Date (Web): 11 October 2007

Downloaded from <http://pubs.acs.org> on February 14, 2009



More About This Article

Additional resources and features associated with this article are available within the HTML version:

- Supporting Information
- Links to the 29 articles that cite this article, as of the time of this article download
- Access to high resolution figures
- Links to articles and content related to this article
- Copyright permission to reproduce figures and/or text from this article

[View the Full Text HTML](#)

Nucleophilic Carbene and HOAt Relay Catalysis in an Amide Bond Coupling: An Orthogonal Peptide Bond Forming Reaction


Harit U. Vora and Tomislav Rovis*

Department of Chemistry, Colorado State University, Fort Collins, Colorado 80523

Received August 25, 2007; E-mail: rovis@lamar.colostate.edu

The ubiquity of amides throughout organic, biological, and materials chemistry mandates the development of more efficient methods for their synthesis.¹ Conventional amide bond formation utilizes acids and amines as coupling partners and relies on stoichiometric activating agents for the acid functionality.^{2,3} A recent survey of process scale reactions cites a "...pressing need for the development of catalytic environmentally friendly acylation processes".^{4,5} We⁶ and others⁷ have recently illustrated that nucleophilic carbenes⁸ catalyze an internal redox reaction whereby α -reducible aldehydes provide α -reduced ester derivatives under catalytic conditions.⁹ Surprisingly, among the many nucleophiles reported to participate in this process, there are only two amines: we^{6a} have shown that aniline participates, and Scheidt^{7b} has shown that a vinylogous imide could be used.¹⁰ Clearly, the salient features of this redox manifold, a waste free catalyzed acylation, provide a strong impetus to identify a general solution to the problem of NHC catalyzed amidation. Herein, we report one such solution relying on relay catalysis by a nucleophilic carbene and a common peptide catalyst such as 1-hydroxy-7-azabenzotriazole¹¹ (HOAt).

Outside of aniline, our efforts at using amines as nucleophiles in the α -redox reaction were met with uniform failure. Since we had established that phenols are competent partners, we hypothesized that the use of a cocatalyst such as HOAt could provide a relay shuttle.^{12,13} HOAt should participate in the redox chemistry to generate activated ester which would undergo the in situ amidation, thereby regenerating the catalyst. The viability of a concerted catalytic system using N-heterocyclic carbenes and HOAt to generate amides was investigated utilizing 2,2-dichloro-3-phenylpropanal as the redox substrate and benzyl amine as the nucleophile. The desired chemical transformation took place to afford the benzyl amide **2a** in 93% yield (eq 1). In the absence of HOAt, only minor amidation product is observed.¹⁴ A cocatalyst screen revealed that 1-hydroxybenzotriazole (HOBt), 4-(dimethylamino)pyridine (DMAP), imidazole, and pentafluorophenol (PFPOH) are effective at promoting the reaction, affording the desired amide products.



Experiments that probe the scope of useful amine partners are summarized in Table 1. A variety of primary and secondary amines partake in the reaction (entries 1–5, Table 1) to afford the desired amide in good to excellent yields. Of particular interest is the generation of the Weinreb amide **2f** in 72% yield (entry 5, Table

Table 1. Amine Scope

Entry ^a	Yield (%)	Entry ^a	Yield (%)
1	89	6	87
2	85	7	82
3	73	8	83
4	89	9	85 ^{b,c}
5	72		

^a Catalyst **A** (20 mol %), HOAt (20 mol %), Et₃N (1.2 equiv), THF (0.5 M), *t*-BuOH (1.0 equiv), 25 °C, 6 h, unless otherwise noted. ^b HOBt (20 mol %) and Et₃N (2.1 equiv) were used. ^c Diastereomeric ratio 2:1.

1). Electron-rich and -poor aryl amines **2f–h** (entries 6–8, Table 1) also undergo the transformation readily to give the desired anilides in 82–87% yield. Amino esters are also competent partners (entry 9, Table 1).

A variety of α -haloaldehydes are suitable partners in the α -redox amidation. The reaction is tolerant of branching at the α and β position: α,α -dichloroisovaleraldehyde provides **3** in 72% yield, and α -bromocyclohexancarboxaldehyde provides **4** in 80% yield (Figure 1).

One of the strengths of the redox amidation reaction manifold is that the appropriate choice of α -reducible aldehyde provides an opportunity for a waste free amidation. Treatment of α,β -epoxy and aziridino aldehydes under the redox amidation conditions affords β -hydroxy and β -amino amides (entries 1–3, Table 2) in good yields and excellent diastereoselectivities. α,β -Unsaturated aldehydes provide the alkanamides in good yield (entries 4 and 5,





Figure 1. α -Haloaldehyde substrate scope.

Table 2. Atom-Economical Amidation


Entry	Substrate	Product	Yield (%)	dr
1			86 ^a	>19:1
2			75 ^a	15:1
3			72 ^a	>19:1
4			80 ^b	-
5			82 ^b	-

^a Catalyst **A** (10 mol %), imidazole (10 mol %), DIPEA (30 mol %), *t*-BuOH (0.1 M), 40 °C, 24 h. ^b Catalyst **A** (10 mol %), HOAt (10 mol %), DIPEA (10 mol %), THF (1.0 M), 45 °C.



Scheme 1. Proposed Catalytic Cycle

Table 2). Importantly, in each case, the only stoichiometric waste generated is derived from solvent; even the base is used in catalytic amounts.

The catalytic cycle is postulated to initiate upon formation of carbene **I**, which undergoes nucleophilic addition to the aldehyde (Scheme 1). Generation of the acylazolium intermediate **II** sets the stage for an acyl transfer event with cocatalyst **III** to furnish the activated carboxylate **IV**. Nucleophilic attack by the amine affords the amide and regenerates the cocatalyst.

Experimental support for the proposed mechanism is provided by the use of chiral carbenes in this process. The use of catalyst **B** leads to an asymmetric α -chloroamide synthesis in modest ee (eq 2), validating the role of the carbene in controlling the protonation event. In contrast, the use of **B** provides no selectivity in the kinetic resolution of α -methylbenzyl amine (eq 3). In addition, the use of stoichiometric HOAt in the absence of amine provides the HOAt ester **IV** in 64% yield. Addition of BnNH_2 generates the amide quantitatively.

In summary, we have developed a waste free amide bond forming reaction using α -reducible aldehydes and amines catalyzed by carbenes in conjunction with common peptide additives as cocatalysts.



Acknowledgment. We thank NIGMS (GM72586), Eli Lilly, Johnson and Johnson, and Boehringer Ingelheim for support. T.R. is a fellow of the Sloan Foundation and thanks the Monfort Family Foundation for a Monfort Professorship.

Note Added after ASAP Publication. Reference 11c was added on October 12, 2007.

Supporting Information Available: Experimental procedures and full characterization of all new compounds (PDF). This material is available free of charge via the Internet at <http://pubs.acs.org>.

References

- Humphrey, J. M.; Chamberlin, A. R. *Chem. Rev.* **1997**, *97*, 2243.
- Larock, R. C. *Comprehensive Organic Transformations*; VCH: New York, 1999.
- For some recent exceptions, see: (a) Shangguan, N.; Katukojvala, S.; Greenberg, R.; Williams, L. J. *J. Am. Chem. Soc.* **2003**, *125*, 7754. (b) Bode, J. W.; Fox, R. M.; Baucom, K. D. *Angew. Chem., Int. Ed.* **2006**, *45*, 1248. (c) Carrillo, N.; Davalos, E. A.; Russak, J. A.; Bode, J. W. *J. Am. Chem. Soc.* **2006**, *128*, 1452.
- (a) Carey, J. S.; Laffan, D.; Thomson, C.; Williams, M. T. *Org. Biomol. Chem.* **2006**, *4*, 2337. (b) Also see: Dugger, R. W.; Ragan, J. A.; Brown Ripin, D. H. *Org. Process Res. Dev.* **2005**, *9*, 253.
- For a sampling of recent contributions in this area, see: (a) Ishihara, K.; Ohara, S.; Yamamoto, H. *J. Org. Chem.* **1996**, *61*, 4196. (b) Eldred, S. E.; Stone, D. A.; Gellman, S. H.; Stahl, S. S. *J. Am. Chem. Soc.* **2003**, *125*, 3422–3423. (c) Yoo, W. J.; Li, C. J. *J. Am. Chem. Soc.* **2006**, *128*, 13064. (d) Ekoue-Kovi, K.; Wolf, C. *Org. Lett.* **2007**, *9*, 3429.
- (a) Reynolds, N. T.; Read de Alaniz, J.; Rovis, T. *J. Am. Chem. Soc.* **2004**, *126*, 9518. (b) Reynolds, N. T.; Rovis, T. *J. Am. Chem. Soc.* **2005**, *127*, 16406.
- (a) Chow, K. Y.-K.; Bode, J. W. *J. Am. Chem. Soc.* **2004**, *126*, 8126. (b) Chan, A.; Scheidt, K. A. *Org. Lett.* **2005**, *7*, 905. (c) Sohn, S. S.; Bode, J. W. *Org. Lett.* **2005**, *7*, 3873. (d) Sohn, S. S.; Bode, J. W. *Angew. Chem., Int. Ed.* **2006**, *45*, 6021. (e) Zeitler, K. *Org. Lett.* **2006**, *8*, 637. (f) Grasa, G. A.; Güveli, T.; Singh, R.; Nolan, S. P. *J. Org. Chem.* **2003**, *68*, 2812.
- Reviews: (a) Enders, D.; Balensiefer, T. *Acc. Chem. Res.* **2004**, *37*, 534. (b) Johnson, J. S. *Angew. Chem., Int. Ed.* **2004**, *43*, 1326. (c) Zeitler, K. *Angew. Chem., Int. Ed.* **2005**, *44*, 7506. (d) Marion, N.; Díez-González, S.; Nolan, S. P. *Angew. Chem., Int. Ed.* **2007**, *46*, 2988.
- This reactivity was first described by Wallach, who found that treatment of chloral with aqueous cyanide provides dichloroacetic acid: Wallach, O. *Ann. Chem.* **1873**, *6*, 114.
- Movassaghi has shown that amide products can be obtained from treatment of ester with amino alcohols using N-heterocyclic carbenes. This reactivity has been shown to proceed via an intramolecular O \rightarrow N acyl transfer to generate the amide. Movassaghi, M.; Schmidt, M. A. *Org. Lett.* **2005**, *7*, 2453.
- (a) König, W.; Geiger, R. *Ber. Dtsch. Chem. Ges.* **1970**, *103*, 788. (b) König, W.; Geiger, R. *Ber. Dtsch. Chem. Ges.* **1970**, *103*, 2024. (c) Carpino, L. A. *J. Am. Chem. Soc.* **1993**, *115*, 4397.
- Catalytic HOAt and HOBt in ester amidation: Han, C.; Lee, J. P.; Lobkovsky, E.; Porco, J. A. *J. Am. Chem. Soc.* **2005**, *127*, 10039.
- Cascade catalysis: (a) Hourri, A. F.; Xu, Z.; Cogan, D. A.; Hoveyda, A. H. *J. Am. Chem. Soc.* **1995**, *117*, 2943. (b) Bruggink, A.; Schoevaart, R.; Kieboom, T. *Org. Process Res. Dev.* **2003**, *7*, 622. (c) Huang, Y.; Walji, A. M.; Larsen, C. H.; MacMillan, D. W. C. *J. Am. Chem. Soc.* **2005**, *127*, 15051. (d) Concurrent tandem catalysis: Wasilke, J.-C.; Obrey, S. J.; Baker, R. T.; Bazan, G. C. *Chem. Rev.* **2005**, *105*, 1001.
- In the absence of cocatalyst, the reaction generates significant amounts of imine and α -reduced carboxylic acid.

JA0764052