


Alkane Hydroxylation by a Nonheme Iron Catalyst that Challenges the Heme Paradigm for Oxygenase Action

Anna Company, Laura Gmez, Mireia Gell, Xavi Ribas, Josep M. Luis, Lawrence Que,, and Miquel Costas

J. Am. Chem. Soc., **2007**, 129 (51), 15766-15767 • DOI: 10.1021/ja077761n

Downloaded from <http://pubs.acs.org> on February 9, 2009



More About This Article

Additional resources and features associated with this article are available within the HTML version:

- Supporting Information
- Links to the 8 articles that cite this article, as of the time of this article download
- Access to high resolution figures
- Links to articles and content related to this article
- Copyright permission to reproduce figures and/or text from this article

[View the Full Text HTML](#)

Alkane Hydroxylation by a Nonheme Iron Catalyst that Challenges the Heme Paradigm for Oxygenase Action

Anna Company,^{†,‡} Laura Gómez,[†] Mireia Güell,[†] Xavi Ribas,[†] Josep M. Luis,[†] Lawrence Que, Jr.,^{*,‡} and Miquel Costas^{*,†}

Departament de Química and Institut de Química Computacional, Campus de Montilivi, Universitat de Girona, E-17071 Girona, Spain, and Department of Chemistry and Center for Biocatalysis, University of Minnesota, Minneapolis, Minnesota 55455

Received October 9, 2007; E-mail: miquel.costas@udg.es

Selective oxidation of hydrocarbons under mild conditions constitutes a major challenge of modern chemistry.¹ Nonheme iron enzymes, such as methane monooxygenase² and Rieske dioxygenases,³ catalyze such reactions and have inspired the development of synthetic models as alkane oxidation catalysts.⁴ Exceptional nonheme iron catalysts are those containing tetradentate N₄ ligands belonging to the tripodal TPA and linear BPMEN family that can perform efficient stereospecific alkane hydroxylation using H₂O₂ as oxidant.⁴ These bioinspired catalysts have provided key insights into the mechanisms by which alkanes are oxidized by enzymes. The common mechanism invoked in these studies follows the heme paradigm⁵ in which a high-valent iron-oxo species first abstracts a H atom from a substrate C–H bond and the incipient alkyl radical then rebounds with the nascent Fe–OH moiety to form the product C–O bond (Scheme 1).^{2,4} Labeling studies indicate that the O atom introduced into the substrate C–H bond derives mainly from the H₂O₂ oxidant. However, a minor fraction incorporates an O atom from water into the product, which is explained by oxo-hydroxo tautomerism of the high-valent HO–Fe=O oxidant.^{5,6} Herein we describe a new remarkably efficient nonheme iron catalyst in the stereospecific oxidation of alkanes using H₂O₂ as oxidant that yields alcohol products with an unexpectedly large fraction of their oxygen atoms derived from water. This unprecedented result challenges the established heme paradigm and sheds new light into how nonheme iron centers may differ from heme centers in their mode of oxidative action.

Iron complex [Fe^{II}(CF₃SO₃)₂(Me₂PyTACN)] (**1**) (see Figure 1) was prepared and structurally characterized (for details, see Supporting Information). Its crystal structure presents a distorted octahedral iron(II) center having a tetradentate Me₂PyTACN ligand and two triflate anions coordinated *cis* to each other.

Complex **1** is a very efficient catalyst for the hydroxylation of cyclohexane by H₂O₂. When 10 equiv of H₂O₂ were delivered by syringe pump together with 1000 equiv of water, 6.0 turnover numbers (TN) of cyclohexanol (*A*) and 0.5 TN of cyclohexanone (*K*) (*A*/*K* = 12) were obtained, corresponding to a 65% efficiency in the conversion of H₂O₂ into products. Both efficiency and *A*/*K* selectivity exhibited by **1** were comparable to values reported for [Fe^{II}(BPMEN)(CH₃CN)₂]²⁺ (63% yield and *A*/*K* = 8), considered the prototypical example of an efficient stereospecific alkane hydroxylation catalyst.^{4b} Additional evidence implicated a metal-based oxidant rather than HO•. The intermolecular kinetic isotope effect evaluated in the hydroxylation of a 1:3 molar mixture of cyclohexane and its deuterio-analogue was 4.3. In addition, **1** oxidized adamantane with a large 3°/2° normalized selectivity (30). Last, the oxidation of *cis*-1,2-dimethylcyclohexane (*cis*-DMCH)

Scheme 1






Figure 1. Left: schematic representation of the ligand. Right: thermal ellipsoid plot (50% probability) of **1**; triflate ligands (except for the O atoms bound to Fe) are omitted for clarity.

afforded the tertiary alcohol product with 93% retention of configuration. Overall, these characteristics are consistent with those of some previously described nonheme iron complexes.⁴


Interestingly, ¹⁸O-labeling experiments in the hydroxylation of different alkanes catalyzed by **1** revealed intriguing differences between this catalyst and previously reported nonheme N₄Fe systems. The oxidation of cyclohexane by **1** with 10 equiv of H₂¹⁸O₂ in the presence of 1000 equiv of H₂¹⁶O afforded 47% ¹⁸O-labeled cyclohexanol. Complementary experiments with 10 equiv of H₂O₂ and 1000 equiv of H₂¹⁸O yielded 42(3)% ¹⁸O-labeled cyclohexanol. The same levels of label incorporation from H₂¹⁸O were observed in the hydroxylation of cyclooctane (Table 1). This level of oxygen incorporation from water was the highest for any synthetic nonheme catalyst thus far. Furthermore, when oxidations of cyclohexane were carried out with different H₂¹⁸O concentrations, the fraction of ¹⁸O-labeled cyclohexanol (% R¹⁸OH) increased linearly with the amount of H₂¹⁸O at lower H₂¹⁸O concentrations, but plateaued at 42(3)% at higher concentrations (Figure S2), suggesting the involvement of a water binding pre-equilibrium, as preceded for Fe(TPA).^{4b} These results support the mechanism shown in Scheme 1 proposed for Fe(TPA) and Fe(BPMEN) catalysts in which a HO–Fe=O oxidant is formed via water-assisted heterolysis of the O–O bond of a H₂O–Fe^{III}–OOH intermediate.^{4b,7} If complete oxo-hydroxo tautomerism^{5,6} of the HO–Fe^V=O moiety occurred, a maximum value of 50% would be expected for the amount of labeled water incorporated into the alcohol product. The H₂¹⁸O-labeling results described above clearly approach this maximum, so that the C–O bond that is formed can be made with either the oxygen from H₂O₂ or the oxygen from H₂O with almost equal probability.

[†] Universitat de Girona.

[‡] University of Minnesota.

Table 1. Percentage of ^{18}O Incorporation into Alcohol Products by Fe(L) Catalysts in the Presence of 1000 equiv of H_2^{18}O

substrate	Me ₂ PyTACN	TPA ^{4b}
cyclohexane	42	29
cyclohexane- <i>d</i> ₁₂	40	35
cyclooctane	44	23
<i>cis</i> -DMCH (3° C–H)	79	6
adamantane (3° C–H)	74	6
2,3-dimethylbutane	76	

Scheme 2

Surprisingly, even higher levels of H_2^{18}O incorporation (76 ± 3%) were obtained for alcohol products in the oxidation of alkanes with tertiary C–H bonds, such as adamantane, *cis*-DMCH, and 2,3-dimethylbutane. Furthermore, this unexpectedly large value was found to be independent of substrate concentration (25–1000 mM). This observation strongly implicates the $\text{HO–Fe}^{\text{V}}=\text{O}$ species as the only oxidant capable of alkane oxidation in the case of **1**.

The labeling results for **1** differ significantly from those reported for Fe(TPA) (Table 1). For the latter catalyst, label incorporation from H_2^{18}O into substrates with 2° C–H bonds decreased with the C–H bond strength and labeling of 2°-ol products was much higher than for 3°-ol products. These results suggested that C–H bond cleavage and oxo-hydroxo tautomerism were competitive processes, as also found for iron porphyrin complexes.^{4b,5,6} However, this was not the case for **1** since higher water incorporation was observed in the oxidation of the weaker 3° C–H bonds, and this level of incorporation was independent of *cis*-DMCH concentration. To rationalize the high level of label incorporation, we considered the possibility that a carbocation intermediate was formed and subsequently trapped by water, but discarded it because of the high retention of configuration observed for *cis*-DMCH hydroxylation. We also considered the possibility that the $\text{HO–Fe}^{\text{V}}=\text{O}$ oxidant became doubly labeled by rapid, multiple intermolecular exchanges with H_2^{18}O , but rejected it on the basis of a H_2^{18}O -labeling experiment where cyclohexane and *cis*-DMCH were oxidized in competition with each other. *cis*-DMCH reacted much faster than cyclohexane, and so should have a shorter lived oxidant. Nevertheless, the 3°-ol thus formed contained 74% label from water, while cyclohexanol contained only 38%. Thus, the difference in label incorporation cannot be determined by how much ^{18}O is present in the common Fe^{V} oxidant, and a modified mechanistic scenario is required for **1**.

In Scheme 2, we propose a scenario in which the structure of the substrate can play a role in determining the course of C–O bond formation. For **1**, the two $\text{Fe}^{\text{V}}(\text{O})(\text{OH})$ isomers (**A** and **B**) in equilibrium via oxo-hydroxo tautomerism are distinct since the ligands *trans* to the oxo and hydroxo groups are not chemically equivalent and the orientations of the pyridine ligand relative to the $\text{Fe}=\text{O}$ bond are different. No matter which isomer abstracts the H atom from substrate, a common $\text{Fe}^{\text{IV}}(\text{OH})_2$ species is formed. The *cis* configuration of the two hydroxo groups differs from the *trans* configuration required in heme complexes and is a unique

feature of these nonheme iron catalysts. Thus, the alkyl radical can in principle rebound with either OH group, the outcome of which determines the % H_2^{18}O incorporation. Given that these values range from 40% for cyclohexane-*d*₁₂ to 79% for *cis*-DMCH, the energy difference between the two possible rebound trajectories is <2 kcal mol⁻¹, which is consistent with DFT calculations.⁸ Thus, the labeling results indicate that 2°-alkyl radicals do not discriminate between the two OH groups, but 3°-alkyl radicals favor rebound with the OH group derived from water. Perusal of the structure of **1** (Figure 1) suggests steric effects may provide a possible rationale for this preference since the hemisphere surrounding O1 is less sterically congested.

In conclusion, **1** is an efficient alkane hydroxylation catalyst that can incorporate surprisingly large amounts of water into products via an unusual rebound-like mechanism. A similar mechanism may be operating in the hydroxylation of indane to 1-indanol by toluene or naphthalene dioxygenase to account for the high level of H_2^{18}O incorporation (68%) found.⁹ In addition, the α -ketoglutarate-dependent halogenases represent an established biological example in which C–H bond cleavage initiated by an $\text{Fe}=\text{O}$ moiety is consummated by rebound not with the incipient OH group but with the adjacent halide ligand.¹⁰ Thus, **1** may serve as a model for this unusual enzymatic chemistry, which suggests that a broader mechanistic landscape than in heme systems applies to nonheme sites.

Acknowledgment. Financial support from MCYT of Spain (CTQ2006–05367/BQU to M.C.) and from US-DOE (DE-FG02-03ER15455 to L.Q.) is gratefully acknowledged. A.C., L.G., and M.G. thank MEC for Ph.D. grants.

Supporting Information Available: Experimental procedures for the preparation of **1** and for the oxidation reactions and the cif file for **1**. This material is available free of charge via the Internet at <http://pubs.acs.org>.

References

- (1) (a) Tanase, S.; Bouwman, E. *Adv. Inorg. Chem.* **2006**, *58*, 29–75. (b) *Biomimetic Oxidations Catalyzed by Transition Metal Complexes*; Imperial College Press: London, 2000.
- (2) Merkx, M.; Kopp, D. A.; Sazinsky, M. H.; Blazyk, J. L.; Müller, J.; Lippard, S. J. *Angew. Chem., Int. Ed.* **2001**, *40*, 2782–2807.
- (3) (a) Costas, M.; Mehn, M. P.; Jensen, M. P.; Que, L., Jr. *Chem. Rev.* **2004**, *104*, 939. (b) Abu-Omar, M. M.; Loaiza, A.; Hontzeas, N. *Chem. Rev.* **2005**, *105*, 2227.
- (4) (a) Costas, M.; Chen, K.; Que, L., Jr. *Coord. Chem. Rev.* **2000**, *200–202*, 517. (b) Chen, K.; Que, L., Jr. *J. Am. Chem. Soc.* **2001**, *123*, 6327. (c) Britovsek, G. J. P.; England, J.; White, A. J. P. *Inorg. Chem.* **2005**, *44*, 8125. (d) Mekmouche, Y.; Ménage, S.; Toia-Duboc, C.; Fontecave, M.; Galey, J.-B.; Lebrun, C.; Pécaut, J. *Angew. Chem., Int. Ed.* **2001**, *40*, 949.
- (5) (a) Meunier, B.; de Visser, S. P.; Shaik, S. *Chem. Rev.* **2004**, *104*, 3947–3980. (b) Groves, J. T.; Han, Y. Z. In *Cytochrome P450: Structure, Mechanism and Biochemistry*; Ortiz de Montellano, P. R., Ed.; Plenum: New York, 1995. (c) Bernadou, J.; Meunier, B. *Chem. Commun.* **1998**, 2167.
- (6) (a) Bernadou, J.; Fabiano, A.-S.; Robert, A.; Meunier, B. *J. Am. Chem. Soc.* **1994**, *116*, 9375. (b) Groves, J. T.; Lee, J.; Marla, S. S. *J. Am. Chem. Soc.* **1997**, *119*, 6269. (c) Lee, K. A.; Nam, W. *J. Am. Chem. Soc.* **1997**, *119*, 1916.
- (7) (a) Bassan, A.; Blomberg, M. R. A.; Siegbahn, P. E. M.; Que, L., Jr. *J. Am. Chem. Soc.* **2002**, *124*, 11056. (b) Quinonero, D.; Morokuma, K.; Musaev, D. G.; Mas-Balleste, R.; Que, L., Jr. *J. Am. Chem. Soc.* **2005**, *127*, 6548.
- (8) DFT calculations were done using cyclohexane as a substrate. Geometries were optimized at the B3LYP level in junction of the LANL2DZ basis set with associated ECP for Fe. The energies were further refined by single-point calculations using the SDD basis set with associated ECP for Fe and 6-311G(d,p) basis sets on the other atoms as implemented in the Gaussian 03 program.
- (9) Wackett, L. P.; Kwart, L. D.; Gibson, D. T. *Biochemistry* **1988**, *27*, 1360–1367.
- (10) Galonic, D. P.; Barr, E. W.; Walsh, C. T.; Bollinger, J. M., Jr.; Krebs, C. *Nat. Chem. Biol.* **2007**, *3*, 113.

JA077761N