

Addition/Correction

Simple and Effective 3D Recognition of Domoic Acid Using a Molecularly Imprinted Polymer [J. Am. Chem. Soc. 2007, 129, 13626–13632].

Koji Nemoto, Takuya Kubo, Makoto Nomachi, Tomoharu Sano, Takatoshi Matsumoto, Ken Hosoya, Tetsutaro Hattori, and Kunimitsu Kaya

J. Am. Chem. Soc., **2008**, 130 (2), 774-774 • DOI: 10.1021/ja7106788

Downloaded from <http://pubs.acs.org> on February 8, 2009

More About This Article

Additional resources and features associated with this article are available within the HTML version:

- Supporting Information
- Access to high resolution figures
- Links to articles and content related to this article
- Copyright permission to reproduce figures and/or text from this article

[View the Full Text HTML](#)



Simple and Effective 3D Recognition of Domoic Acid Using a Molecularly Imprinted Polymer [*J. Am. Chem. Soc.* **2007**, *129*, 13626–13632]. Koji Nemoto, Takuya Kubo,* Makoto Nomachi, Tomoharu Sano, Takatoshi Matsumoto, Ken Hosoya, Tetsutaro Hattori, and Kunimitsu Kaya

Page 13630. In Table 2, the second column (ΔG of DA) should be removed. The correct table is given below.

Table 2. Gibbs Free Energies of DA on Chromatographic Evaluation

polymers	K' of DA	$\Delta G - \Delta G(\text{on BASE})$ (kJ/mol)	$\Delta G - \Delta G(\text{on BLANK})$ (kJ/mol)
BASE	0.10	—	—
BLANK	1.10	6.04	—
OPA-MIP	9.62	11.5	5.46
2,3-NDA-MIP	13.7	12.4	6.35
2,6-NDA-MIP	5.94	10.3	4.25
EDA-MIP	12.6	12.2	6.14
FDA-MIP	11.6	12.0	5.93
KDA-MIP	14.5	12.6	6.51
1,3,5-PeTA-MIP	53.5	15.8	9.79

JA7106788

10.1021/ja7106788

Published on Web 12/15/2007