

Addition/Correction

Sulfonic Acid-Functionalized Gold Nanoparticles: A Colloid-Bound Catalyst for Soft Lithographic Application on Self-Assembled Monolayers

Xue-Mei Li, Vasile Paraschiv, Jurriaan Huskens, and David N. Reinhoudt

J. Am. Chem. Soc., **2008**, 130 (16), 5606-5606 • DOI: 10.1021/ja801758r • Publication Date (Web): 28 March 2008

Downloaded from <http://pubs.acs.org> on February 8, 2009

More About This Article

Additional resources and features associated with this article are available within the HTML version:

- Supporting Information
- Links to the 1 articles that cite this article, as of the time of this article download
- Access to high resolution figures
- Links to articles and content related to this article
- Copyright permission to reproduce figures and/or text from this article

[View the Full Text HTML](#)

Sulfonic Acid-Functionalized Gold Nanoparticles: A Colloid-Bound Catalyst for Soft Lithographic Application on Self-Assembled Monolayers [*J. Am. Chem. Soc.* **2003**, *125*, 4279–4284]. Xue-Mei Li, Vasile Paraschiv, Jurriaan Huskens,* and David N. Reinhoudt*

Page 4280, Experimental Section. In the first sentence of the paragraph describing the synthetic procedure for bis(ω -trimethylsilyloxyundecyl)disulfide ((TMS-OC₁₁H₂₂S)₂), the amount of iodine (0.5 g) is incorrect. This should be 0.8 g. The amount 3.10 mmol is correct.

JA801758R

10.1021/ja801758r

Published on Web 03/28/2008