


Communication

Mechanistic Insights into the Gold-Catalyzed Cycloisomerization of Bromoallenyl Ketones: Ligand-Controlled Regioselectivity

Yuanzhi Xia, Alexander S. Dudnik, Vladimir Gevorgyan, and Yahong Li

J. Am. Chem. Soc., **2008**, 130 (22), 6940-6941 • DOI: 10.1021/ja802144t • Publication Date (Web): 08 May 2008

Downloaded from <http://pubs.acs.org> on February 8, 2009



More About This Article

Additional resources and features associated with this article are available within the HTML version:

- Supporting Information
- Links to the 2 articles that cite this article, as of the time of this article download
- Access to high resolution figures
- Links to articles and content related to this article
- Copyright permission to reproduce figures and/or text from this article

[View the Full Text HTML](#)

Mechanistic Insights into the Gold-Catalyzed Cycloisomerization of Bromoallenyl Ketones: Ligand-Controlled Regioselectivity

Yuanzhi Xia,^{†,‡} Alexander S. Dudnik,[§] Vladimir Gevorgyan,[§] and Yahong Li^{*†,‡}


Key Laboratory of Organic Synthesis of Jiangsu Province, Department of Chemistry and Chemical Engineering, Suzhou University, Suzhou 215006, China, Qinghai Institute of Salt Lakes, Chinese Academy of Sciences, Xining 810008, China, and Department of Chemistry, University of Illinois at Chicago, Chicago, Illinois 60607-7061

Received March 23, 2008; E-mail: liyahong@suda.edu.cn

In recent years, the homogeneous catalysis by gold complexes received increasing attention. Particularly, cyclizations of alkynes and allenes mediated by gold complexes have been widely explored.¹ Along this line, the Gevorgyan group has recently reported a gold-catalyzed regiodivergent cycloisomerization of bromoallenyl ketones **a** into isomeric bromofurans **b** and **c** (Scheme 1).² According to the originally proposed mechanism, products **b** and **c** result from different activation modes of the substrate by gold complexes.² It was thought that more oxophilic Au(III) catalyst³ coordinates to the carbonyl oxygen (**d**), which, via halirenium intermediate **e**, produces the 3-bromofuran **b**, a product of 1,2-Br migration. Alternatively, the more π -philic Au(I) catalyst coordinates to the distal double bond of the allene (**f**), leading to the gold carbenoid intermediate **g**. Subsequent 1,2-hydride shift in the latter forms 2-bromofuran **c** (Scheme 1).² Great interest has been aroused by this important discovery, since it not only opens an unprecedented access to different types of halofurans but also represents the very first example of reaction, in which employment of gold catalyst in different oxidation states leads to different products.^{1b,c,4–6} A more detailed mechanistic investigation of origins of the observed regioselectivity of this cycloisomerization, properties of the key intermediates, as well as profiles of the overall catalytic cycles, may give better understanding on the important transition metal-catalyzed chemistry of alkynes and allenes. This in turn, may provide a guide for future design of new catalytic transformations. Herein, we wish to report our theoretical and experimental results on the mechanism of Au(III)- and Au(I)-catalyzed cycloisomerization of bromoallenyl ketones, with the emphasis on the origins of the regiochemistry of 1,2-migrations. Our study indicates that the 1,2-H migration could be assisted by the ligand L of the Au(PR'₃)₃L complex, thus, the regioselectivity of 1,2-migrations can be controlled by ligands at Au-complexes.

Thus, DFT calculations to investigate the above-mentioned cycloisomerization have been performed.^{7,8} The computed energy surfaces for the AuCl₃ and Au(PH₃)Cl-catalyzed cycloisomerizations of bromoallenyl ketone **1** are provided in Figure 1. Results of the Au(III)-catalyzed reaction (**I**, Figure 1) indicate that the gold catalyst, upon coordination to the distal double bond of the allene **1**,⁹ instantly gives the cyclic intermediate **2** via a barrierless and highly exergonic process with relative free energies of 32.6 and 33.6 kcal/mol, in the gas phase and in toluene solution, respectively. Next, intermediate **2** undergoes more favorable 1,2-Br migration via the transition state **TS1** to generate the product complex **3** rather than 1,2-H shift through **TS2** to give **4**. Thus, the computed free energy barriers difference of 14.3 kcal/mol for the latter processes is in good agreement with the experimentally observed nearly exclusive 1,2-Br migration.² Although, the formation of the complex **3** from intermediate **2** is endergonic (Figure 1), the latter will be transformed into the product **b** irreversibly via a highly

Scheme 1. Regiodivergent Au(III)- and Au(I)-Catalyzed Cycloisomerizations of Bromoallenyl Ketones



exergonic (30.3 kcal/mol) ligand exchange process of **3** with another molecule of substrate, thus, completing the catalytic cycle.¹⁰

Remarkably, it was found that the AuCl₃-catalyzed cycloisomerization of **1** in wet toluene¹¹ leads to the decreased ratio of 1,2-Br vs 1,2-H migration products. This implies that the H-shift could be catalyzed by a trace amount of water present in the system, analogously to the recently reported H₂O-assisted formal 1,2-H shift in the Au⁺-catalyzed carbocyclization.^{8c} This hypothesis was unambiguously confirmed by the calculations and further by reactions performed in the presence of D₂O.¹⁰ Thus, DFT calculations revealed the H₂O-assisted H-shift from **2** to **4** is a stepwise H-abstraction/donation process with an activation free energy of 17.8 kcal/mol in toluene, which is only about 1.4 kcal/mol higher than that for the 1,2-Br migration.¹⁰

The energy surface for the reaction catalyzed by Au(I) (**II**, Figure 1) is quite different from that for Au(III). Coordination of the less electrophilic Au(PH₃)Cl to the distal double bond of **1** gives π -complex **5** with the free energy increased by 20.8 kcal/mol in toluene solution. The latter, via the **TS3** with 10.5 kcal/mol activation energy barrier, transforms into the zwitterionic intermediate **6**. From intermediate **6**, the reaction may proceed through the direct 1,2-migration steps to generate complexes **7** and **8**. Calculations showed that the activation barrier for the direct 1,2-H shift (**TS5**) is 30.1 kcal/mol, while that for the 1,2-Br migration (**TS4**) is only 14.6 kcal/mol, thus contradicting with the experimental results on the observed predominant 1,2-H shift.² However, the major 1,2-H migration pathway can reasonably be rationalized in terms of the stepwise ligand-assisted H-shift process. Thus, as shown in **II** (Figure 1), **6** rotamerizes to a relatively more stable intermediate **9**. Geometries of **6** and **9** revealed the remarkably prolonged Au–Cl distances, implying quite weak Au–Cl interactions.¹² Consequently, the weakly coordinated chloride is now capable of hydrogen abstraction at C2 via **TS6** with a barrier of 6.3 kcal/mol only to give intermediate **10**. Notably, the HCl moiety is associated with the gold-substituted furan through Cl–H–C3 hydrogen-bonding interaction.¹⁰ Gas phase calculations showed that the formation of **10** is slightly exergonic, though a solvation by toluene substantially destabilizes this intermediate. Protodeauration of **10** at C3 via **TS7** leads to the product complex **8**. The activation free energy of this step is 15.7 and 8.7 kcal/mol in the gas phase and in toluene, respectively. Dissociation of **8** is quite favorable with an exergonicity of 26.3 kcal/

[†] Suzhou University.

[‡] Qinghai Institute of Salt Lakes, Chinese Academy of Sciences.

[§] University of Illinois at Chicago.





Figure 1. Potential energy surfaces for AuCl_3 - and $\text{Au}(\text{PR}'_3)\text{Cl}$ -catalyzed cycloisomerizations ($\text{R} = \text{CH}_3$, $\text{R}' = \text{H}$).⁷

Scheme 2. Experimental Results of Au(I)-Catalyzed Reactions¹⁰



mol in toluene. Thus, the overall barrier for the chloride-assisted H-migration process from **9** to the complex **8** is only 8.7 kcal/mol.

On the basis of these results, we hypothesized that if no labile chloride ligand is present in a Au(I)-catalyzed cycloisomerization (e.g., using cationic Au(I) or AuCl complexes), the reaction may proceed via a predominant 1,2-Br migration. As predicted, our experiments indicated that employment of AuCl or cationic catalysts, such as $\text{Au}(\text{PPh}_3)\text{BF}_4$ and $\text{Au}(\text{PPh}_3)\text{SbF}_6$, produced the 1,2-Br migration product **A** as a major or sole isomer (Scheme 2).¹⁰ Unexpectedly, use of $\text{Au}(\text{PPh}_3)\text{OTf}$ led to exclusive formation of the 1,2-H shift product **B**, thus, exhibiting a dramatic counterion effect on the regioselectivity of the reaction. The DFT calculations shed light on this seeming controversy.^{8d,10} The computations revealed that in both $\text{Au}(\text{PR}_3)^+$ and AuCl-catalyzed reactions, the 1,2-Br migration is more favorable with the activation barrier about 13 kcal/mol. However, 1,2-H shift in $\text{Au}(\text{PR}_3)^+$ -catalyzed reactions can be assisted by the counterions. Thus, it was found that the OTf^- -assisted 1,2-H shift requires an activation free energy of only 9.6 kcal/mol, whereas those for BF_4^- and SbF_6^- are much higher (20.2 and 29 kcal/mol, respectively), which is in a good agreement with the experiments.

In summary, the mechanism of Au-catalyzed cycloisomerization of bromoalleny ketones has been investigated through DFT computational and experimental studies. It was found that both Au(I) and Au(III) catalysts¹³ activate the distal double bond of the allene to produce cyclic zwitterionic intermediates, which undergo a kinetically favored 1,2-Br migration. However, in the cases of $\text{Au}(\text{PR}_3)\text{L}$ ($\text{L} = \text{Cl}, \text{OTf}$) catalysts, the counterion-assisted H-shift is the major process, indicating that the regioselectivity of the Au-catalyzed 1,2-H vs 1,2-Br migration is ligand dependent. The present study may provide better understanding of the reactions of allenes and alkynes activated by Au(I)

and Au(III) complexes,^{14,15} different catalytic performance of Au-complexes in different oxidation states, and particularly the ligand effect, which has received only limited attention.^{1c,8d,16}

Acknowledgment. This work was supported by Hundreds of Talents Program of CAS, Natural Science Foundation of Jiangsu Province (BK2005030), and NIH of the U.S. (Grant GM-64444). We thank Prof. Z.-X. Yu of Peking University for helpful discussion.

Supporting Information Available: Computational and experimental details. This material is available free of charge via the Internet at <http://pubs.acs.org>.

References

- (1) For recent reviews, see: (a) Hashmi, A. S. K.; Hutchings, G. J. *Angew. Chem., Int. Ed.* **2006**, *45*, 7896. (b) Zhang, L.; Sun, J.; Kozmin, S. A. *Adv. Synth. Catal.* **2006**, *348*, 2271. (c) Gorin, D. J.; Toste, F. D. *Nature* **2007**, *446*, 395. (d) Hashmi, A. S. K. *Chem. Rev.* **2007**, *107*, 3180.
- (2) (a) Sromek, A. W.; Rubina, M.; Gevorgyan, V. *J. Am. Chem. Soc.* **2005**, *127*, 10500. (b) Dudnik, A. S.; Sromek, A. W.; Rubina, M.; Kim, J. T.; Ke'lin, A. V.; Gevorgyan, V. *J. Am. Chem. Soc.* **2008**, *130*, 1440.
- (3) For higher oxophilicity of AuCl₃ compared to that of AuCl, see: Yamamoto, Y. *J. Org. Chem.* **2007**, *72*, 7817.
- (4) (a) Kirsch, S. F. *Org. Biomol. Chem.* **2006**, *4*, 2076. (b) Fürstner, A.; Davies, P. W. *Angew. Chem., Int. Ed.* **2007**, *46*, 3410.
- (5) Hashmi, A. S. K. *Angew. Chem., Int. Ed.* **2005**, *44*, 6990.
- (6) Ferrer, C.; Echavarren, A. M. *Angew. Chem., Int. Ed.* **2006**, *45*, 1105.
- (7) B3LYP/6-31G*(LANL2DZ for Au and Br) method was used for all the calculations, and solvation effect was calculated by CPCM model. See Supporting Information for details.
- (8) For selected examples of DFT studies in gold chemistry, see: (a) Nevado, C.; Echavarren, A. M. *Chem. Eur. J.* **2005**, *11*, 3155. (b) Comas-Vives, A.; González-Arellano, C.; Corma, A.; Iglesias, M.; Sánchez, F.; Ujaque, G. *J. Am. Chem. Soc.* **2006**, *128*, 4756. (c) Shi, F.-Q.; Li, X.; Xia, Y.; Zhang, L.; Yu, Z.-X. *J. Am. Chem. Soc.* **2007**, *129*, 15503. (d) Kovács, G.; Ujaque, G.; Lledós, A. *J. Am. Chem. Soc.* **2008**, *130*, 853. (e) Nieto-Oberhuber, C.; López, S.; Muñoz, M. P.; Cárdenas, D. J.; Buñuel, E.; Nevado, C.; Echavarren, A. M. *Angew. Chem., Int. Ed.* **2005**, *44*, 6146. (f) Faza, O. N.; López, C. S.; Álvarez, R.; de Lera, A. R. *J. Am. Chem. Soc.* **2006**, *128*, 2434. (g) Correa, A.; Marion, N.; Fensterbank, L.; Malacria, M.; Nolan, S. P.; Cavallo, L. *Angew. Chem., Int. Ed.* **2008**, *47*, 718.
- (9) Our computational studies confirmed possible formation of the originally proposed Au(III)-coordination complex **d** (Scheme 1);⁷ however, neither cyclization transition state nor halirenium intermediate **e** were located.
- (10) See Supporting Information for experimental and computational details.
- (11) Possible involvement of Brønsted acid as a true catalyst was ruled out by control experiments. Besides, addition of TTBP, a proton scavenger, did not inhibit the Au-catalyzed cycloisomerization.¹⁰
- (12) The Au-Cl distances in $\text{Au}(\text{PH}_3)\text{Cl}$, **5**, **6**, and **9** are 2.327, 2.481, 2.573, and 2.561 Å, respectively. See Supporting Information for bond critical point analysis of the weak interactions.
- (13) Possible involvement of AuCl as a true catalyst in AuCl_3 -catalyzed reactions via an in-situ redox process was ruled out by the observed different reactivity patterns of these complexes.¹⁰
- (14) For selected examples of allene activation by Au(I) and Au(III), see: (a) Hashmi, A. S. K.; Schwarz, L.; Choi, J.-H.; Frost, T. M. *Angew. Chem., Int. Ed.* **2000**, *39*, 2285. (b) Gockel, B.; Krause, N. *Org. Lett.* **2006**, *8*, 4485. (c) Morita, N.; Krause, N. *Eur. J. Org. Chem.* **2006**, 4634. (d) Zhou, C.-Y.; Chan, P. W. H.; Che, C.-M. *Org. Lett.* **2006**, *8*, 325. (e) Zhang, L. *J. Am. Chem. Soc.* **2005**, *127*, 16804. (f) Morita, N.; Krause, N. *Angew. Chem., Int. Ed.* **2006**, *45*, 1897. (g) Hoffmann-Röder, A.; Krause, N. *Org. Lett.* **2001**, *3*, 2537. (h) Morita, N.; Krause, N. *Org. Lett.* **2004**, *6*, 4121. (i) Kang, J.-E.; Lee, E.-S.; Park, S. I.; Shin, S. *Tetrahedron Lett.* **2005**, *46*, 7431. (j) Luzung, M. R.; Mauleon, P.; Toste, F. D. *J. Am. Chem. Soc.* **2007**, *129*, 12402. (k) Huang, X.; Zhang, L. *Org. Lett.* **2007**, *9*, 4627. (l) Dudnik, A. S.; Gevorgyan, V. *Angew. Chem., Int. Ed.* **2007**, *46*, 5195.
- (15) For examples of alkyne activation by gold proceeding via allenyl intermediates, see ref 2b and (a) Marion, N.; Nolan, S. P. *Angew. Chem., Int. Ed.* **2007**, *46*, 2750. (b) Zhang, L.; Wang, S. *J. Am. Chem. Soc.* **2006**, *128*, 1442. (c) Wang, S.; Zhang, L. *Org. Lett.* **2006**, *8*, 4585. (d) Wang, S.; Zhang, L. *J. Am. Chem. Soc.* **2006**, *128*, 8414. (e) Buzas, A.; Istrate, F.; Gagosz, F. *Org. Lett.* **2006**, *8*, 1957. (f) Buzas, A.; Gagosz, F. *J. Am. Chem. Soc.* **2006**, *128*, 12614.
- (16) (a) Markham, J. P.; Staben, S. T.; Toste, F. D. *J. Am. Chem. Soc.* **2005**, *127*, 9708. (b) Zhang, Z.; Liu, C.; Kinder, R. E.; Han, X.; Qian, H.; Widenhofer, R. A. *J. Am. Chem. Soc.* **2006**, *128*, 9066. (c) Marion, N.; Díez-González, S.; de Frémond, P.; Noble, A. R.; Nolan, S. P. *Angew. Chem., Int. Ed.* **2006**, *45*, 3647. (d) Sherry, B. D.; Toste, F. D. *J. Am. Chem. Soc.* **2004**, *126*, 15978.

JA802144T