


Evidence for Protein Radical-Mediated Nuclear Tunneling in Fatty Acid β -Oxygenase

Ankur Gupta, Arnab Mukherjee, Kenji Matsui, and Justine P. Roth

J. Am. Chem. Soc., **2008**, 130 (34), 11274-11275 • DOI: 10.1021/ja8042273 • Publication Date (Web): 02 August 2008

Downloaded from <http://pubs.acs.org> on February 8, 2009



More About This Article

Additional resources and features associated with this article are available within the HTML version:

- Supporting Information
- Access to high resolution figures
- Links to articles and content related to this article
- Copyright permission to reproduce figures and/or text from this article

[View the Full Text HTML](#)



ACS Publications
 High quality. High impact.


Evidence for Protein Radical-Mediated Nuclear Tunneling in Fatty Acid α -Oxygenase

Ankur Gupta,[†] Arnab Mukherjee,[†] Kenji Matsui,[‡] and Justine P. Roth*[†]

Department of Chemistry, Johns Hopkins University, 3400 North Charles Street, Baltimore Maryland 21218 and
Department of Biological Chemistry, Yamaguchi University, Yamaguchi 753-8515, Japan

Received June 4, 2008; E-mail: jproth@jhu.edu

The α oxidation of fatty acids is associated with essential metabolic, protective, and cell signaling functions in higher organisms.^{1,2} Here we provide evidence for an α oxidation mechanism involving a protein-derived radical. The reaction of interest is the first step in eq 1 where 2-*R*-hydroperoxide compounds are formed upon insertion of O₂ into fatty acid C _{α} -H bonds. In heme-containing dioxygenases,¹ such as rice α -oxygenase (R α O), the catalytic oxidant is proposed to be a tyrosyl radical.^{1c} This idea derives from (i) the homology of R α O to cyclooxygenases (COX-1 and COX-2) believed to use a Tyr \bullet to oxidize arachidonic acid to prostaglandin H³ and (ii) the loss of activity upon mutation of the conserved Tyr379 to Phe.^{1c,4-6}



We have found that treatment of R α O with H₂O₂ enhances the enzyme's activity and results in formation of a persistent organic radical.⁶ Experiments under anaerobic conditions suggest that the Tyr379 \bullet is thermodynamically capable of H \bullet abstraction from the fatty acid C _{α} -H. Large, weakly temperature dependent deuterium kinetic isotope effects (KIEs, ^Dk_{cat} and ^Dk_{cat}/K_M) are observed, consistent with nuclear tunneling. The description of such quantum effects poses an important challenge for current theories of proton-coupled electron transfer.^{7,8}

Though a crystal structure is not yet available,⁹ we have constructed a model of R α O using a COX-1 mutant with ~15% homology (Figure 1).¹⁰ Overlaying the R α O model structure with the monomeric subunit of wild-type (wt) COX-1 reveals similarly positioned His ligands to the Fe^{III} protoporphyrin IX (Fe^{III}Por) and virtually identical positions for the proposed catalytic Tyr residues (~6 Å from the porphyrin edge). In spite of overlapping Arg residues, the fatty acid is expected to bind differently in R α O with C _{α} -H pointing toward Tyr379 possibly due to an interaction between the carboxylate and His311.

Exposure of Fe^{III}-R α O to H₂O₂ results in oxidation of Tyr379 but not exogenous reductants;^{1c} this is a marked contrast to the peroxidase activity of COX.³ Reacting R α O (50 μ M) with H₂O₂ under N₂ results in an electron paramagnetic resonance (EPR) signal that persists for ≥ 1 h (Figure 2a). The spectral features^{6,12} at ambient and low temperatures are consistent with a Tyr \bullet in a yield of ~25% spin/heme after 20 min. The signal is diminished upon adding substrates such as decanoic or palmitic acids during preparation but not the inhibitor 5-(tetradecyloxy)-2-furoic acid (TOFA), which does not possess C _{α} -H bonds (Figure 2b).⁶ The putative substrate-derived radical is not detectable.

The Tyr379Phe mutant does not form a stable protein radical upon treatment with H₂O₂ (Figure 2c). The reactions of the wt and mutant R α O also differ spectrophotometrically (Figure 2d,e). Under conditions




Figure 1. Model of the active site of R α O (green, blue) compared to COX-1 (red) with bound arachidonate (yellow) (PDB: 1DIY).¹¹




Figure 2. X-band EPR spectra (22 °C) after 20 min (center = 3450 G): (a) wt-R α O (50 μ M) + 10 equiv of H₂O₂. (b) Sample (a) + 40 equiv of decanoate. (c) Tyr379Phe R α O (50 μ M) + 10 equiv of H₂O₂. UV-vis absorption spectra: (d) wt-R α O (7 μ M) + 10 equiv of H₂O₂ (initial = red, final = blue). (e) Tyr379Phe R α O (7.5 μ M) + 10 equiv of H₂O₂, (initial = red, intermediate = blue, final = green).

analogous to those of the EPR experiment, a red shift and slight bleach of the Soret band occur rapidly in wt-R α O.¹³ In contrast, Tyr379Phe R α O forms an intermediate resembling Fe^{IV}(O)Por⁺ which is reduced to Fe^{IV}(O)Por. Three Trp residues in the active site may provide the reducing equivalent(s).

Steady-state kinetics using an O₂ electrode reveals that palmitate and decanoate are fast and slow substrates, respectively, with K_M values that vary by $> 10^2$. Full enzyme activity was ensured by preincubating R α O with H₂O₂ (10–50 equiv) prior to the initial rate measurements. Experiments with peroxidases and reducing cosubstrates indicate that peroxide impurities present in the reaction mixtures are required for enzyme activity.^{3,6}

The kinetics of palmitate oxidation is only slightly affected by changes in pH: at pH 10, $k_{cat} = 18 \pm 1.5$ s⁻¹, $k_{cat}/K_M(\text{palmitate}) = (3.5 \pm 1.0) \times 10^6$ M⁻¹ s⁻¹ and $k_{cat}/K_M(\text{O}_2) = (4.0 \pm 0.5) \times 10^5$ M⁻¹ s⁻¹; at pH 7.2, $k_{cat} = 9.0 \pm 0.5$ s⁻¹, $k_{cat}/K_M(\text{palmitate}) = (2.2 \pm 0.4) \times 10^6$ M⁻¹ s⁻¹ and $k_{cat}/K_M(\text{O}_2) = (3.0 \pm 0.5) \times 10^5$ M⁻¹ s⁻¹. Errors are reported throughout as $\pm 1\sigma$. Experiments were mostly conducted at the higher pH where enzyme activity and substrate solubility are enhanced, while kinetic complexity appears to be minimized (see below). Under the experimental conditions, the 2-*R*-

[†] Johns Hopkins University.

[‡] Yamaguchi University.




Figure 3. Determination of Dk_{cat}/K_M by GC/MS analysis. The resolution of d_{29} and h_{29} pentadecanal at (a) 100% and (b) \sim 1% product formation.




Figure 4. Temperature dependence of k_{cat} for h_{31} - and d_{31} -palmitate at pH 10 (a) and pH 7.2 (b). Errors are shown for visibility as $\pm 2\sigma$.

hydroperoxide products are unstable^{1c} losing CO_2 to afford $\text{C}_{\text{N}-1}$ aldehydes which were identified and quantified using gas chromatography/mass spectrometry (GC/MS).

Palmitate oxidation is characterized by a large competitive KIE at pH 10. Experiments were performed by analyzing the product ratios from solutions initially containing 5:1 or 10:1 mixtures of h_{31} : d_{31} -palmitate (Figure 3).¹⁴ Five independent determinations at $<5\%$ reaction conversion indicated Dk_{cat}/K_M (22 °C) = 53 ± 5 .

Large KIEs were also determined uncompetitively at pH 10 with saturating ($\geq 10K_M$) concentrations of O_2 and palmitate ($>98\%$ isotopic purity).⁶ Identical k_{cat} values were observed in H_2O and D_2O indicating the absence of isotope exchange into the α -position. Dk_{cat} (palmitate) = 54 ± 7 (22 °C) was determined using α,α - d_2 -palmitate or d_{31} -palmitate. None of the experiments showed a kinetic burst due to protio substrate contamination or an induction phase due to insufficient peroxide initiator.⁶ The similar KIEs, Dk_{cat} and Dk_{cat}/K_M , imply that C_{α} -H cleavage is the first irreversible as well as the turnover-controlling step at pH 10. At pH 7.2 and 22 °C, Dk_{cat} is diminished to 31 ± 5 . This is presumably the result of a downstream unimolecular step that contributes to k_{cat} , reducing the KIE from the intrinsic value.

The temperature dependence of the intrinsic KIE reveals the quantum mechanical nature of C-H oxidation as discussed by Klinman et al.¹⁵ Arrhenius plots reveal $E_a(\text{H}) = 9.3 \pm 0.2 \text{ kcal mol}^{-1}$ and $A(\text{H}) = (1.3 \pm 0.4) \times 10^8 \text{ s}^{-1}$, $E_a(\text{D}) = 10.4 \pm 0.2 \text{ kcal mol}^{-1}$ and $A(\text{D}) = (1.3 \pm 0.3) \times 10^7 \text{ s}^{-1}$ at pH 10; $E_a(\text{H}) = 12.6 \pm 0.3 \text{ kcal mol}^{-1}$, $A(\text{H}) = (2.1 \pm 0.9) \times 10^{10} \text{ s}^{-1}$, $E_a(\text{D}) = 12.9 \pm 0.3 \text{ kcal mol}^{-1}$ and $A(\text{D}) = (1.1 \pm 0.6) \times 10^9 \text{ s}^{-1}$ at pH 7.2 (Figure 4). The intrinsic KIE is more fully expressed at pH 10 where $E_a(\text{D}) - E_a(\text{H}) = 1.1 \pm 0.3 \text{ kcal mol}^{-1}$ and $A(\text{H})/A(\text{D}) = 10 \pm 4$. The $A(\text{H})/A(\text{D})$ deviates from 1 within the $\pm 1\sigma$ error weighted limits, as seen in some other systems.^{6,15c,d}

This study has demonstrated large deuterium KIEs upon C-H oxidation by $\text{R}\alpha\text{O}$ and supported the intermediacy of the $\text{Tyr}379\bullet$ in catalysis. Though $\text{Fe}^{\text{IV}}(\text{O})\text{Por}\bullet+$ and $\text{Fe}^{\text{IV}}(\text{O})\text{Por}$ were observed, a persistent radical was absent in the $\text{Tyr}379\text{Phe}$ mutant for which no activity could be detected in solutions containing up to $5 \mu\text{M}$ protein. Future efforts will concentrate on the kinetic mechanism. At this stage, we cannot rigorously exclude the possibility that the $\text{Tyr}379\bullet$ is a side product or the intermediacy of another catalytic oxidant, possibly an unstable peroxy radical. A reversible reaction with O_2 to form an amino acid peroxy radical that is regenerated with each enzyme turnover would still be consistent with the stoichiometry of eq 1.

Consideration of such a species is warranted in view of the thermodynamics; the $\text{H}\bullet$ affinity of a $\text{Tyr}\bullet$ is expected to be less than the inverse bond dissociation energy of the fatty acid $\text{C}_{\alpha}\text{-H}$ ($\sim 90 \text{ kcal mol}^{-1}$) making initial $\text{H}\bullet$ abstraction endothermic.¹⁶

Though increasingly observed in enzymes that oxidize substrate C-H bonds,¹⁵ the large KIEs in $\text{R}\alpha\text{O}$ are somewhat surprising. The analogous reaction in the homologous COX-1 is characterized by a small tritium KIE of ~ 4 .³ Kinetically reversible $\text{H}\bullet$ abstraction by the $\text{Tyr}\bullet$ has been proposed in COX-1,¹⁷ a reaction that involves oxidation of a weak, bis-allylic C-H with a dissociation energy of $\sim 80 \text{ kcal mol}^{-1}$. The irreversible reaction in $\text{R}\alpha\text{O}$ suggests that the oxidant's thermodynamic affinity for $\text{H}\bullet$ may be significantly altered relative to COX.

$\text{R}\alpha\text{O}$, thus, presents a novel example where a protein-derived radical may effect homolysis of a robust C-H bond by nuclear tunneling.⁸ Although a close distance between $\text{H}\bullet$ donor and acceptor is anticipated for radical reactions,^{7c} the KIEs from 40 to 60 (5–40 °C) reveal nonadiabatic behavior due to vibrational overlap below the activation barrier. In view of these unique results, it will be interesting to determine how current theories of proton-coupled electron transfer are able to account for the isotopic activation parameters seen in $\text{R}\alpha\text{O}$.⁷

Acknowledgment. Support was provided by NSF CHE-044990, Research Corp. Cottrell Scholar, Alfred P. Sloan Fellowship, and Camille Dreyfus Teacher Scholar awards to J.P.R. We thank Prof. C. A. Townsend for the gift of TOFA and Dr. T. Koeduka and Mr. K.-M. Chang for assistance with experiments.

Supporting Information Available: Experimental details. This material is available free of charge via the Internet at <http://pubs.acs.org>.

References

- (1) (a) Hamberg, M.; Ponce de Leon, I.; Rodriguez, M. J.; Castresana, C. *Biochem. Biophys. Res. Commun.* **2005**, *338*, 169. (b) Liu, W.; Wang, L.-H.; Fabian, P.; Hayashi, Y.; McGinley, C. M.; van der Donk, W. A.; Kulmacz, R. J. *Plant. Phys. Biochem.* **2006**, *44*, 284. (c) Koeduka, T.; Matsui, K.; Akakabe, Y.; Kajiwara, T. *J. Biol. Chem.* **2002**, *277*, 22648. (d) Su, C.; Sahlin, M.; Oliw, E. H. *J. Biol. Chem.* **1998**, *273*, 20744.
- (2) Jansen, G. A.; Wanders, R. J. A. *Biochem. Biophys. Acta* **2006**, *1763*, 1403.
- (3) Rouzer, C. A.; Marnett, L. J. *Chem. Rev.* **2003**, *103*, 2239.
- (4) An error in the wt sequence has since been corrected (Genbank#AF229813).
- (5) The His-tagged heme-containing protein was expressed as in ref 1c. The enzyme concentration was determined from $\epsilon_{410} = 123 \text{ mM}^{-1} \text{ cm}^{-1}$.
- (6) Further details are provided in the Supporting Information.
- (7) (a) Kuznetsov, A. M.; Ulstrup, J. *Can. J. Chem.* **1999**, *77*, 1085. (b) Hammes-Schiffer, S. *ChemPhysChem* **2002**, *3*, 33–42. (c) Cukier, R. I. *J. Phys. Chem. B* **2002**, *106*, 1746.
- (8) For other examples, see: (a) Whittaker, M. M.; Ballou, D. P.; Whittaker, J. W. *Biochemistry* **1998**, *37*, 8426. (b) Chowdhury, S.; Banerjee, R. *J. Am. Chem. Soc.* **2000**, *122*, 5417.
- (9) Lloyd, T.; Krol, A.; Campanaro, D.; Malkowski, M. *Acta Crystallogr., Sect. F* **2006**, *62*, 365.
- (10) Constructed using PHYRE: www.sbg.bio.ic.ac.uk/phyre/index.cgi and the COX-1 mutant (PDB: 1U67); Fe^{III} protoporphyrin IX ($\text{Fe}^{\text{III}}\text{Por}$) is replaced by $\text{Co}^{\text{III}}\text{Por}$ in the crystallized protein; Harman, C. A.; Rieke, C. J.; Garavito, R. M.; Smith, W. L. *J. Biol. Chem.* **2004**, *279*, 42929.
- (11) Malkowski, M. G.; Ginell, S. L.; Smith, W. L.; Garavito, R. M. *Science* **2000**, *289*, 1933.
- (12) (a) Østdal, H.; Andersen, H. J.; Davies, M. J. *Arch. Biochem. Biophys.* **1999**, *362*, 105. (b) Svistunenko, D. A. *Biochim. Biophys. Acta* **2005**, *1707*, 127. (c) Davies, M. J.; Puppo, A. *Biochem. J.* **1992**, *281*, 197.
- (13) Similar behavior has been reported for ascorbate peroxidase: Hiner, A. N. P.; Martinez, J. I.; Arnao, M. B.; Acosta, M.; Turner, D. D.; Raven, E. L.; Rodriguez-Lopez, J. N. *Eur. J. Biochem.* **2001**, *268*, 3091.
- (14) Similar methodology using HPLC is described in: Lewis, E. R.; Johansen, E.; Holman, T. R. *J. Am. Chem. Soc.* **1999**, *121*, 1395.
- (15) (a) Klinman, J. P. *Philos. Trans. R. Soc. London, Ser. B* **2006**, *361*, 1323. (b) Klinman, J. P. *Pure Appl. Chem.* **2003**, *75*, 601. (c) Meyer, M. P.; Tomchick, D. R.; Klinman, J. P. *Proc. Natl. Acad. Sci. U.S.A.* **2008**, *105*, 1146. (d) Knapp, M. J.; Rickert, K.; Klinman, J. P. *J. Am. Chem. Soc.* **2002**, *124*, 3865.
- (16) Estimated using acetate as a model: (a) Wenthold, P. G.; Squires, R. R. *J. Am. Chem. Soc.* **1994**, *116*, 11890. The Tyr O-H bond strength is estimated to be $\sim 85 \text{ kcal mol}^{-1}$. (b) Blomberg, M. R. A.; Siegbahn, P. E. M.; Styring, S.; Babcock, G. T.; Aakermark, B.; Korall, P. *J. Am. Chem. Soc.* **1997**, *119*, 8285.
- (17) Mukherjee, A.; Brinkley, D. W.; Chang, K.-M.; Roth, J. P. *Biochemistry* **2007**, *46*, 3975.

JA8042273